



Effects of endocrine disrupting chemicals on gonad development: Mechanistic insights from fish and mammals

G. Delbes^{a,*}, M. Blázquez^b, J.I. Fernandino^c, P. Grigorova^d, B.F. Hales^e, C. Metcalfe^f, L. Navarro-Martín^g, L. Parent^d, B. Robaire^{e,h}, A. Rwigemera^a, G. Van Der Kraakⁱ, M. Wade^j, V. Marlatt^k

^a Centre Armand Frappier Santé Biotechnologie, Institut National de la Recherche Scientifique (INRS), Laval, Canada

^b Institute of Marine Sciences (ICM-CSIC), Barcelona, Spain

^c Instituto Tecnológico de Chascomús (CONICET-UNSAM), Chascomús, Argentina

^d Université TELUQ, Montréal, Canada

^e Department of Pharmacology and Therapeutics, McGill University, Montreal, Canada

^f School of Environment, Trent University, Trent, Canada

^g Institute of Environmental Assessment and Water Research (IDAEA-CSIC), Barcelona, Spain

^h Department of Obstetrics and Gynecology, McGill University, Montreal, Canada

ⁱ Department of Integrative Biology, University of Guelph, Guelph, Canada

^j Environmental Health Science & Research Bureau, Health Canada, Ottawa, Canada

^k Department of Biological Sciences, Simon Fraser University, Burnaby, Canada

ARTICLE INFO

Keywords:

Development
Endocrine disrupting chemicals
Fish
Gonadal differentiation
Mammals
Ovary
Sex determination
Testis

ABSTRACT

Over the past century, evidence has emerged that endocrine disrupting chemicals (EDCs) have an impact on reproductive health. An increased frequency of reproductive disorders has been observed worldwide in both wildlife and humans that is correlated with accidental exposures to EDCs and their increased production. Epidemiological and experimental studies have highlighted the consequences of early exposures and the existence of key windows of sensitivity during development. Such early in life exposures can have an immediate impact on gonadal and reproductive tract development, as well as on long-term reproductive health in both males and females. Traditionally, EDCs were thought to exert their effects by modifying the endocrine pathways controlling reproduction. Advances in knowledge of the mechanisms regulating sex determination, differentiation and gonadal development in fish and rodents have led to a better understanding of the molecular mechanisms underlying the effects of early exposure to EDCs on reproduction. In this manuscript, we review the key developmental stages sensitive to EDCs and the state of knowledge on the mechanisms by which model EDCs affect these processes, based on the roadmap of gonad development specific to fish and mammals.

Introduction

Concerns have arisen regarding the impact of endocrine disrupting chemicals (EDCs) on reproductive health. There is widespread exposure to populations when these chemicals, found in a large variety of consumer products, including electronic equipment, building materials, cosmetics, medical devices and food packaging, leach out into the environment (Metcalfe et al., this issue). Many of these chemicals are used as plasticizers, flame retardants, or pesticides, while others are pharmaceuticals or industrial by-products. Correlating with accidental

exposures to EDCs and the rise in their production, an increased frequency of reproductive disorders has been observed worldwide in both wildlife and humans (Marlatt et al. this issue; WHO, 2002). Such disorders include feminization in fish, abnormal development of the reproductive tract, precocious puberty, testicular and ovarian cancer, low sperm count and poor gamete quality (Marlatt et al. this issue). A growing number of epidemiological and wildlife studies have highlighted the impact of early exposures and the existence of key windows of sensitivity during development. This is in line with the concept of the developmental origins of health and diseases (DOHaD) that early

* Corresponding author. Institut national de la recherche scientifique, Centre Armand-Frappier Santé Biotechnologie, 531 boulevard des Prairies, Laval (Quebec), H7V 1B7, Canada.

E-mail address: geraldine.delbes@inrs.ca (G. Delbes).

<https://doi.org/10.1016/j.envres.2021.112040>

Received 27 May 2021; Received in revised form 3 September 2021; Accepted 8 September 2021

Available online 9 September 2021

0013-9351/© 2021 The Authors.

Published by Elsevier Inc.

This is an open access article under the CC BY-NC-ND license

(<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

perinatal exposures to environmental influences affect long-term health and disease susceptibility (Gillman, 2005). According to this principle, immediate effects may be observed shortly after exposure, but it is also possible that the phenotypic consequences of molecular effects on the programming of a tissue may be visible only later, in the adult. In fact, it has been shown that, depending on the type of EDCs, the dose and the precise time of exposure, early exposure to EDCs can induce immediate effects on the establishment of the different cell types of the gonads or on the development of the reproductive tract; later effects may be revealed on hormonal homeostasis, somatic cell differentiation, gamete production and gamete quality (Johansson et al., 2017; Wohlfahrt-Veje et al., 2009).

EDCs exert their effects by altering endocrine pathways, including those that control reproduction. Traditionally, the mechanisms of action of EDCs are defined as affecting receptor binding, synthesis or transport of hormones. Thus, many studies are focussed on studying activation/inhibition of estrogen, androgen, and thyroid receptor signaling and effects on steroidogenesis (EATS-pathways) (Martyniuk et al. this issue; Robitaille et al. this issue; Amir et al., 2021). However, non-EATS pathways are now gaining importance and interest as a modality by which EDCs impact reproductive health (Martyniuk et al. this issue). A better understanding of the underlying physiology is required to anticipate the potential effects of environmental chemicals such as EDCs. In recent years, fundamental studies, primarily in fish and rodents, have led to a better understanding of sex determination, gonadal differentiation, and the mechanisms underlying the events during the embryo and early life stages that are critical to the development of the reproductive system and germ cell lineage. These advances in knowledge increase our understanding of the molecular mechanisms by which EDCs affect key events and induce long-term effects on reproduction.

In this manuscript, we will review the key events in the establishment of the gonads and the state of knowledge on the mechanisms by which EDCs affect these processes. This will be based on the roadmaps of development specific to fish and mammals, from gonad development and differentiation to the entry of germ cells into meiosis. The selection of EDCs reported here, is based on the prevalence of exposure in the study model and known reproductive effects; our focus is on EDCs for which we have some understanding of the mechanisms involved in gonadal dysregulation during development.

Part 1: evidence from Fish

An increasing number of chemicals coming from wastewater treatment plants, leaching from agricultural lands or present in industrial effluents or in urban runoff, among others, are contaminating aquatic ecosystems and producing adverse effects on fish populations (reviewed by Segner, 2011). In the past decades, the effects of EDCs on reproductive function in several fish species have been the focus of an increasing number of investigations (Marlatt et al. this issue). But, only a few studies have addressed the effect of EDCs in fish prior to pubertal onset and little is known about their precise mode of action. Yet, evidence demonstrates that in fish, exposure to EDCs is responsible for: 1) the disruption of gonad development and sex differentiation, inducing intersex; 2) abnormal gonad differentiation, affecting the number of germ cells, leading to episodes of sterility; and, 3) alterations in the timing of puberty. Here we summarize the physiology of these key stages of development and review how early exposures to EDCs affect fish reproduction from sex determination to entry into puberty (Figs. 1 and 2, Table 1).

1. Gonad development

1.1. Development of germ cells

The establishment of the germ line is a crucial process for the reproductive success of an individual. This includes several well-orchestrated steps, beginning with specification of the germ cell identity in the early stages of development (Ewen-Campen et al., 2010). These germ cell precursors of gametes, named primordial germ cells (PGCs), migrate actively through the body via the coelom, reaching their final location, the gonadal primordium (Nishimura and Tanaka, 2016; Weidinger et al., 2002). This primordium presents an exclusive characteristic in terms of organ development because it differentiates into an ovary or a testis, two dissimilar organs from the same origin (Capel, 2017). When PGCs reach the gonadal primordium, they are named embryonic germline stem cells (EGSCs) and are arrested at G1/G0 phase of the cell cycle. These germ cells become the precursors of spermatogonial stem cells (SSCs) in the testis and oogonial stem cells (OSCs) in the ovary (Nakamura et al., 2010). During migration and even after reaching the gonadal primordium, some of these EGSCs will start asynchronous proliferation in both sexes to give rise to the primary population of germ cells in the gonads (Chen and Liu, 2015c; Nakamura




Fig. 1. Timing of testicular development in zebrafish, rodents and humans. The diagram illustrates the fate and development of germ cells in the testis, as they proliferate by mitosis, and organize to form tubules (cysts in zebrafish) and then enter spermatogenesis.




Fig. 2. Timing of ovarian development in zebrafish, rodents and humans. The diagram illustrates the fate and development of germ cells in the ovary as they proliferate by mitosis, organize to form ovarian follicles, and ultimately undergo folliculogenesis. One major difference between fish versus mammals is that fish maintain a population of germ stem cells in the adult ovary whereas in mammals at birth all germ cells are arrested in meiotic prophase, forming primordial follicles, and are incapable of dividing mitotically subsequently.

et al., 2010; Tanaka, 2019), while others remain in a quiescent state. This first phase of germline proliferation is regulated by the Anti-Müllerian hormone (Amh), as described by Morinaga et al. (2007). Any alterations of this proliferation and of the number of EGSCs are involved in gonadal sex reversal. For example, in Japanese medaka (*Oryzias latipes*) hyperproliferation of EGSCs induces male-to-female reversal, while in zebrafish (*Danio rerio*), the lack of this proliferation induces the female-to-male sex reversal (Kurokawa et al., 2007; Rodríguez-Marí et al., 2010). Before meiosis, EGSCs enter a second round of proliferation, this time synchronous, which generates a group of cells known as cysts or nests, surrounded by somatic cells (Saito et al., 2007, Figs. 1 and 2). Timing of when this second proliferation begins is specific to sex; mostly, it begins first in females and is an initial marker of sexual dimorphism (Tanaka, 2019). Although this second round of proliferation is not related to the establishment of gonadal sex, the early increase in the number of germ cells is a necessary step for reproductive success, as changes can alter the number of eggs in females at spawning, and the fertilization rate and mating behavior of males (Padilla et al., 2021).

1.2. Regulation and impacts of EDCs

The possibility that PGCs are a target of EDC action has been studied in zebrafish embryos (Hu et al., 2014; Lombó et al., 2019). Embryos were injected with the gfp-nanos-3'UTR mRNA and were immediately exposed to different concentrations of the potent synthetic estrogen 17 α -ethynylestradiol (EE2). The migration and distribution of PGCs were affected by exposure to high doses of EE2 (500 ng/L) as the embryos displayed ectopic PGCs. Knocking down the estrogen receptor *esr2a* or application of a ESR2 inhibitor significantly reduced the percentage of embryos with ectopic PGCs after exposure to EE2, indicating that ESR2a may play an important role in EE2-related PGC phenotypes. In other studies, zebrafish embryos were exposed to Bisphenol A (BPA) during the first 24 h of development. Vasa immunostaining of PGCs revealed that exposure to BPA impaired their migration to the genital ridge and two pivotal genes of PGCs migration (C-X-C Motif Chemokine

Receptor 4 (*cxc4b*) and C-X-C Motif Chemokine Ligand 12 (*sdf1a*) were highly dysregulated (Lombó et al., 2019). Interestingly, once the embryos reached adulthood normal testes were developed. Collectively, these studies show that PGCs may be targeted by EDCs in fish.

2. Sex-determination and gonadal differentiation

2.1. Control of sex determination and gonadal differentiation

Fish are a group of vertebrates with a large range of mechanisms for sex determination, including genetic and environmental sex determination mechanisms (GSD and ESD, respectively; Devlin and Nagahama, 2002; Hattori et al., 2020). In species with GSD, sex is determined at fertilization, driven by a gene or chromosomal difference between sexes. Different heterogametic systems have been characterized in fish. However, the genes that initiate the development of males (XX/XY system) or females (ZZ/ZW system) vary. For example, male heterogametic systems have been identified that exhibit different sex determining genes in closely related families (Hattori et al., 2019; Maitre et al., 2017; Matsuda, 2005). Moreover, a similar range has been observed in female heterogametic systems. Many closely related species exhibit both types of heterogamety (Capel, 2017), showing that genes that direct gonadal fate have appeared several times, independently, in fish. In ESD strategies, gonadal fate is determined at embryo or juvenile stages, driven by the effects of changes in the environment, such as temperature, dissolved oxygen or pH (Hattori et al., 2020).

Interestingly, GSD and ESD can coexist under environmentally relevant conditions, exhibiting a continuum of genetic and environmental mechanisms (Xiong et al., 2020; Yamamoto et al., 2014). As an example, it was recently established that the brain, through the hypothalamus, detects changes in the environment by raising cortisol levels (Castañeda Cortés et al., 2019). This stress hormone produces, as a by-product, changes to the ratio of sex steroids. On one hand, it can increase androgen levels by the up-regulation of the *hsd11b2* gene that encodes for the 11 β -hydroxysteroid dehydrogenase type 2 enzyme

Table 1
Impacts of early exposure to EDCs on gonad development and puberty in fish.

Species	Exposure Window	Compounds	Gonadal effects	Endocrine and Molecular effects	References	
Zebrafish (<i>Danio rerio</i>)	1 hpf - 60dph	EE2	Delayed puberty in both males and females	Increase in VTG in both males and females	Baumann et al. (2014), 2013	
		4-Tert-Pentylphenol	Delayed puberty in both males and females	Decrease in VTG levels in females, increase in males		
		DHT	Delayed puberty in females and acceleration in males	Decrease in VTG in both males and females		
		Tb	Delayed puberty in females and acceleration in males	Decrease in VTG in both males and females		
	2 hpf-56 dph 1 - 60dph	EE2	Prochloraz	Hormesis-like effects with no clear results	Hormesis-like effects with increased VTG levels at intermediate concentrations	Schulz et al. (2007) Örn et al. (2006)
			EE2	Decreased testicular development	Decreased amh and decreased dmrt1	
		Tb	Precocious puberty in females, determined by ovaries showing only pre-vitellogenic oocytes	Increased VTG levels at 38 dph		
	1 dpf - 60dph	Tb	Precocious puberty in males, determined by increased testicular area and increased area with spermatozoa in the testis lumen	Decreased VTG levels at 38 dph	Morthorst et al. (2010)	
	6 dpf - 5 months	Natural mixture of POPs	Precocious puberty, determined by treated fish taking less time to first spawning	No effects on VTG levels at 60 dph	Nourizadeh-Lillabadi et al. (2009)	
	2 - 60dph	NP + EE2	Precocious puberty, determined by acceleration of spermatogenesis	Transcriptomic alterations in testis and liver related to steroid and thyroid hormone functions, insulin signaling and metabolic homeostasis.	Lin and Janz (2006)	
	20 - 80 dpf	Levonorgestrel	Progesterone	Precocious puberty, determined by advanced gonadal maturation in males but not in females	No differences in HSP70 expression, increasing NP concentration acted antagonistically to EE2 in terms of VTG induction at 60 dph	Svensson et al. (2016)
			EE2	Sex reversal (all-male populations). Precocious puberty, determined by advanced gonadal maturation in males but not in females	Increase in the expression of <i>amh</i> , <i>cyp11b</i> , <i>fshb</i> , and <i>lhb</i> and decrease of <i>cyp19a1a</i> at 44 dpf	
20-115 dpf	DBP + EE2	EE2, DHP and the combination decreased testicular development (Reduced spermatozoa and increased spermatogonia and spermatocytes)	No effects on gene expression levels	Chen et al. (2015b)		
1-120dpf	triadimefon	EE2, DHP the combination reduced ovariaian development through reduction in vitellogenic oocytes and degenerative changes	Increased vtg protein at 45 d No effect on vtg protein at 45 d Increased vtg protein at 45d			
1-60dpf	prochloraz	EE2 and EE2 plus DHP induced 100% females whereas DHP had no effect on sex ratio	Reduced vtg expression but no effects on testosterone or estradiol levels	Liu et al. (2014)		
1-60dpf	prochloraz	Sex ratio of fish skewed to male and female exposed to 0.5 g/mL triadimefon had immature ovary. No obvious effects on males. Decreased spawning and fertility success	Decreased vtg in females; in males low levels of prochloraz increased vtg in males whereas high levels decreased vtg	Kinnberg et al. (2007)		
1-60dpf	prochloraz	Increased numbers of males and a higher proportion of intersex fish	Decreased vtg in females; in males low levels of prochloraz increased vtg in males whereas high levels decreased vtg	Song et al. (2020)		
Wild-type and <i>cyp19a1a</i> and nER mutant zebrafish (<i>Danio rerio</i>)	8 - 9wpf	DES	Increased numbers of males and a higher proportion of intersex fish	Decreased vtg in females; in males low levels of prochloraz increased vtg in males whereas high levels decreased vtg	Lessman and Brantley (2020)	
Casper mutant transparent zebrafish (<i>Danio rerio</i>)	1 - 100 dph	Genistein	Delayed puberty, determined by delayed oocyte maturation, atresia, reduced oocyte number, larger ovarian lumen and higher incidence of PGCs in females. No effects on		Kiparissis et al. (2003)	

(continued on next page)

Table 1 (continued)

Species	Exposure Window	Compounds	Gonadal effects	Endocrine and Molecular effects	References
	1 dph - 60dph	Equol	spermatogenesis but increase in testis connective tissue, lobular fibrosis and decreased spermatozoa densities Delayed puberty in females, determined by delayed oocyte maturation, atresia, larger ovarian lumen and development of somatic stromal tissue in females. Delayed puberty in males, determined by retarded spermatogenesis EE2	Precocious puberty, determined by higher proportion early vitellogenic oocytes	Increased VTG levels at 38 dph
Tb	Precocious puberty, determined by increased area with spermatozoa in the testis lumen	Örn et al. (2006)	Decreased VTG levels at 38 dph		
Rainbow trout (<i>Oncorhynchus mykiss</i>)	6 month to 18 mo (first gametogenesis and spawning)	Genistein	Delayed puberty in females, determined by delays in oocyte maturation. Acceleration of spermatogenesis and decrease in sperm density and motility	Induction of VTG, decrease in, 17a,20b-dihydroxyprogesterone, testosterone and <i>fshb</i> and <i>lhb</i> in both females and males.	Bennetau-Pelissero et al. (2001)
Roach (<i>Rutilus rutilus</i>)	1 year-old- 28 days exposure (first gametogenesis)	Levonorgestrel	No effect in ovaries Decrease in numbers of spermatogonia type B in testis	Increased <i>vtg</i> and <i>er1</i> expression in both sexes Increased <i>lhb</i> and suppression of <i>fshb</i> expression in both sexes Decrease of 11 KT and E2 and increase of T in females Decrease 11 KT in males	Kroupova et al. (2014)
White suckers (<i>Catostomus commersoni</i>)		Pulp mill effluents	Delayed puberty, determined by increased age to maturity	Decreased testosterone, 17a,20b-dihydroxyprogesterone in both sexes, 11-KT in males and E2 in females.	McMaster et al. (1991); Munkittrick et al. (1992)

11 KT, 11ketotestosterone; Amh, antimüllerian hormone, BPA, Bisphenol A; cyp11b, cytochrome P450 11 subfamily B; cyp19a1a, cytochrome P450 19 subfamily A member 1a (aromatase); DES, Diethylstilbestrol; dph, days post hatch, dpf, days post fertilization; DHT, Dihydrotestosterone; E2, 17 β estradiol; EE2, 17 α -Ethinylestradiol; fshb, Follicle stimulating hormone subunit beta; hpf, hours post fertilization; lhb, luteinizing hormone subunit beta; NP, Nonylphenol; POP, persistent organic pollutants; Tb, 17b-Trenbolone; VTG, vitellogenin; wpf, week post fertilization.

(Note that examples of studies are presented here; the list is not comprehensive).

(11 β -HSD2) involved in the synthesis of 11-oxygenated androgen, which is the most potent androgen in fish (Castañeda-Cortés et al., 2020). On the other hand, it affects estrogen levels by the down-regulation of cytochrome P450 family 19 subfamily A member 1 (*cyp19a1a*), which is the gene encoding the aromatase enzyme involved in the biosynthesis of estradiol (Yamaguchi et al., 2010). This complex interaction between the environment, stress hormones and alterations of gene expression induces the development of testis.

The diversity of sex determination mechanisms, and subsequent gonadal differentiation, is possible in fish due to a high gonadal plasticity and to the similarity of extremely simple structures in both sexes - mainly the germline cells surrounded by somatic supporting-cells (Nishimura et al., 2016). Although the triggers of gonadal development vary enormously between fish, the molecular networks that promote the development of testes or ovaries are highly conserved, even with mammals (Capel, 2017). Depending on the way in which the differentiation of the gonads progresses, they are classified into two reproductive strategies, gonochorism and hermaphroditism (Devlin and Nagahama, 2002).

Gonochoristic species differentiate as males or females and remain the same sex throughout their lifespan, while hermaphroditic species develop both the male and female phenotype at some point in their life history (Devlin and Nagahama, 2002). In the gonochoristic group, the gonadal primordium activates the gonadal differentiation network, based on the sex determination system, establishing a balance between testis- and ovary-related genes. Within these genes, which include several transcription and growth factors, the male-related genes, such as *amh*, mab3-related transcription factor 1 (*dmrt1*), sry-box containing gene 3 (*sox3*) and gonadal soma derived factor (*gsdf*) are up-regulated in the bipotential gonadal primordium and during gonadal differentiation

to induce and maintain the development of testis. Alternately, the female-related genes, such as R-spondin 1 (*rspo1*), forkhead box protein L2 and L3 (*foxl2*, *foxl3*) and cytochrome P450 1A1 (*cyp11a1*), are up-regulated to induce the development of ovaries (Guiguen et al., 2018; Nishimura et al., 2015). Moreover, other hormones, receptors and enzymes that regulate gonadal steroidogenesis, such as follicle-stimulating hormone receptor (*fshr*), Cytochrome P450 Family 11 Subfamily B Member 1 (*cyp11b1*) and *hsd11b2*, are expressed downstream within the gonadal gene regulatory network to initiate and/or maintain testis development. The female-related gene *cyp19a1a* is necessary for ovary development and/or maintenance (Guiguen et al., 2018; Hattori et al., 2020). Once the gonad has differentiated into a testis or ovary, the gonadal fate in gonochoristic fish species cannot subsequently be reversed under normal conditions. However, in adult gonochoristic fish, intersex gonads have been observed upon hormonal manipulation; that is, ovaries that had varying amounts of male germline cells or testes that had varying amounts of primary oocytes (Devlin and Nagahama, 2002), establishing that the lability of the gonadal somatic and germline cells persists even in adulthood. For example, the transplantation of germ stem cells, such as SSCs or OSCs, into a host of the opposite sex results in the generation of oocytes or spermatocytes, respectively (Yoshizaki and Yazawa, 2019).

Gonadal differentiation in fish with hermaphrodite strategies can shift from one to another gonadal biological sex, depending mainly on social factors (Gemmell et al., 2019). Hermaphrodites that develop sequentially are either protandrous, if they first mature as males, or protogynous, if they first mature as females. This plasticity can occur in fish because, unlike mammals, gonads contain germ cells as stem cells during adulthood in both sexes (Nakamura et al., 2010). Like in gonochoristic fish, a conserved gonadal gene regulatory network has been

characterized in hermaphroditic fish during sex change (Ortega-Recalde et al., 2020). However, it is unusual among teleosts for the same gonad to display stages of both spermatogenesis and oogenesis; when this is observed, it is usually referred to as gonadal “intersex”, or sometimes “testis-ova”, or “ovotestes” among gonochoristic or hermaphroditic teleost (see Marlatt et al. this issue, Fig. 2).

2.2. Impact of EDCs

Intersex and other alterations to gonadal development have been observed in several populations of wild fish, and this is often viewed as evidence of exposure to EDCs (Metcalfe et al., 2010 and Marlatt et al. this issue). In a review of the literature on gonadal intersex in populations of teleosts in the wild, Bahamonde et al. (2013) tabulated reports of this condition in 37 fish species from field surveys conducted in 24 countries. However, there are questions concerning whether this condition is definitively linked to exposures of wild fish to EDCs and whether a “background” prevalence of gonadal intersex is a natural condition in some species of gonochoristic teleosts (Bahamonde et al., 2013).

With the high fecundity and the availability of large numbers of fertilized eggs and larvae, a common approach for studies of the effects of EDCs and environmental contaminants on reproductive development in fish involves their treatment, grow out and evaluation of subsequent effects on sexual development. Many studies have reported gender bias and effects on gonad developmental in fish treated with EDCs. Santos et al. (2017) summarized the studies testing the effects of more than 25 suspected EDCs on sex development and the sex ratio of zebrafish. These studies showed that treatment with compounds that mimic the action of estrogens [e.g. EE2, BPA] result in a shift in the sex ratio towards females. Exposure to androgenic EDCs, such as 17 β -trenbolone and aromatase inhibitors [prochloraz and fadrozole], masculinize zebrafish, leading to male skewed sex ratios. The timing of exposure was critical in that complete masculinization occurred when zebrafish were exposed from hatch or from 20 to 60 days post-hatch (dph), which corresponds to the period of gonad differentiation.

Exposure occurring outside of the period of sexual differentiation leads to changes in secondary sex characteristics or effects on the reproductive system, such as inhibition of spermatogenesis, degenerative changes in gonads, reduced fecundity and alterations of the transcription of genes involved in reproductive development. Yet, there is evidence that sex reversal in fish can be induced outside the period of gonad differentiation if exposure to EDCs is longer and in higher doses. Takatsu Takatsu et al. (2013) exposed sexually mature adult female zebrafish to the aromatase inhibitor fadrozole (0.2 mg/g) for 5 months and observed an ovarian retraction followed by the development of testes-like organs, which contained sperm heads without tails. Moreover, even after discontinuing the treatment, normal and fertilization-competent sperm were produced by these females. This study suggested that undifferentiated PGCs may persist in adult zebrafish female which enables the differentiation to male germ cells and confer sexual plasticity (Takatsu Takatsu et al., 2013).

Increasingly studies of the effects of EDCs on fish undergoing sex differentiation involve monitoring changes in the expression of genes and measurement of sex steroid hormone levels related to sex differentiation. As one example, Yang et al. (2018) exposed zebrafish embryos/larvae to 1–1000 μ g/L of Bisphenol F (BPF) for 0–60 days post-fertilization (dpf) and showed alterations in survival, growth, the balance of steroid hormones, and sex differentiation from 100 μ g/L of BPF. This led to a skewed sex ratio in favor of females in the high BPF exposure groups and a higher frequency of fish that exhibited ovotestis. Testosterone levels decreased and 17 β -estradiol levels increased in zebrafish in response to BPF. BPF exposure suppressed the expression of doublesex and mab-3 related transcription factor 1, (*dmrt1*), fushi tarazu factor 1d (*ff1d*), sry-box transcription factor 9a (*sox9a*) and *amh*; induced expression of *foxl2*, leading to increased expression of *cyp19a1a*, which promoted production of estrogens, and further caused

phenotypic feminization of zebrafish. In other studies, various EDCs, including nonylphenol, EE2, BPA, phthalates and 17 β -trenbolone or runoff from lands fertilized with animal manure from concentrated animal feeding operations (a source of EDCs), have been shown to influence sex differentiation in various fish species (Hill and Janz, 2003; Leet et al., 2015; Ye et al., 2014). Collectively, these results illustrate the sensitivity of fish to EDCs during sexual differentiation and identify effects on multiple target genes.

The permanency of EDC effects on gonad development varies depending on the compound and its mode of action, concentration, and the period of exposure (see Santos et al., 2017). Zebrafish often recover from estrogenic EDCs exposures whereas more permanent effects are induced by exposures to aromatase inhibitors. The basis of these differences is not fully understood but it has been proposed that estrogens only delay the normal sexual differentiation of males, rather than promoting real sex reversal. Alternately, androgens promote the loss of PGCs and this is significant in that PGCs are critical to ovarian development. In the absence of PGCs, development of females would not be possible. There are also concerns of possible multigenerational effects of EDCs in fish, such that the resulting progeny may have a higher sensitivity than the exposed parental generation (Santos et al., 2017; Robaire et al., this issue). For example, female-biased sex ratios were reported in F1 and F2 generation zebrafish following exposure to 1 nM (0.228 μ g/L) BPA (Chen et al., 2015a); female-biased sex ratios were reported in the untreated F1 and F2 generations following TCDD treatment of just the parents (Baker et al., 2014). While the mechanisms responsible for such effects are not known, these may involve modification of the epigenome (Robaire et al., this issue).

3. Puberty

3.1. Physiology of puberty

Like other vertebrates, puberty in fish is a transitional period in the life history that includes all the processes by which an individual attains, for the first time, the capability to reproduce as an adult, integrating life history traits, environmental and internal cues, gene expression and signaling pathways (Carrillo et al., 2014; Okuzawa, 2002; Strüssmann and Nakamura, 2002). Pubertal development starts after sex differentiation and is completed by the time of the first gametogenesis, requiring the fine tuning of the hormonal system regulated by the hypothalamic-pituitary-gonad axis (Taranger et al., 2010). The activation of this axis involves the secretion of gonadotropin releasing hormone (GnRH) by hypothalamic neurons. GnRH neurons directly innervate the pituitary and stimulate the synthesis and secretion of two gonadotropins i.e., follicle-stimulating hormone (FSH), and luteinizing hormone (LH). Both gonadotropins are secreted from the pituitary into the bloodstream, ultimately reaching the gonads to control gametogenesis and steroidogenesis (Carrillo et al., 2014; Schulz et al., 2010; Taranger et al., 2010). FSH is the first hormone to be secreted, suggesting that it is a key player in the initial stages of pubertal onset (Molés et al., 2012; Prat et al., 1996). Some studies show that the administration of FSH to prepubertal male Sea bass (*Sebastes* sp.) results in spermatogenesis progression (Mazón et al., 2014), reinforcing its role in the onset of puberty.

At the onset of puberty, SSCs and OSCs leave their quiescent stage and enter a process of proliferation that includes several cycles of mitosis followed by a differentiation into spermatogonia and oogonia, respectively (Lacerda et al., 2014; Lubzens et al., 2010, Figs. 1 and 2). These differentiated germ cells initiate a unique process in sexual reproduction, termed meiosis, considered as the starting event leading to puberty, during which their genetic content will be reduced to half (from 2n to n) for the necessary formation of gametes (n). At the same time, specialized Sertoli and Leydig cells in the testis, and follicular cells in the ovaries are activated to provide structural support, growth factors, and trigger the synthesis and secretion of sex steroids, thus providing the proper hormonal milieu needed for maturation of the gonad and gametes (Alix

et al., 2020; Kagawa, 2013; Lubzens et al., 2010; Schulz et al., 2010, Figs. 1 and 2). This process, in fish species with seasonal reproduction, is repeated cyclically throughout their life span and may be regulated by similar mechanisms (Blázquez et al., 2017).

In several fish species, the entry into meiosis is marked by significant increases in plasma levels of 11-ketotestosterone (Blázquez et al., 2017; Rolland et al., 2013; Schulz et al., 2010). Different genes also appear to be altered during pubertal onset, as is the case for *amh*, a potent inhibitor of FSH secretion, and, therefore an inhibitor of spermatogenesis (Blázquez et al., 2017; Crespo et al., 2016; Rolland et al., 2013). Moreover, expression profiling studies have revealed changes in the expression of several genes that control signaling pathways, cell proliferation-, cell cycle-, and meiosis progression at the start of pubertal development (Blázquez et al., 2017; Rolland et al., 2013). In this regard, the role of $17\alpha,20\beta$ -dihydroxy-4-pregnen-3-one ($17,20\beta P$), a natural progestin inducing spermatogonia entering into meiotic prophase, has been a topic of research in several fish species (Lubzens et al., 2010; Miura et al., 2006, 2007, 2006; Schulz et al., 2010). Moreover, retinoic acid (RA), an active derivative of vitamin A, has been reported to play a key role as a trigger of meiosis onset, not only for different fish species (Adolfi et al., 2016; Blázquez et al., 2017; Medina et al., 2019; Peng et al., 2020; Rodríguez-Marí et al., 2013), but also in rodents, birds (Bowles et al., 2008; Smith et al., 2008) and amphibians (Wallacides et al., 2009). RA homeostasis, regulated by *cyp26a1*, has an essential role in the maintenance of germ cell production. In tetrapods, the RA-signaling pathway activates the transcription of a meiosis specific gene, stimulated by retinoic acid gene 8 protein (*Stra8*) (Griswold et al., 2012). Although this gene is absent in the majority of fish species (Medina et al., 2019; Pasquier et al., 2016; Rodríguez-Marí et al., 2013), a *stra8* homolog has been identified in Southern catfish, *Silurus meridionalis* (Dong et al., 2013) and other fish species (Pasquier et al., 2016), suggesting that in fish two different mechanisms might regulate the RA-mediated entry into meiosis, one dependent and the other independent of the presence of *stra8* (Feng et al., 2015; M. Li et al., 2016b; Medina et al., 2019).

3.2. Impact of EDCs

Spermatogenesis in fish is highly dependent on steroid hormones with estrogens, androgens and progestins having distinct and critical roles (Chen et al., 2013; Melo et al., 2015; Schulz et al., 2010). Estrogens play a critical role in stem cell renewal, androgens are critical for spermatogonial differentiation, and progestins promote the proliferation of early spermatogonia and their differentiation into late spermatogonia and spermatocytes. Similarly, oogenesis and follicle development are highly dependent on steroids (Clelland and Peng, 2009; Devlin and Nagahama, 2002) with estrogens responsible for cell proliferation and vitellogenesis, androgens for feedback mechanisms in the hypothalamus and pituitary and progestins in oocyte maturation and ovulation. Given that many EDCs act by modulating steroid hormone synthesis or action, it is perhaps not surprising that there are many examples of alterations in sexual development in fish. Table 1 summarizes the available literature on the effects of EDCs on the onset of puberty in fish. Traditionally, attention has focussed on the study of the effects of natural and synthetic sex steroids, such as EE2, 17β -trenbolone, diethylstilbestrol (DES), and dihydrotestosterone (DHT). These studies have reported that sex steroids are capable of inducing precocious puberty and delays in puberty in both females and males in zebrafish and Japanese medaka (Baumann et al., 2013, 2014; Lessman and Brantley, 2020; Morthorst et al., 2010; Örn et al., 2006; Tokumoto et al., 2004, 2005).

Although a full understanding of mechanisms by which EDCs effect gonad development during puberty have yet to be realized, several studies have shown that EDCs affect folliculogenesis. For example, EE2 affected follicular development in the zebrafish, as evidenced by reduced ovarian size and a reduction in the proportion of cortical alveolus, vitellogenic and mature follicles compared to controls (Cosme et al., 2015). In addition, a more recent study reported that exposure to

1–10 $\mu\text{g/L}$ of DES for 6 days in juvenile *Casper* zebrafish delayed puberty between 5 and 10 weeks in females and males, respectively (Lessman and Brantley, 2020). These effects were found to be reversible since after cessation of DES exposure animals were capable of reaching puberty and spawning. Long-term exposure to BPA in juvenile zebrafish during gonad differentiation and puberty maturation was found to induce adverse effects on reproduction as well as delays and failure of pubertal onset in male and female zebrafish (Song et al., 2020). In this regard, chronic exposure to 1 and 10 μM BPA induced a delay in the formation of previtellogenic follicles, considered as markers of female puberty onset; in BPA treated males only a few cysts of meiotic spermatocytes and the absence of mature spermatozoa were observed. Interestingly, both males and females exposed to 10 μM BPA showed little signs of post-pubertal folliculogenesis or spermatogenesis, leading the authors of the study to conclude that exposure to BPA blocked the first step in gonad maturation (Song et al., 2020). In the same study, transgenic zebrafish lines null for *cyp19a1a* and the nuclear form of the estrogen receptor (nER: *esr1*^{-/-}, *esr2a*^{-/-} and *esr2b*^{-/-}) were used to elucidate the mechanism of action of BPA; these experiments indicated that nERs, were involved in the impairment of spermatogenesis induced by BPA. Thus, these recent studies suggest that delayed pubertal onset is due to estrogenic EDCs exerting actions prior to folliculogenesis; for some EDCs this appears to cause severe, irreversible effects on reproductive capacity. However, further investigations and the development of endpoints focussing on the effects of different EDC exposures on the early stages of testis and ovary development are warranted to identify the mechanisms of action and better understand the severity of such impacts.

Growing evidence from environmental epidemiological studies report an association between EDCs and alterations in pubertal timing (reviewed by Zawatski and Lee, 2013). In earlier studies focussing on the effects of pulp mill effluents in white suckers (*Catostomus commersoni*), alterations of gonadal development and delayed puberty were linked to reduced levels of sex steroids (McMaster et al., 1991; Munkittrick et al., 1992). Later studies determined the presence of vitellogenin (VTG) as a biomarker of alterations in gonadal maturation, particularly in male fish (Denslow et al., 1999; Garcia-Reyero et al., 2009; Sumpter and Jobling, 1995). For example, gonadal VTG protein levels and the degree of gonad maturation in juvenile zebrafish were found to be directly correlated in females and inversely correlated in males (Baumann et al., 2013), indicating that high levels of VTG in females and low levels in males are needed at the time of pubertal onset. For this reason, VTG levels have been monitored in many studies as a biomarker of the effects of EDCs on reproduction and some studies have reported VTG levels measured during puberty (Table 1). Synthetic progestagens and their derivatives, widely used for example in contraceptive pills, for pregnancy maintenance in humans or as growth promoters in livestock, are other EDCs with proven ecotoxicological effects that appear as compounds of highest environmental concern after EE2 (Fent, 2015; Kumar et al., 2015; Liu et al., 2012). For example, juvenile rainbow trout (*Oncorhynchus mykiss*) exposed to effluents from sewage treatment plants in Sweden (Fick et al., 2010) and fish from marine aquaculture farms in China (Liu et al., 2017) had elevated levels of plasma progestagens, reinforcing the need to increase the focus of research on these compounds. Although no mechanistic studies on gonad development in rainbow trout are available, juvenile zebrafish exposed during sex differentiation and puberty to the androgenic progestin, levonorgestrel (5.5–834 ng/L) and the non-androgenic progestin, progesterone (3.7–1122 ng/L) showed clear signs of precocious puberty in males (Svensson et al., 2016). Moreover, a study on the effects of 20 different progestagens demonstrated reproductive impairment in fish exposed to environmentally relevant concentrations (reviewed by Kumar et al., 2015). Although, natural progestagens have a direct effect on the onset of meiosis and are considered key players in this process (reviewed by Regidor, 2018), the molecular mechanism by which they alter the first meiosis and onset of puberty in fish remains to be elucidated. Thus, to date eco-epidemiological field studies, combined with controlled

experimental studies, support the impacts of EDCs on gonad development during puberty.

Whether EDCs act exclusively by disrupting the neuroendocrine system and/or by altering gonadal development *per se* is still unclear. Possible modes of action of EDCs include genomic and non-genomic mechanisms, as well as estrogen, androgen and thyroid receptor-mediated effects (Faheem and Bhandari, 2021), yet other mechanisms could also play a major role in EDCs-induced gonadal disruption and puberty onset alteration. For example, RA a key player in the onset of meiosis and spermatogenesis in fish (Adolfi et al., 2016; Blázquez et al., 2017; Medina et al., 2019; Peng et al., 2020; Rodríguez-Marí et al., 2013), could be a target of EDCs. In fact, effluents from pulp and paper mills that are known to contribute to impaired reproduction and delayed puberty are a significant source of ligands that bind to the RA receptors and retinoid X receptors (Alsop et al., 2003). Toxicogenomic approaches, together with the development of new non-target molecular methodologies, have proven to be a valuable approach to elucidate new molecular key events related to responses to EDCs (Caballero-Gallardo et al., 2016). However, to the best of our knowledge, very few studies in the field of fish puberty research have used this approach. Nourizadeh-Lillabadi et al. (2009) reported that precocious puberty was related to alterations of transcriptomic key regulators of steroid and thyroid hormone functions, insulin signaling and metabolic homeostasis. Therefore, we suggest that further investigations using toxicogenomic tools are required to characterize the molecular mechanisms modulated by EDCs, and to elucidate if EDCs can directly trigger the gonad to enter (or not) into meiosis and proceed towards (or arrest) gonadal development and maturation in fish.

Part 2: evidence from Mammals

Exposure to a wide variety of EDCs has been associated with adverse effects on gonad development in mammals (Marlatt et al. this issue), but there is very little certainty about their molecular mechanism of action on perinatal gonads. Here, we aimed to highlight key sensitive events of gonad development mammals (Figs. 1 and 2), and targeted our review on 4 families of EDCs for which human exposure has been tested and for which we have some insight into their mode of action, both in males (Table 2) and females (Table 3). These include: 1) estrogenic substances from therapeutic sources (i.e. DES or EE2) or epoxy resin-derived bisphenols (i.e. BPA); 2) phthalates, used to make plastics flexible, found in building materials, cosmetics, food packaging, toys and medical devices; 3) flame retardants, such as polybrominated diphenyl ethers (PBDE) and organophosphate esters (OPEs); and, 4) perfluoroalkyl and polyfluoroalkyl substances (PFAS) that comprise a large group of chemicals used for a variety of applications (i.e. stain and water repellants in clothing, carpets and paper; nonstick cookware; food packaging; firefighting foams; photographic and electronic equipment; and, industrial coatings on tiles, stones and versatile consumer products).

1. Gonad differentiation

1.1. Gonad formation

The early stages of gonadogenesis in mammals, understood primarily based on work in rodents, consist of the formation of genital crests on the ventromedial surface of the embryonic kidneys (mesonephros; Pelosi and Koopman, 2017). These crests arise from the accelerated proliferation of celiac epithelial cells, transforming the monolayer of celiac epithelium into a dense, pseudostratified form. This phenomenon occurs around gestation day (GD) 10.5 in mice, GD12.5 in rats and 4–5 weeks of gestation (GW4-5) in humans. The primordial gonads continue to develop through extensive cell proliferation, by ingression of cells from the coelomic epithelium and by recruitment of cells from the mesonephros. These cells will be the origin of the different somatic cells present in the gonads (reviewed Piprek et al., 2016). In parallel, PGCs, which have differentiated from the epiblast under the influence of bone

morphogenic protein signals from the extra-embryonic ectoderm (Lawson et al., 1999; Ying et al., 2000), will migrate from the allantois to the genital crests via the posterior part of the endoderm while multiplying rapidly (reviewed in Saitou and Yamaji, 2012; Svingen and Koopman, 2013). PGCs arrive in the genital ridges at around GD11.5 in mice, 13.5 in rats and GW6-7 in humans (Pelosi and Koopman, 2017). At that stage, the DNA of PGCs is mostly demethylated (Hill et al., 2018).

1.2. Sex determination and gonad differentiation

The newly formed gonad is bipotential and can develop into a testis or an ovary. Sex determination in mammals mostly depends on the presence of the Y chromosome and the timely expression of the gene sex-determining region of Y chromosome (*Sry*), a transcription factor of the SOX family, in cells of the coelomic epithelium in the genital crests (Gubbay et al., 1990; Sinclair et al., 1990). If *SRY* is not expressed or is transcribed in sufficient quantities, the coelomic epithelium differentiates into pre-granulosa cells and the gonads develop into ovaries (Buaas et al., 2009; Kato et al., 2013; Larney et al., 2014; Wu et al., 2012). Expression of *SRY* triggers a cascade of gene expression leading to differentiation of Sertoli cells which will induce the differentiation of other testicular cell lines and form aggregates around PGCs, creating the characteristic structures of the testis: the seminiferous cords, surrounded by a basement membrane and the interstitium (reviewed in Cool et al., 2012, Fig. 1). In the interstitium, fetal Leydig cells differentiate shortly after seminiferous cord formation, which triggers androgen production, including testosterone (Habert and Picon, 1984). In males, PGCs are called gonocytes, while female PGCs are called oogonia (Pelosi and Koopman, 2017, Figs. 1 and 2).

1.3. Impact of EDCs

To date, cases of sex reversal or, very rarely, cases of ovotestes in mammalian species (including humans), are due to genetic mutations (Dewing et al., 2002). However, a study by Yasuda et al. (1985) showed that oral administration of 0.02–2 mg/kg/d of EE2, a strong estrogen agonist, to Jc1:ICR mice from GD11 to GD17 induced the appearance of ovotestes in male fetuses and ovarian hypoplasia in female fetuses (Yasuda et al., 1985). More recently, *in vitro* experiments in a human testis-derived cell line, NT2/D1, showed that exposure to EE2 significantly decreased the mRNA levels of key genes for Sertoli cell differentiation and male gonad differentiation; these include *SOX9*, *SRY*, *AMH*, *Fibroblast Growth Factor 9 (FGF9)* and *Prostaglandin D2 Synthase (PTGD)*, and later, significantly increased the mRNA levels of key granulosa cell genes, such as *FOXL2* and *Wnt family member 4 (WNT4)* (Stewart et al., 2020). This suggests that an excess of estrogen may affect Sertoli cell differentiation. However, in rodents most *in utero* EDC exposures at the time, or including the time, of gonad differentiation, did not induce sex reversal in gonad differentiation. We can therefore conclude that there is no evidence to date that endocrine disruptors can reverse genetic sex determination, but there is evidence that they may have effects on early somatic cell differentiation. However, there is now evidence that epigenetic mechanisms play a role in sex determination (García-Moreno et al., 2018), leading to new hypotheses on environmental influence, and possible molecular targets involved in the deregulation of this process.

2. Testis development

The testis differentiates at GD12.5 in mice, GD14.5 in rats and GW6-8 in humans (Pelosi and Koopman, 2017). At this stage of development, the testis consists of two compartments representing the two main functions of the testis, gametogenesis in the seminiferous cords and steroidogenesis in the interstitium (Fig. 1).

The seminiferous cords contain the gonocytes, surrounded by Sertoli cells, and are composed of an epithelium bordered by peritubular cells and a basement membrane. Sertoli cells proliferate and differentiate until after birth when the blood-testis barrier forms (Enders, 1993). They

Table 2
Early exposure to EDCs and testicular development in mammals.

Compound	Species	Dose	Exposure Window (Route)	Effects on Testis	Molecular effects	Reference
DES	CD-1 mice	10, 50, 100 µg/kg/d	GD10.5–17.5 (Sc. Injection)	Decrease in the intratesticular testosterone level of GD18.5 testes treated with 10 µg/kg/d.	Decrease in the expression of <i>P450c17</i> , <i>StAR</i> (mRNA and protein) in GD18.5 testes treated with 100 µg/kg/d.	Guyot et al. (2004)
DES or Zeranol	NMRI mice	150 µg/kg/d	GD9-10 (Sc. Injection)	Presence of multinucleated gonocytes in GD17-18 testes. Cryptorchidism		Perez-Martinez et al. (1996)
DES or 17β-estradiol	CD-1 mice	20 µg DES or 6 mg 17β-estradiol	GD11.5, 13.5 and 15.5 (Sc. Injection)		Decrease in expression of <i>Insl3</i> at GD 17.5 and PND0	Nef et al. (2000)
DES	Sprague-Dawley Rats	1.5, 15 µg/kg/d 0.01–2 µg/kg/d	GD7-21 (Sc. Injection) GD14 – birth (Gavage)	Decrease in blood testosterone levels at 6 weeks after birth.	Increase in expression of <i>Pdgfra</i> at PND3 Decrease in expression of <i>Pdgfrβ</i> at PND3 (2 µg/kg/d)	Yamamoto et al. (2003) Thuillier et al. (2003)
EE2	CF-1 mice	0.002–20 µg/kg/d	GD14 – birth (Gavage)	Decrease in sperm count, motility and altered morphology at PND75		Ahmad et al. (2014)
	Jc1:ICR mice	0.02, 0.2 mg/kg/d	GD0-17 (micropipette) GD11-17 (Gavage)	Decrease in daily sperm production at PND50 Decrease in intratesticular testosterone level and LC hyperplasia in 20–22-month testes exposed <i>in utero</i> to 0.2 mg/kg/d.		Thayer et al. (2001) Yasuda et al. (1988)
	Sprague-Dawley Rats	0.001–10 µg/kg/d	GD11-20 (Sc. Injection)	No morphological changes in the developing reproductive system	Decrease in expression of <i>Cyp11a1</i> , <i>StAR</i> , <i>Cyp17a1</i> at the highest dose on GD20	Naciff et al. (2005)
Bisphenol A	ICR mice	5, 50 mg/kg	GD7 and GD14 (Sc. Injection)	At 6 weeks: Decrease in sperm count, motility and morphology Decreased number of SC per seminiferous tube.	Decrease in expression of genes associated with Sertoli cell functions (<i>Msi1h</i> , <i>Ncoa1</i> , <i>Nid1</i> , <i>Hspb2</i> , <i>Gata6</i>) at the highest dose.	Tainaka et al. (2012)
	C57BL/6J mice	5, 50 µg/mL (1, 10 mg/kg/d)	GD1-birth (Drinking water)	Increase in gonocyte apoptosis at PND14 (both doses) and PND35 (50 µg/mL).	Decrease in expression of <i>StAR</i> , <i>Cyp11a1</i> , <i>3βhsd</i> in PND1 testes.	Yang et al. (2019)
	Kunming mice	2.5–40 mg/kg/day	GD0.5–17.5 (Gavage)	Decrease in post-natal testis weight with the high dose Decrease in serum T and E2 at PND21 and PND56	upregulated testicular expression of <i>Dnmt1</i> , <i>Esr1</i> and <i>Ar</i> and genes related to apoptosis at PND21 Decreased testicular expression of <i>Dnmt3A</i> and <i>Dnmt3B</i> at PND 21	Wei et al. (2019)
	Sprague-Dawley Rats	4, 40, 400 mg/kg/d	GD16-21 (Gavage)	Decrease in testosterone production and the number of LC at GD 21 (40 and 400 mg/kg/d)	Decrease in expression of <i>Insl3</i> , <i>Hsd17b3</i> (protein and mRNA) at GD21.5 Decrease in expression of <i>Lhcgr</i> , <i>Cyp11a1</i> , <i>Cyp17a1</i> , <i>Amh</i> (protein and mRNA) at GD21.5 (400 mg/kg/d)	Lv et al. (2019)
		0.002–400 mg/kg/d	GD11-20 (Sc. Injection)	No morphological changes in the developing reproductive system	Decrease in expression of <i>Cyp11a1</i> , <i>StAR</i> , <i>Cyp17a1</i> in GD20 testes treated with the highest dose at GD20	Naciff et al. (2005)
		0.2–200 µg/mL	GD1 -2 h after birth (Drinking Water)	Dose-dependant decrease in blood testosterone level		Tanaka et al. (2006)
		0.1–200 mg/kg/d 200 mg/kg/d	GD14- birth (Gavage) GD14- birth (Gavage)		Increase in expression of <i>Pdgfra</i> and <i>Pdgfrβ</i> at PND3 (≥1 mg/kg/d) Decrease in expression of <i>Erk1</i> (protein and mRNA)	Thuillier et al. (2003) Thuillier et al. (2009)
	Wistar Rats	25, 250 µg/kg/d	GD10-21 (Gavage)	Decrease in the percentage of proliferating gonocytes at PND3. Decrease in sperm count and viability at PND120		Olukole et al. (2019)
Dibutyl phthalate (DBP)	Wistar Rat Wistar Rat	500 mg/kg/d	GD13-21 (Gavage)	Decrease of 90% in testosterone secretion at GD19 associated with LC hyperplasia. High rate (>60%) of cryptorchidism, hypospadias, infertility, and testicular impairment. Presence of malformed tubules and LC inside seminiferous tubules. Presence of multinucleated gonocytes up to PND10 and immature SC.		Fisher et al. (2003)
		100–500 mg/kg/d	GD13.5–20.5 (Gavage)	Decrease in testosterone and changes in LC distribution at		Mahood et al. (2007)

(continued on next page)

Table 2 (continued)

Compound	Species	Dose	Exposure Window (Route)	Effects on Testis	Molecular effects	Reference				
Di (2-ethylhexyl) phthalate (DEHP)	Wistar Rats	750 mg/kg/d	GD15.5–18.5 (Gavage)	GD21.5 Increase in occurrence of multinucleated gonocytes at GD21.5		van den Driesche et al. (2012)				
				Decrease in intratesticular testosterone level. Changes in LC distribution due to large LC aggregates in GD21.5 testes.						
		500 mg/kg/d	GD13.5–20.5 (Gavage)	Germ cells are aggregate at GD21.5 Increased occurrence of multinucleated gonocytes at GD21.5			van den Driesche et al. (2015)			
		750 mg/kg/d	GD15.5–18.5 (Gavage)	Decreased number of germ cells at GD21.5 when exposed from GD13.5–15.5 Decrease of 47–48% in intratesticular testosterone level at GD17.5 associated with aggregation of LC at GD 17.5 and 21.5			van den Driesche et al. (2017)			
	Sprague-Dawley Rats	405 mg/kg/d	GD6 –PND21 (Gavage)	Multinucleated gonocytes observed at PND1	dose-dependent increases in SOX9 and AMH at PND10 Increased CYP19A1, AR and DNMT3B in adult testis Decreased expression of <i>Cyp11a1</i> , <i>Cy17a1</i> , <i>Star</i> , and <i>Tspo</i> at GD19 Increased expression of <i>Cyp11a1</i> , <i>Cy17a1</i> , <i>Star</i> , and <i>Tspo</i> from PND3 to adults	Abdel-Maksoud et al. (2015)				
				750 mg/kg/d			GD7 –PND17 (Gavage)	Decrease in intratesticular and circulating levels of testosterone at GD21.		
		234–1250 mg/kg/d	GD14.5–birth (Gavage)	No effect on post-natal testis weight Increased serum AMH in adults No change in Serum T to E2 ratios Hyperplasia of LC No change in absolute volume of germ cells Significant decrease of fetal testosterone production and adult serum testosterone levels associated to			55 DEG in adult testis, including some related to estrogen function and signaling (Nr5a2, Ltf and Runx 2)	Albert et al. (2018); Nardelli et al. (2017)		
				30 or 300 mg/kg/day					GD8- PND21 (Gavage)	Decreased Anogenital index Increased incidence of hemorrhagic testis and multinucleated gonocytes No effect on steroid level or sperm quality in adult rats
		10 mg/kg/day	GD14.5–birth (Gavage)	No change in anogenital distance No change in adult serum testosterone Increased adult testis weight					Altered expression of mRNA coding for steroidogenic enzymes, markers of Leydig progenitor, Sertoli cells and germ cells, proliferation, redox balance and xenobiotic transporter markers from PND3 to adulthood	Jones et al. (2015), 2014
		Diisononyl phthalate (DiNP)	Wistar Rats	750 mg/kg/d					GD7-21 (Gavage)	Decrease in testosterone production and intra-testicular testosterone level at GD21
1000 mg/kg/d	GD12-21 (Gavage)			Decrease in testosterone secretion LC hyperplasia and aggregation Presence of multinucleated gonocytes						
2,2,4,4,5-pentabromodiphenyl ether (BDE-99)	Long-Evans Rat	1 or 10 mg/kg/d	GD10-18 (Sc. Injection)	Reduced anogenital distance Decrease serum testosterone		Lilienthal et al. (2006)				
BDE-47	Wistar rats	0.2 mg/kg body weight/day	GD8-birth (micropipette)	Smaller testes, Decreased sperm production, Increased percentage of morphologically abnormal spermatozoa	Decreased expression of transition proteins and protamine genes Increased expression of TNF α and IL3 Changes in sperm DNA methylation	Khalil et al. (2017) Suvorov et al. (2018)				
Hexafluoropropylene oxide dimer acid (HFPO-DA)	Sprague-Dawley rat	1–500 mg/kg-body	GD14-18 (Gavage)	No change in fetal testis testosterone production Decreased adult testis and epididymis weights	No effect on gene expression in fetal testis (genes tested are known to be affected by phthalates)	Conley et al. (2019)				
Perfluorotridecanoic acid (PFTTrDA)	Sprague-Dawley rat	1, 5, 10 mg/kg/d	GD14-21 (Gavage)	Reduced anogenital distance (high dose)	Decreased mRNA and protein of <i>Insl3</i> , <i>Lhcgr</i> , <i>Scarb1</i> , <i>Star</i> , <i>Hsd3b1</i> ,	Li et al. (2021)				

(continued on next page)

Table 2 (continued)

Compound	Species	Dose	Exposure Window (Route)	Effects on Testis	Molecular effects	Reference
				Decreased serum testosterone levels Abnormal aggregation of fetal Leydig cells	Cyp17a1, Nr5a1, and Dhh Increased antioxidants (SOD1, CAT, and GPX1), induced autophagy (increased levels of LC3II and beclin 1, and reduced the phosphorylation of mTOR)	

DES: Diethylstilbestrol; EE2: Ethynylestradiol; GD: Gestation day; PND: Post-natal Day; LC: Leydig cells; SC: Sertoli cells; #: number; Sc. Injection: Subcutaneous injection; Amh, Anti Mullerian hormone; bw, body weight.

(Note that examples of studies are presented here; the list is not comprehensive).

AR, Androgen receptor; CAT, Catalase; Cyp11a1, cytochrome P450 11 subfamily A member 1; Cyp17a1, cytochrome P450 17 subfamily A member 1; Cyp19a1, cytochrome P450 19 subfamily A member 1; Dhh, Desert Hedgehog; Dnmt, DNA methyltransferase; Esr1: Estrogen receptor 1; Erk1, Extracellular Signal-Regulated Kinase 1; Gata6, GATA Binding Protein 6; GPX1, Glutathione Peroxidase 1; Hspb2, Heat Shock Protein Family B (Small) Member 2; Hsd17b3, Hydroxysteroid 17-Beta Dehydrogenase 3; Hsd3b1, Hydroxy-Delta-5-Steroid Dehydrogenase, 3 Beta- And Steroid Delta-Isomerase 1; Insl3, Insulin like 3; Lhcgr, luteinizing hormone/choriogonadotropin receptor; LC3II, microtubule-associated protein light chain 3; Msi1h, Musashi RNA Binding Protein 1; mTOR, Mechanistic Target Of Rapamycin Kinase; Ncoa1, Nuclear Receptor Coactivator 1; Nid1, Nidogen 1; Nr5a1, Nuclear Receptor Subfamily 5 Group A Member 1; P450c17, Cytochrome P450 Family 17 Subfamily A Member 1; Pdgfra, Pdgfrb, Platelet Derived Growth Factor Receptor Alpha and beta; Scarb1, Scavenger Receptor Class B Member 1; SOD1, Superoxide Dismutase 1; StAR, steroidogenic acute regulatory protein; Tspo, Translocator Protein 18-kDa.

act as support and feeder cells for the germ cells, notably by contributing to gonocyte survival and preventing their entry into meiosis (Bowles et al., 2006; Koubova et al., 2006; Li et al., 1997). They synthesize and secrete the AMH responsible for the regression of the Müllerian ducts (Magre and Jost, 1984). Gonocytes continue to proliferate for a few days until they enter a quiescent phase characterized by a cell cycle arrest in the G0/G1 phase at about GD16.5 in mice, GD18.5 in rats and GW18-19 in humans; the cell cycle resumes after birth in rodents (Culty, 2009, Fig. 1). Around this time, gonocytes undergo the epigenetic reprogramming that corresponds to the establishment of new DNA methylation marks and chromatin remodelling (Ly et al., 2015; Rwigemera et al., 2021; Wu et al., 2015). This contributes to the establishment of the male germ cell transcriptome and the formation of the spermatogonial stem cell pool in the neonatal testis (Culty, 2009; Manku and Culty, 2015; Rwigemera et al., 2021).

In the interstitium, fetal Leydig cells differentiate and contribute to the production of androgens and insulin-like 3 (INSL3) that are key to the further development of the testis and differentiation of the reproductive tract, inducing the maturation of the Wolffian duct and testicular descent. In rodents, Sertoli cells also contribute to steroidogenesis by converting androstenedione (A4) to testosterone, as fetal Leydig cells do not express the enzyme catalyzing this conversion, namely 17 β -HSD (17 β -hydroxysteroid dehydrogenase; Shima et al., 2013). In primates, substantial A4 is also produced by the fetal adrenal; in contrast, no androgens are produced by fetal rodent adrenals. Leydig cell steroidogenesis is responsive to LH stimulation from the fetal pituitary but, in rodents, it is mostly regulated by paracrine and autocrine factors, while in primates, chorionic gonadotropins enhance fetal Leydig cell androgen production (reviewed by Sharpe, 2020).

2.1. The male masculinization window – a key stage of sensitivity to EDCs

After the testis is formed, further masculinization of the reproductive tract is driven mostly by testosterone, AMH and INSL3 (Sharpe, 2006). At this stage, all testicular cell types express both the estrogen and androgen receptors; data from murine null mutation studies for the various forms of these receptors have shown that the proper development of these cells is in part regulated by steroid hormones (Rouiller-Fabre et al., 2015). Given this, and the fact that the masculinization of the reproductive system is sensitive to androgens, it is not surprising that this period of testicular and male genital tract development is particularly sensitive to potential endocrine disruption. This hypothesis was proposed in 1993 by Sharpe and Skakkebaek, who suggested that an impairment of androgen production or an excess of estrogen during this window of sensitivity could be the origin of developmental abnormalities observed in the testicular dysgenesis syndrome (Sharpe and

Skakkebaek, 1993). Since then, experimental studies using flutamide, an anti-androgen, have allowed a better definition of the window of development during which androgen action is key for the establishment of reproductive functions in the male. This window of masculinization occurs between GD 15.5 and 19.5 in rats (corresponding to GW8 - 14 in humans) (Welsh et al., 2008). Developmental changes in testicular sensitivity to estrogens have also been observed in experiments with organ culture of perinatal testis sampled at different times during fetal and neonatal life, identifying a similar period of sensitivity to both androgens and estrogens (Delbès et al., 2007).

2.2. Impact of EDCs on testicular steroidogenesis

The effects of EDCs on steroidogenesis have been reviewed recently (Walker et al., 2021). EDCs have been implicated in impairing steroid hormone synthesis by the fetal testis, thereby affecting testicular development and masculinization. Table 2 summarizes some of the available *in vivo* exposure studies in rodents that have been particularly useful in characterizing the effects of EDCs on testicular steroidogenesis. Since there are differences between rodent and human testicular development, endocrinology and metabolism, a xenograft model in which human fetal testis is grafted to mice exposed to EDCs is also of interest (Heger et al., 2012), although a limitation of this approach is the age range at which fetal human testicular tissue can be obtained. Other tools for screening the impact of EDCs on testicular steroidogenesis are available. These include organ cultures of fetal testis (rat, mouse, human (early stages)) (Habert et al., 2014) and testicular cultures of primary cells or cell lines. There are currently no human Leydig cell models or fetal Leydig cell models, but several rodent adult Leydig cell lines, have been used to characterize the effects of chemicals on steroidogenesis (mouse LTC-1, MA-10 and TM3 (Nikula et al., 1999; Schang et al., 2016; Walsh et al., 2000) and rat R2C (Balbuena et al., 2013; Heneweer et al., 2004).

Table 2 highlights several studies in rats showing the negative impact of *in utero* exposure to **estrogenic substances**, such as EE2, DES and BPA on testosterone secretion. Some of these studies, including transcriptomic analyses of the testis, have shown an association with decreased expression of genes encoding steroidogenesis-related transcripts, such as steroidogenic acute regulatory protein (StAR), cytochrome P450 family 11 subfamily A member 1 (Cyp11a1), cytochrome P450 family 17 subfamily A member 1 (Cyp17a), hydroxysteroid 17 beta dehydrogenase 3 (Hsd17b3) and cytochrome P450 family 17 subfamily A member 1 (P450c17; Guyot et al., 2004; Lv et al., 2019; Naciff et al., 2005; Yang et al., 2019). Interestingly, the deleterious effects on steroidogenesis observed in the fetal (GD18) and post-natal (PND1) testes also persist into adulthood suggesting that early exposure to EDCs may

Table 3
Early exposure to EDCs and ovarian development in mammals.

Compound	Species	Dose	Exposure Window (Route)	Effects on Ovary	Molecular Effects	Reference
DES	CD-1 mice	0.5 mg/kg/d	PND 1–5 (Sc. injection)	Breakdown of nests of oogonia; reduced number of small antral follicles; induced multioocyte follicle formation	Decreased activin β subunit expression	Iguchi et al. (1986); Kipp et al. (2007)
EE2	C57BL/6J mice	0.0067–0.067 mg/kg/d	GD 10–18 (Sc. injection)	Reduced survival of foetuses and newborns; polyovular follicles; ovary-independent vaginal epithelial stratification		Kirigaya et al. (2006)
Bisphenol A	ICR/Jcl mice	10 mg/kg/d	GD 10–18 (Sc. injection)	Reduced the number of ovarian corpora lutea		Suzuki et al. (2002)
Bisphenol A	ICR/Jcl mice	0.150 mg/kg/d	Birth-PND 5 (Sc. injection)	Ovary-independent vaginal epithelial stratification		Suzuki et al. (2002)
Diethylhexyl phthalate	CD-1 mice	0.04 mg/kg/d	GD 1-birth (oral)	Delayed meiotic progression of early germ cells (17.5 DPC); delayed follicle assembly in newborn ovary; decreased number of primordial and increased number of secondary follicles (PND21) in F1 and F2 offspring	Reduced expression of STRA 8 (gene and protein) at 13.5 DPC; significant downregulation of <i>Cyp17a1</i> and <i>Cyp19a1</i> gene expression in fetal ovaries	Zhang et al. (2015)
	CD-1 mice	0.02, 0.2, 200, 500, or 750 mg/kg/d	GD 10.5-birth (oral)	Decreased folliculogenesis and increased serum estradiol (F1); dysregulated folliculogenesis and disrupted serum progesterone (F2); accelerated folliculogenesis (F3)		Rattan et al. (2018)
	CD-1 mice	0.02, 0.2, 500, or 750 mg/kg/d	GD 10.5-birth (oral)	Transgenerational disruption of DNA methylation (F1, F2 and F3)	Transgenerational (F1, F2 and F3) suppression of gene expression pathways required for folliculogenesis and steroidogenesis	Rattan et al. (2019)
	BalB/C mice	0.0025, 0.005, 0.010 mg/kg/d	PND 0–4 (IP injection)	Impaired germ nest breakdown and primordial follicle assembly		Mu et al. (2015)
	BalB/C mice	0.6 mg/kg	PND 0–4 (IP injection)	Severe disruption of primordial follicle formation; premature ovarian senescence and reduced fertility	Enhanced autophagy: increased autophagy-related gene expression and recognizable autophagosomes	Zhang et al. (2018)
	CD-1 mice	0.02, 0.04 mg/kg/d	PND 7–14 (hypodermal injection) (treatment 1) PND 5,10,15, and 20 (hypodermal injection) (treatment 2)	Decreased number of primordial follicles and increased secondary (treatment 2) and antral follicles.	Reduced and/or delayed methylation of imprinted genes such as <i>Igf2</i> and <i>Peg3</i> in oocytes Increased abnormal metaphase II spindles in oocytes matured <i>in vitro</i> (treatment 2)	Zhang et al. (2013)
	CD-1 mice	0.02, 0.04 mg/kg/d	PND 5, 10, and 15 (IP injection)	Inhibited antral follicle enlargement process	Increased mRNA levels of the apoptosis related genes; disturbed oxidative status	L. Li et al. (2016)
Mono-2-ethylhexyl phthalate (MEHP)	C57/Bl6 mice	100, 500, or 1000 mg/kg	GD 17–19 (gavage)	Reduced reproductive lifespan in F1 (1000 mg/kg)	Altered mRNA for the LHCGR and steroidogenic genes (aromatase, StAR)	Moyer and Hixon (2012)
Pentabrominated PBDE-99	Wistar rats	0.06, 0.3 mg/kg	GD 6 (gavage)	Increased rate of resorptions		Talsness et al. (2005)
Pentabrominated PBDE-99	Long Evans rats	1, 10 mg/kg/d	GD 10–18 (Sc. Injection)	Decreased ovarian primordial and secondary follicles		Lilienthal et al. (2006)
Tetrabrominated PBDE-47	Wistar rats	0.14, 0.7 mg/kg	GD 6 (gavage)	Decreased numbers of secondary and antral follicles and decreased ovary weights; decreased serum estradiol		Talsness et al. (2008)
Perfluorononanoic acid (PFOA)	Mice	3 mg/kg/d	GD 1–18 (oral)	Decreased ovary size; decreased numbers of primary, secondary and antral follicles, and corpora lutea; delayed onset of sexual maturity		Zhang et al. (2020)
Perfluorobutane sulfonate (PFOS)	Mice	200, 500 mg/kg/d	GD 1–20 (oral)	Reductions in all stages of ovarian follicles; decreased ovary and body size and weight; decreased serum estradiol; delayed onset of sexual maturity		Feng et al. (2017)
	Sprague-Dawley rats	0.1, 1, 10 mg/kg/d	PND 1–5 or 26–30 (Sc. Injection)	Reduction in the number of primordial and growing follicles and corpora lutea		Du et al. (2019)

DES: Diethylstilbestrol; EE2: Ethynylestradiol; GD: Gestation day; PND: Post-natal Day; Sc. Injection: Subcutaneous injection; IP: intraperitoneal. (Note that examples of studies are presented here; the list is not comprehensive).

Cyp17a1, cytochrome P450 17 subfamily A member 1; *Cyp19a1*, cytochrome P450 19 subfamily A member 1; *Igf2*, insulin like growth factor 2 receptor; LHGCR, luteinizing hormone/choriogonadotropin receptor; *Peg3*, paternal expressed gene 3; StAR, steroidogenic acute regulatory protein; Stra8, stimulated by retinoic acid gene 8 protein.

impair the establishment and function of adult Leydig cells (Yamamoto et al., 2003; Yang et al., 2019; Yasuda et al., 1988). The inhibitory effect of estrogenic substances on testosterone production was also observed throughout fetal life in dispersed testicular cell cultures and in organ cultures of rat and mouse fetal testes (Delbès et al., 2007; N'Tumba-Byn et al., 2012). However, the effects on testosterone that were observed in early fetal stage whole rat testes were masked by GD20.5, due to local production of estrogens (Delbès et al., 2007).

The human fetal testis seems to have a different sensitivity than rodents since it has been reported to be insensitive to the effects of DES in both the organ culture (N'Tumba-Byn et al., 2012) and the xenograft models (Mitchell et al., 2013), but more sensitive to the effects of BPA, at least in organ culture (N'Tumba-Byn et al., 2012). Studies with estrogen receptor knock-out (ERKO) mice have revealed that the effect of DES on steroidogenesis is mediated by ESR1, while the impact of BPA is independent of estrogen receptor (Delbès et al., 2007; N'Tumba-Byn et al., 2012). These data strongly suggest that the human testis is insensitive to effects mediated by the ESR1 at these early stages (Rouiller-Fabre et al., 2015). Other estrogen mediated pathways activated by BPA, such as estrogen related receptor gamma (ERRγ) (Liu et al., 2014), may play important roles, but have not yet been fully characterized in fetal rodent or human testes.

Similarly, many studies have described the negative impact of **phthalates** on testicular steroidogenesis. Although the exact molecular mechanism of action of phthalates on steroidogenesis has yet to be resolved, there is a growing body of evidence that such suppression of steroidogenesis is mediated by the activation of peroxisome proliferator-activated receptor (PPAR) α and γ nuclear receptors (Martinez-Arguelles et al., 2013). Fetal exposure during the masculinization window to dibutyl phthalate (DBP), diethylhexyl phthalate (DEHP) or diisononyl phthalate (DiNP) leads to underdevelopment and/or malformations of the male reproductive tract and genitalia as a consequence of reduced testosterone secretion by fetal Leydig cells (Table 2; Borch et al., 2004; Chen et al., 2021; Culty et al., 2008; Fisher et al., 2003; Mahood et al., 2007; van den Driesche et al., 2017, 2015). Outcomes in adult male rats associated with fetal exposures to these phthalates include decreased weights of reproductive organs, reduced anogenital distance (AGD), disrupted seminiferous epithelium with reduced spermatogenesis and decreased numbers of Sertoli and germ cells; in addition, malformations such as prostatic or epididymal agenesis, hypospadias, cryptorchidism, retained thoracic areolas or nipples have been observed after fetal exposure to phthalates. This suite of effects has been labelled the Phthalate syndrome (NRC, 2008). These effects are often associated with hyperplasia and aggregation of fetal Leydig cells and decreased expression of steroidogenic enzymes. Among the earliest gene expression changes in the fetal rat testis following *in vivo* exposure to an effective dose of phthalate are transcripts and proteins involved in the synthesis and uptake of the steroid hormone precursor cholesterol and in steroid hormone synthesis (Johnson et al., 2011). The potency by which phthalate esters induce these effects is highly dependent on the structure and location of the ester side chains; the more potent phthalates have side chains between 4 and 9 carbons, with potency diminishing with shorter chains (Gray et al., 2000). In contrast with the effects in rats, DBP exposure of mice during the masculinization window does not result in reduced testis androgen production (Albert and Jégou, 2014; Johnson et al., 2012), or alter the expression of steroidogenesis associated genes (Johnson et al., 2011). Yet, DBP exposure results in aggregation of fetal Leydig cells and increased incidence of hypospadias in exposed mice (Albert and Jégou, 2014). Similarly, studies using organ culture and xenograft of human fetal testes suggest that phthalates have little impact on fetal human testis steroidogenesis (Hallmark et al., 2007; Heger et al., 2012; Mitchell et al., 2012).

Effects of phthalate exposures on a range of other endpoints of human male development are less clear but have received considerable attention; this issue has been extensively reviewed (Albert and Jégou, 2014; Arzuaga et al., 2020; Habert et al., 2014; Howdeshell et al., 2008;

Johnson et al., 2012; National Toxicology Program, 2003). The ubiquitous presence of diverse phthalates means that the majority of fetuses through the world have measurable exposure to some combination of phthalates, mostly at doses below those observed to disrupt male reproductive tract development in rats. Multiple studies have examined the correlation between phthalate exposures during early pregnancy (corresponding to the human masculinization window) and AGD; the results of these studies have been inconsistent. While several studies revealed a robust correlation between human fetal exposures to phthalates and reduced AGD in male infants (Swan et al., 2005, 2015), others have not found such a relationship (Jensen et al., 2016) or found that exposure was correlated with longer AGD (Arbuckle et al., 2019). When these data were considered in a meta-analysis, fetal exposures to DEHP was significantly associated with reduced AGD in human male infants (Dorman et al., 2018; Zarean et al., 2019). These results suggest that *in utero* exposures to certain phthalate esters may be associated with impaired androgen-dependent male reproductive tract development although fetal testis androgen production may be unaffected.

The effects of **PBDEs and OPEs**, chemicals commonly used as flame retardants, on testicular development have been reviewed recently (Hales and Robaire, 2020). There is evidence from several epidemiological studies that exposure to brominated flame retardant (BFRs) during gestation may affect endpoints in male offspring such as cryptorchidism or anogenital distance. In a study from Sweden, an association between an elevated incidence of cryptorchidism and PBDE concentrations in breast milk was significant (Main et al., 2007). Similarly, a study in a Canadian population reported that every 10-fold increase in maternal hair 2,2,4,4,5-pentabromodiphenyl ether (BDE-99) or 2,2',4,4',6-pentabromodiphenyl ether (BDE-100) was associated with at least a doubling in the risk of having a child with cryptorchidism (Goodyer et al., 2017). Data from Shanghai have suggested that prenatal exposure to 2,2',4,4'-tetrabromodiphenyl ether (BDE-47) and Σ4PBDEs, even at low environmental levels, may be associated with shorter AGD in boys (Luan et al., 2019). Studies in several experimental models have provided evidence that exposure to PBDEs may affect testicular steroidogenesis, suggesting altered steroidogenesis as a possible causal factor inducing the phenotypes observed in human cohorts. In fetal porcine Leydig cells, several BFRs reduced testosterone synthesis by inhibiting specific enzymes in the steroidogenesis pathway (Mankidy et al., 2014). Using the mouse LTC-1 Leydig cells, Han et al. (2019) demonstrated an inhibition of steroid production after exposure to deca-brominated diphenyl ether (BDE-209). Treatment of Long-Evans rat dams with a single dose of BDE-99 on gestation day 6 reduced AGD and decreased circulating sex steroids in male offspring (Lilienthal et al., 2006). The underlying mechanism(s) for such effects have yet to be elucidated.

The persistence, bioaccumulation and toxicity of PBDEs has led to regulations of their use and created a need for their replacement to meet the flammability standards in many jurisdictions; the OPEs are substances that are commonly used as replacement flame retardants. While studies of the mechanism of action of these chemicals during fetal development on steroidogenesis are scarce, a few have demonstrated that these chemicals do affect steroid production in cultured Leydig cells. Tris (methylphenyl) phosphate (TMPP) was found to suppress TM3 Leydig cell testosterone secretion (Liu et al., 2016). Another OPE, 2-(butoxyethyl) phosphate (TBEP), was shown to induce an increase in oxidative stress and a decrease in steroid production in TM3 cells (Jin et al., 2016). Schang et al. (2016) compared the effects of seven OPEs on MA10 Leydig cell functions with those of BDE-47. All the OPEs affected mitochondrial activity, cell survival, and superoxide production. Five of the seven OPEs caused an increase in basal or cAMP-stimulated steroid secretion and several OPEs affected the expression of genes involved in steroid biosynthesis. Thus, the limited data available to date suggest that both PBDEs and OPEs may affect Leydig cell functions.

The effects of **PFAS** on testicular steroidogenesis have been reviewed recently (Zhu et al., 2020), highlighting three *in utero* exposure studies of perfluorooctanoate (PFOA), perfluorooctane sulfonate (PFOS) and

perfluorononanoic acid (PFNA) that reported negative effects on Leydig cell development, testosterone secretion, and expression of steroidogenic enzymes. Similarly, a recent study of gestational exposure to perfluorotridecanoic acid (PFTrDA) in rats reported inhibition of the differentiation of fetal Leydig cells mainly through increasing oxidative stress and inducing autophagy (Li et al., 2021). *In utero* exposure of rats from GD14-18 to another perfluoroalkyl substance, hexafluoropropylene oxide dimer acid (HFPO-DA, also known as GenX; 1–500 mg/kg/day) was not reported to affect fetal testis development but negative effects on pup birth weights and the liver were observed (Conley et al., 2019, 2021).

2.3. Impact of EDCs on gametogenesis in males

Table 2 summarizes some of the available *in vivo* exposure studies in rodent models. Evidence from *in vivo* and *in vitro* experimental models range from immediate effects on germ cell proliferation or death, the appearance of multinucleated germ cells, changes in gene expression, or effects on the epigenetic reprogramming that takes place prior to the establishment of SSCs in males, to long term impacts on sperm production.

Experiments of early exposure to **estrogenic substances** in rats have shown immediate and long-term impacts on male germ cells. Gonocyte proliferation, essential for germline development, has been shown to be sensitive to exogenous estrogens such as DES, with high sensitivity during the fetal proliferation phase (Delbès et al., 2007; Lassurguère et al., 2003). Long term effects of early exposure to BPA, DES and EE2 on the adult germline have also been observed; these included increased germ cell apoptosis, reduced sperm count and motility and increased sperm morphological abnormalities (Ahmad et al., 2014; Olukole et al., 2019; Tainaka et al., 2012; Thayer et al., 2001; Yang et al., 2019). Using both *in vitro* and *in vivo* models of exposure to estradiol, DES or BPA, Culty and collaborators have shown that estrogens can target PGDF signaling in gonocytes and alter germ cell proliferation and differentiation after birth (Li et al., 1997; Manku et al., 2015; Thuillier et al., 2003, 2009, 2003; Wang and Culty, 2007). Similarly, one study reported an impact of *in utero* low-dose exposure to EE2 or BPA on the testis transcriptome, suggesting these chemicals may affect the expression of genes important for germ cell differentiation, even at doses at which no immediate phenotypical or histological changes were observed (Naciff et al., 2005). Interestingly, multinucleated gonocytes were observed in murine testes at GD17-18 after exposure to DES or Zeranol (150 µg/kg/day) from GD9-10, yet the mechanism is unknown (Peréz-Martínez et al., 1996). Organ culture and xenograft experiments have revealed that exposure to 10 µM BPA may have negative effects on germ cells density and differentiation, suggesting similar sensitivity in human and rodents (Eladak et al., 2018).

There are several reports of the negative impact of **phthalates** on the male germline with many observations of decreased numbers of germ cells and the presence of multinucleated gonocytes after birth, immature Sertoli cells and focal testicular dysgenesis (Table 2) (Andrade et al., 2006; Fisher et al., 2003; Mahood et al., 2007; Nardelli et al., 2017; van den Driesche et al., 2017, 2015, 2012). Early exposure to DEHP has been reported to have immediate and long-term effects on testicular gene expression, some of which being related to the differentiation of germ cells (Jones et al., 2014, 2015). Interestingly, after MEHP exposure similar effects were observed in both rodent and human fetal testes; these included an increase in germ cell death in the absence of an effect on the proliferation or apoptosis of Sertoli cells (Habert et al., 2014). As MEHP does not affect testosterone production in humans, this suggests that MEHP may act directly on germ cells. Exposure of ERKO mice or AR-deficient mice to MEHP revealed that MEHP-induced germ cell damage is not predominantly mediated by steroid receptors (Rouiller-Fabre et al., 2015). Instead, MEHP targets PPAR and liver X receptor LXR pathways in the fetal testis; these pathways are also linked to the effects of MEHP in adults (Rouiller-Fabre et al., 2015).

Exposures during gestation to **PBDEs** have also been found to affect

male germ cells. In a study on Wistar rats exposed to BDE-47 during gestation from day 8 onward, Khalil et al. (2017) reported that offspring had significantly smaller testes, decreased sperm production, and an increased percentage of morphologically abnormal spermatozoa. The expression of genes essential for spermatogenesis, such as transition proteins and protamines, was suppressed and immune response genes, such as the tumor necrosis factor 3 (Tnf 3) and interleukin 3 (Il3), were increased; these authors suggested that this treatment might be producing an aberrant sperm epigenome. A subsequent study revealed that BDE-47 exposure altered sperm DNA methylation (Suvorov et al., 2018). Using a stem cell-based spermatogenesis model, Greeson et al. (2020) showed that exposure to 2,2',4,4',5,5'-hexabromodiphenyl ether (BDE-153) decreased *de novo* and maintenance DNA methylation at regulatory elements controlling imprinted genes. Furthermore, PBB-153 affected DNA methylation as well as the expression of genes critical to proper human development (Greeson et al., 2020). Studies of the effects of two other flame retardants, hexabromocyclododecane (HBCDD) and tetrabromobisphenol A (TBBPA), on a human stem cell-based model of spermatogenesis, revealed that mitochondrial membrane potential was perturbed and reactive oxygen species were increased, leading to the apoptosis of spermatogonia and primary spermatocytes (Stevens et al., 2018).

A recent review of the potential impact of **PFAS** on male fertility has linked increased exposure to PFAS with effects on semen quality (Tarpore and Ouyang, 2021). Yet, observations in experimental rodent exposure studies showed inconsistency from no effect to some negative effects on sperm production. Such variation may be due to species- and strain-specific differences in PFAS metabolism, the rat being the least adequate to compare PFAS toxicity studies relative to human health.

Together, these studies suggest that there are deleterious effects induced by a variety of EDCs on germ cells. Whether all of these are entirely direct or indirect (i.e. mediated by effects on somatic cell impairment) that consequently affect germ cells, remains to be determined. With recent development of flow cytometry-based methods to purify mouse and human germ cells, a few investigations have reported molecular impacts specifically on germ cells. Most report little immediate impact on germ cells' transcriptome and DNA methylation (Iqbal et al., 2015; Muczynski et al., 2012), but many other studies reported long-term impacts of *in utero* exposure on the offspring sperm epigenome (Robaire et al., this issue; Ho et al., 2017). These observations support the original TDS hypothesis proposed by Niels E. Skakkebaek and colleagues that it is the immediate early effects on the testicular somatic environment that subsequently influence germ cell development (Skakkebaek et al., 2016).

3. Ovary development

3.1. Primordial follicle formation

When the ovary is formed, the female PGCs differentiate into oogonia undergoing extensive mitotic proliferation and forming clusters of interconnected cells known as germ cell cysts or nests (Pepling, 2012; Pepling et al., 1999; Wang et al., 2017, Fig. 2). Around GD13.5 in mice, GD17.5 in rats and GW12-13 in humans, the germ cells enter meiosis (Motta, 1997; Wang et al., 2017). Oogonia become oocytes and progress through meiotic prophase I until they become arrested at the diplotene stage. Approximately two thirds of the oocytes in the germ cell nests undergo programmed cell death (involving both apoptosis and autophagy), germ cell cysts break apart and the remaining single oocytes become surrounded by a flattened (squamous) layer of intruding somatic pre-granulosa cells, forming the primordial follicles (Johansson et al., 2017; Pepling, 2012). Germ nest breakdown and primordial follicle formation are prerequisites for the establishment of the ovarian reserve; among the many factors implicated in this process are programmed cell death regulators (BCL2 associated X, apoptosis regulator and BCL2 apoptosis regulator), growth factors and signaling molecules (activin A, AMH, notch signaling pathway, the

phosphoinositide-3-kinase (PI3K) pathway), transcription factors (aryl hydrocarbon receptor), hormones (estradiol, progesterone, FSH), meiotic regulators (RA, activin A, WNT4) and cell adhesion molecules (Pepling, 2012; Sun et al., 2017; Wang et al., 2017). In humans, primordial follicle assembly is initiated around mid-gestation and completed during fetal life, while in rodents the process continues until PND6 (Johansson et al., 2017). Fetal and neonatal periods are, therefore, critical windows for the establishment of the primordial follicles and any disturbance of the process may result in premature ovarian senescence in the offspring.

3.2. Folliculogenesis

Folliculogenesis occurs after puberty (Fig. 2). Four major developmental events can be distinguished in this process: primordial follicle recruitment; preantral follicle development; selection and growth of the antral follicle; and follicle atresia (Williams and Erickson, 2012). Folliculogenesis is irreversible once it has begun; thus, after primordial follicle recruitment is initiated, the follicle is destined to either ovulate or undergo atresia (Johansson et al., 2020; Zama and Uzumcu, 2010). In fact, approximately 99% of follicles undergo atresia, with only a small number reaching the preovulatory stage. This process is controlled by a precise balance of pro- and anti-apoptotic factors (Williams and Erickson, 2012). The primordial-to-primary follicle transition is characterized by oocyte growth and granulosa cell differentiation from a squamous to cuboidal form; these cells start to express FSH receptors (Edson et al., 2009). In the secondary or pre-antral follicles, granulosa cells continue to proliferate and form two or more layers. It is at this stage that the thecal cells begin to emerge and form a layer around the granulosa-oocyte structure (Zama and Uzumcu, 2010); the main function of thecal cells is the production of androgens for neighboring granulosa cells to convert to estrogens (Edson et al., 2009). This phase is known as gonadotropin-independent growth (or the pre-antral phase) and is under the control of autocrine and paracrine regulatory factors (Edson et al., 2009; Hannon et al., 2015). Follicles then develop into antral follicles that are characterized by more than five layers of granulosa cells, a fluid filled cavity (antrum), and an oocyte enclosed by cumulus cells that are derived from the granulosa cells. Further, thecal cells develop into the theca externa and theca interna layers; the theca interna start to express receptors for LH. At this stage, follicles become gonadotropin-dependent (controlled mainly by pituitary FSH and LH) and continue to grow until they reach a preovulatory size (Edson et al., 2009). A surge of LH can result in maturation and completion of the first meiotic division and ensuing ovulation (Edson et al., 2009; Johansson et al., 2020).

Intraovarian triggers are responsible for primordial follicle activation (Mark-Kappeler et al., 2011). A number of growth factors, hormones, and cytokines have been shown to be important for primordial follicle recruitment and maintenance. PTEN (phosphatase and tensin homolog), Foxo3a (a transcription factor produced in both oocytes and granulosa cells), and SDF-1 (stromal derived factor-1) generated by primordial oocytes restrain their own activation (Williams and Erickson, 2012). Kit ligand (KL, also known as stem cell factor secreted by pre-granulosa cells) has been found to induce the transition of primordial to primary follicles (Oktem and Urman, 2010; Parrott and Skinner, 1999) exerting its effects through the PI3K pathway, a critical determinant of primordial follicle activation (Crain et al., 2008; Edson et al., 2009; Sarraj and Drummond, 2012, Fig. 2). Oppositely, AMH, produced by the granulosa cells of growing follicles, exerts an inhibitory influence on primordial oocytes, keeping them in a dormant state (Johansson et al., 2020; Zama and Uzumcu, 2010) and thus negatively regulating recruitment (Williams and Erickson, 2012). Androgens also have a clear and important physiological role in follicle development at all stages (preantral and antral follicle development), including the promotion of granulosa cell proliferation (Franks and Hardy, 2018).

3.3. Impact of EDCs on ovarian development and function

Reviews of the impact of EDCs on ovarian development and function

have discussed specific time windows when the ovary is sensitive to disruption. These times/events, include gonadal sex determination, meiosis, follicle assembly and the first wave of follicle recruitment (Johansson et al., 2017). EDCs also affect folliculogenesis and steroidogenesis in the adult ovary (Patel et al., 2015).

Studies in rodents have revealed that exposures to **estrogenic substances** during early life influence the number of normal primordial follicles and increase the number of abnormal ovarian follicles containing multiple oocytes or multi-ovular follicles (MOF) (Table 3). MOF contain two or more oocytes, each with distinct zona pellucida, within the same granulosa cell layer, basement membrane and theca layer. Rodents exposed to DES, a pharmaceutical estrogen, early in life exhibit reduced primordial follicles (Kipp et al., 2007; Rodríguez et al., 2010). The frequency of MOF in the post pubertal or adult ovaries of mice was increased by DES exposure during late fetal development and prior to weaning (Iguchi et al., 1986, 1990; Iguchi and Takasugi, 1986; Kipp et al., 2007). The most dramatic increase in the frequency and relative number of MOF occurred in females exposed in the first 5 postnatal days (Iguchi et al., 1986), coinciding with the breakdown of nests of oogonia and early follicle formation (Fig. 2). The formation of MOF induced by DES exposure appeared to be independent of the effects of DES on the developing hypothalamic-pituitary-gonad axis, as DES caused a similar effect in neonatal ovaries implanted in ovariectomized adult females or cultured *ex vivo* (Iguchi et al., 1990). In addition, oocytes collected from super-ovulated juvenile mice treated as neonates with DES had substantially reduced capacity for fertilization *in vitro* (Iguchi et al., 1990) and *in vivo* (Iguchi et al., 2002) compared to those from untreated mice. No MOF were observed in ovaries of mice lacking functional nuclear ESR2 while ESR1 knockout mice remained vulnerable (Kirigaya et al., 2006) suggesting that MOF formation is mediated via DES signaling through ESR2. DES treatment of the neonatal ovary reduced activin mRNA and protein (Kipp et al., 2007) and increased the mRNA of inhibin, an activin antagonist (Oikawa et al., 2019). Like DES, many other estrogenic substances have similar impacts on follicular formation in rodents. Exposure of neonatal mice or rats to chemicals with estrogenic activity, such as BPA (Suzuki et al., 2002), EE2 (Kirigaya et al., 2006), genistein (Losa et al., 2011) and tamoxifen (Irisawa and Iguchi, 1990), all cause disruption of follicular formation with increased MOF in the ovaries after puberty.

Despite extensive study, no specific ovarian phenotype has been described for women exposed to DES as fetuses. In extrapolating from rodent studies to humans, it is important to consider that oocyte nest breakdown occurs mid-gestation in the human fetus when the fetal zone produces estrogens from adrenal androgens; while in rodents oogonial nest breakdown and early follicle formation occur very late in gestation and continue into the early post-natal period (Pepling and Spradling, 2001, Fig. 2). This process in mice is believed to be initiated by a reduction in estrogen levels in the late fetal ovary (Chen et al., 2007; Dutta et al., 2014). In a study in baboons, treatment with an aromatase inhibitor through mid gestation resulted in a reduction in follicle number, and surviving follicles were in a poor condition at puberty (Pepe et al., 2006). This suggests not only that estrogen has a critical role in follicle survival but also that the timing and presence of endogenous estrogens may render follicular formation in the primate ovary less sensitive than in the rodent, where nest breakdown is driven by germ cell apoptosis due to a precipitous decline in estrogen exposure perinatally.

A number of studies have shown that early life exposure to **phthalates** can affect ovarian development in female offspring, disrupting fundamental processes like meiotic initiation, primordial follicle assembly and folliculogenesis (Table 3, Fig. 2). One of the most studied phthalates is di (2-ethylhexyl) phthalate (DEHP) and its active metabolite, mono (2-ethylhexyl) phthalate (MEHP). In mice, the daily oral administration of DEHP (40 µg/kg) throughout gestation led to reduced serum estradiol levels in the dams and significantly downregulated expression of *Cyp17a1* and *Cyp19a1* in the fetal ovaries at GD12.5. A

delay in the meiotic progression of early germ cells was observed in the fetal ovaries on GD17.5. This effect was associated with considerably reduced mRNA and protein expression of *Stra8* at GD13.5, suggesting that DEHP exposure can deregulate meiosis related genes, although the exact molecular mechanisms remain to be determined (Johansson et al., 2020; Zhang et al., 2015). Large regions of germ-cell cysts and rare follicles were also observed in newborn ovaries, indicative of delayed follicle assembly (Zhang et al., 2015). As in rodents, meiotic progression and primordial follicle assembly are related; reduced *Stra8* expression can explain reduced numbers of assembled primordial follicles (Johansson et al., 2017). However, DEHP impaired primordial follicle formation might also be caused by oxidative stress in germ cells, as well as by disturbances in the interaction between germ cells and pre-granulosa cells mediated by transforming growth factor-beta (TGF-beta) signaling, as revealed recently by Wang et al. (2021). Further, gestational DEHP exposure significantly decreased the number of primordial follicles and increased the number of secondary follicles at PND21, indicating an accelerated rate of follicle recruitment (Zhang et al., 2015). Similarly, exposure to MEHP at 100–1000 mg/kg during a narrow but relevant window of exposure before parturition (GD17–19) resulted in an increase in preantral and antral follicles in F1 female mice at PND56. The premature reproductive senescence observed at the highest dose was attributed to accelerated folliculogenesis (Moyer and Hixon, 2012). It has been suggested that overactivation of PI3K signaling can most likely explain MEHP accelerated primordial follicle recruitment (Hannon et al., 2015). Exposure to DEHP (20 µg/kg/day–750 mg/kg/day) during the second half of gestation (GD10.5 – birth) resulted in decreased folliculogenesis in adult F1 ovaries (Rattan et al., 2018). In a subsequent study, numerous signaling pathways necessary for healthy ovarian function were analyzed in the F1; this study revealed disrupted expression of cell cycle regulators, PPARs, and increased *Pten* expression in the PI3K pathway (Rattan et al., 2019).

The early postnatal period of ovary development is also a window of susceptibility to DEHP. Indeed, newborn female mice injected intraperitoneally with DEHP (2.5–10 µg/g b. w./day) on PND0–4 displayed impaired germ nest breakdown and follicle assembly (Mu et al., 2015). This effect was shown to be mediated through both ER dependent and independent mechanisms (i.e., Notch 2 signaling components). Additionally, in a neonatal *in vivo* model and in an ovary culture model, DEHP exposure was shown to induce autophagy in the newborn mouse ovary and to up-regulate the expression of autophagy-related genes and the key components of AMPK-SKP2-CARM1 signaling, thus further reducing the primordial follicle pool and female fertility (Zhang et al., 2018). When the exposure took place from PND 7–14, DEHP (20 and 40 µg/kg body weight (bw)) significantly decreased the number of primordial and increased antral follicles at PND 15 in female mice (Zhang et al., 2013). Much stronger depletion of the primordial follicle pool and an increased number of secondary and antral follicles were observed at PND 21 after single day exposure to DEHP (20 and 40 µg/kg bw) on PND 5, 10, 15 and 20 (Zhang et al., 2013). The DEHP-induced acceleration in primordial follicle recruitment was proposed to be mediated by its active metabolite, MEHP, via overactivation of the PI3K signaling pathway (Hannon et al., 2015; Johansson et al., 2020). A significant increase in abnormal metaphase II spindles that may result in aneuploidy was observed in mature oocytes (Zhang et al., 2013). Further, DEHP was found to inhibit the antral follicle enlargement process in pre-pubertal mice exposed to 20 or 40 µg/kg DEHP at PND 5, 10 and 15 (Li et al., 2016a). This inhibition was linked to disturbed oxidative status and increased follicle somatic cell apoptosis caused by DEHP exposure.

Several studies provide evidence that *in utero* exposure to PBDEs may have an impact on development of the ovaries in progeny. After the treatment of Wistar rats with BDE-99 (60 or 300 µg/kg bw) on gestation day 6, the F1 female progeny that were mated to unexposed males had an increased rate of resorptions (Talsness et al., 2005). In another study, Long Evans rats exposed to BDE-99 (1 or 10 mg/kg body weight daily) from GD10 to 18 had F1 female offspring with decreased ovarian

primordial and secondary follicles (Lilienthal et al., 2006). A decrease in the numbers of secondary and antral follicles and a decrease in ovary weights was also observed in the F1 progeny of Wistar rats treated on GD6 with BDE-47 (140 or 700 µg/mL); this effect was accompanied by a reduction in serum estradiol concentrations, although ovarian aromatase activity was not affected (Talsness et al., 2008).

“Real life” exposure to environmental chemicals is to complex mixtures, yet the safety data that we have available are usually for individual compounds. When adult female Sprague-Dawley rats were exposed to a “house dust” mixture of brominated flame retardants before mating and during gestation, ovarian folliculogenesis and steroidogenesis were disrupted in the dams (F0; Lefevre et al., 2016). Interestingly, multi-oocyte follicles were observed in the ovaries of their F1 progeny, and this was accompanied by advanced puberty (Allais et al., 2020). Postnatal PBDE exposures may also affect the ovary. The treatment of female Sprague Dawley rats on PND10 with BDE-47 (10 mg/kg) decreased ovary weights and altered expression of markers for the apoptosis/stress response in the adult F1 ovary (Wang et al., 2016).

The mechanisms by which exposure to PBDEs affect ovary development are not clear. Since PBDEs, and especially their hydroxylated metabolites, are potent ligands for PPAR γ receptors, one possibility is that they disrupt PPAR γ signaling in the developing ovary (Fang et al., 2015). There is also evidence from studies with cell lines and follicle cultures that exposure to PBDEs may affect steroidogenesis in ovarian granulosa cells. Experiments with a human granulosa cell line (KGN granulosa cells) have provided evidence that exposure to the mixture of PBDEs detected in follicular fluid affects steroidogenesis and induces oxidative stress (Lefevre et al., 2016). Other studies have provided evidence that exposure to PBDEs alters steroid secretion in cells from ovarian follicles (Gregoraszczyk et al., 2008; Karpeta et al., 2011; Karpeta and Gregoraszczyk, 2010). It has been proposed that PBDE exposures may disrupt epigenetic programming during oocyte development or by affecting mitochondria and inducing oxidative stress (Sun et al., 2020).

There is mounting evidence, based on laboratory studies in rodents, that early life stage exposure to PFAS impacts the developing ovary, causing ovarian dysfunction and leading to altered folliculogenesis, abnormal hormone levels and delayed puberty in the female rodent model (Table 3, Fig. 2). For example, Zhang et al. (2020) demonstrated that *in utero* exposure of mice to PFNA, from GD 1 to 18 via maternal oral dosing (3 mg/kg bw), led to a decrease in ovary size, in the numbers of primary, secondary and antral follicles, and corpora lutea, along with delayed vaginal opening and first estrus. In addition, in these same offspring, Zhang et al. (2020) reported increased liver weight and fibroblast growth factor 21 from postnatal day 1–21 (FGF21; a PPAR-mediated hepatokine), decreased pubertal activation of kisspeptin neurons and GnRH neurons, leading to the hypothesis that these effects of PFNA were key in suppressing puberty and altering ovarian folliculogenesis. Similarly, neonatal exposure of female rats to 0.1 and 1 mg/kg/day of PFOA or to 0.1 and 10 mg/kg/day of PFOS caused a significant reduction in the number of primordial, growing and corpora lutea (Du et al., 2019). Female mice exposed *in utero* to 200 or 500 mg/kg day of perfluorobutane sulfonate (PFBS) from GD 1–20 exhibited similar effects, with reductions in all stages of ovarian follicles, in addition to decreased ovary and body size and weight, delayed vaginal opening, onset of sexual maturity and decreased serum estradiol (Feng et al., 2017). Interestingly, Feng et al. (2017) also observed decreased thyroid hormones, triiodothyronine (T3) and thyroxine (T4), in the dams on gestation day 20, as well as in the female offspring exposed *in utero*.

The mechanistic evidence for ovarian development as a target of PFAS toxicity is not clear, but one hypothesis is that it is due to the induction of oxidative stress (Ding et al., 2020). Indeed, several *in vivo* and *in vitro* experimental models have demonstrated that PFAS induces oxidative stress in various tissues via increased production of reactive oxygen species (ROS) (Wielsøe et al., 2015; Chen et al., 2017; Lu et al.,

2016; Mashayekhi et al., 2015; Suh et al., 2017). Although ROS are produced by all living organisms as a result of normal cellular metabolism, ROS-induced injury is mitigated in cells by enzymatic and non-enzymatic antioxidant systems. However, when antioxidant system capacity is exceeded due to abnormally high ROS production, damage to various cellular components may ensue (i.e. carbohydrates, nucleic acids, proteins, lipids), resulting in many pathological conditions and diseases (Birben et al., 2012; Qian et al., 2010). Some recent studies have shown that PFAS induces ROS in early stages of ovarian development, particularly, during germ cell nest formation and mitosis. For example, fetal mouse ovaries explanted on GD17, cultured for 7 days *in vitro* and then exposed for 24 h to 28.2 or 112.86 μM PFOA, exhibited a concentration-dependent increase in ROS levels as well as apoptosis and necrosis (López-Arellano et al., 2019). There is also considerable evidence to indicate that PFAS may disrupt the developing ovary by disturbing lipid metabolism via PPAR signaling pathways (Ding et al., 2020). The three known PPARs, α , β/δ and γ , are found in the mammalian ovary with α , β/δ primarily expressed in the thecal and stromal cells, and γ at higher levels in the granulosa cells and corpora lutea (Braissant et al., 1996; Komar, 2005). The PPAR α pathway plays a major role in maintaining lipid and glucose homeostasis, as well as regulating inflammatory response, cell proliferation, and differentiation (Escher and Wahli, 2000).

In summary, there is evidence that exposure to EDCs during ovarian development may delay meiotic progression and primordial follicle assembly, increase the numbers of abnormal ovarian follicles containing multiple oocytes, and alter the rate of follicle recruitment. There is a possibility that these exposures may affect the quality of the oocytes that are produced, resulting in a decrease in the rate on *in vitro* fertilisation. Steroidogenesis may also be affected. A number of signaling pathways have been implicated, including, disrupted expression of steroidogenic enzymes, disrupted PPAR signaling, disrupted PI3K pathway, cell cycle regulators, deregulation of meiosis related genes, apoptosis, and oxidative stress factors. This early life disruption of ovarian development can have long lasting effects on female reproductive health, lead to premature ovarian senescence and various comorbidities that can arise from deprivation of ovarian sex hormones, thus affecting women's overall health (Marlatt et al., this issue).

Conclusions

Based on experimental studies from different species, there is evidence that early exposure to EDCs can cause immediate change in gonad development with long-term consequences on reproductive functions. Effects range from impacts on sex determination, gonad differentiation and gonad development in fish. In mammals, *in utero* EDCs exposures do not alter sex determination or gonad differentiation; however, they can affect gonad development and steroidogenesis and may alter the programming of the germline. The discrepancy between effects observed in fish and mammals are largely due to the differences in the regulation of sex determination and timing of germline development outlined in Figs. 1 and 2. Taking into consideration that fish have many different reproductive strategies, from opportunistic (characterized by early maturation, frequent reproduction over an extended spawning season) to periodic (delayed maturation with large egg batches), while humans have a limited and relatively inefficient reproductive system (millions of spermatozoa for a single oocyte per ovulation cycle leading to a majority of singleton pregnancies), the impact of EDCs on human reproduction may carry a greater impact. Yet, the negative impact of EDCs on reproduction appears to be global in wildlife and humans (Marlatt et al. this issue).

Our review of the literature allows us to identify common mechanisms of endocrine disruption in fish and mammalian gonads; some of these are related to the importance of steroid hormone homeostasis in gonad development and function across all species. However, other signaling pathways have been shown to be important for gonadal

development and appear to be altered by EDCs, inducing various short and long-term phenotypes. Pathways related to PPAR, ERR, ROS, PI3K, AHR, RA, among others, are demonstrated targets of EDCs and may play an important role. There is also some evidence that early exposures to EDCs affect DNA methylation in mature gametes but the molecular mechanism for this is not understood. Despite progress in improving our understanding of the molecular events leading to gonadal dysgenesis after exposure to EDCs, there are still many gaps in our knowledge. In addition, other associated challenges that need to be addressed include: 1) linking mechanisms identified in experimental models to actual exposure levels requires better screening of population exposure levels; 2) the consequences of cumulative effects of exposure to multiple EDCs is not well studied; and 3) many new alternative substances that are replacing regulated EDCs are already in our environment and we have limited knowledge of their potential effects. Together, these challenges highlight the pressing need to better identify the molecular targets of EDCs and understand their mode of action.

Recently much attention has been directed to developing adverse outcome pathways (AOPs) that describe the chain of causally linked events at different levels of biological organization that lead to an adverse health or ecotoxicological effect. In the future, more emphasis should be directed to this approach as it affords the opportunity to proactively support chemical risk assessment based on mechanistic understanding of the targets and pathways affected by chemical exposure. Many of the molecular pathways controlling gonadal development are conserved across species as illustrated by the common features underpinning sex determination and gonadal differentiation in fish and mammals, including humans. Furthermore, Marlatt et al. (this issue) report consistent trends in ovarian and testicular abnormalities across adult vertebrate taxa for several well studied estrogenic EDCs, strongly supporting direct effects on EDCs on gonads and a link to organism level adverse effects on reproductive capacity. Given the common features such as the roles of estrogens and androgens and downstream regulators in mediating these processes, it is not surprising that environmental chemicals that interact with these receptors have overlapping toxicological consequences across vertebrates. The AOP framework may provide insight into targets and pathways that should be considered in conducting hazard assessments.

Credit author statement

All authors participated in the conceptualization and organization of this review. All contributed equally to the data collection and writing. GD led the project, centralized the different sections and prepared the first draft for editing by VM first followed by all the authors for final approval.

Declaration of competing interest

The authors declare that they have no known competing financial interests or personal relationships that could have appeared to influence the work reported in this paper.

Acknowledgements

The authors are grateful to the Intersectoral Centre for Endocrine Disruptor Analysis (ICEDA)'s researcher network that facilitated this Special Issue. LNM was supported by a H2020-Marie Skłodowska-Curie Action MSCA-IF-RI- 2017 awarded by the European Commission (ref. 797725-EpiSTOX).

References

- Abdel-Maksoud, F.M., Leasor, K.R., Butzen, K., Braden, T.D., Akingbemi, B.T., 2015. Prenatal exposures of male rats to the environmental chemicals bisphenol A and di (2-ethylhexyl) phthalate impact the sexual differentiation process. *Endocrinology* 156, 4672–4683. <https://doi.org/10.1210/en.2015-1077>.

- Adolfi, M.C., Herpin, A., Regensburger, M., Sacquegno, J., Waxman, J.S., Scharl, M., 2016. Retinoic acid and meiosis induction in adult versus embryonic gonads of medaka. *Sci. Rep.* 6, 1–13. <https://doi.org/10.1038/srep34281>.
- Ahmad, R., Gautam, A.K., Verma, Y., Sedha, S., Kumar, S., 2014. Effects of in utero dibutyl phthalate and butyl benzyl phthalate exposure on offspring development and male reproduction of rat. *Environ. Sci. Pollut. Res.* 21, 3156–3165. <https://doi.org/10.1007/s11356-013-2281-x>.
- Albert, O., Jégou, B., 2014. A critical assessment of the endocrine susceptibility of the human testis to phthalates from fetal life to adulthood. *Hum. Reprod. Update* 20, 231–249. <https://doi.org/10.1093/humupd/dmt050>.
- Albert, O., Nardelli, T.C., Lalancette, C., Hales, B.F., Robaire, B., 2018. Effects of in utero and lactational exposure to new generation green plasticizers on adult male rats: a comparative study with di(2-ethylhexyl) phthalate. *Toxicol. Sci.* 164, 129–141. <https://doi.org/10.1093/toxsci/kfy072>.
- Alix, M., Kjesbu, O.S., Anderson, K.C., 2020. From gametogenesis to spawning: how climate-driven warming affects teleost reproductive biology. *J. Fish. Biol.* 97, 607–632. <https://doi.org/10.1111/jfb.14439>.
- Allais, A., Albert, O., Lefèvre, P.L.C., Wade, M.G., Hales, B.F., Robaire, B., 2020. Utero and lactational exposure to flame retardants disrupts rat ovarian follicular development and advances puberty. *Toxicol. Sci.* 175, 197–209. <https://doi.org/10.1093/toxsci/kfaa044>.
- Alsop, D., Hewitt, M., Kohli, M., Brown, S., Van Der Kraak, G., 2003. Constituents within pulp mill effluent deplete retinoid stores in white sucker and bind to rainbow trout retinoic acid receptors and retinoid X receptors. *Environ. Toxicol. Chem.* 22, 2969–2976. <https://doi.org/10.1897/02-566>.
- Amir, S., Shah, S.T.A., Mamoulakis, C., Docea, A.O., Kalantzi, O.I., Zachariou, A., Calina, D., Carvalho, F., Sofikitis, N., Makrigiannakis, A., Tsatsakis, A., 2021. Endocrine disruptors acting on estrogen and androgen pathways cause reproductive disorders through multiple mechanisms: a review. *Int. J. Environ. Res. Publ. Health* 18 (4), 1464. <https://doi.org/10.3390/ijerph18041464>.
- Andrade, A.J.M., Grande, S.W., Talsness, C.E., Grote, K., Golombiewski, A., Sterner-Kock, A., Chahoud, I., 2006. A dose-response study following in utero and lactational exposure to di-(2-ethylhexyl) phthalate (DEHP): effects on androgenic status, developmental landmarks and testicular histology in male offspring rats. *Toxicology* 225, 64–74. <https://doi.org/10.1016/j.tox.2006.05.007>.
- Arbuckle, T.E., MacPherson, S., Barrett, E., Muckle, G., Séguin, J.R., Foster, W.G., Sathyanarayana, S., Dodds, L., Fisher, M., Agarwal, A., Monnier, P., Walker, M., Fraser, W.D., 2019. Do stressful life events during pregnancy modify associations between phthalates and anogenital distance in newborns? *Environ. Res.* 177, 108593. <https://doi.org/10.1016/j.envres.2019.108593>.
- Arzuaga, X., Walker, T., Yost, E.E., Radke, E.G., Hotchkiss, A.K., 2020. Use of the Adverse Outcome Pathway (AOP) framework to evaluate species concordance and human relevance of Dibutyl phthalate (DBP)-induced male reproductive toxicity. *Reprod. Toxicol.* 96, 445–458. <https://doi.org/10.1016/j.reprotox.2019.06.009>.
- Bahamonde, P.A., Munkittrick, K.R., Martyniuk, C.J., 2013. Intersex in teleost fish: are we distinguishing endocrine disruption from natural phenomena? *Gen. Comp. Endocrinol.* 192, 25–35. <https://doi.org/10.1016/j.ygcen.2013.04.005>.
- Baker, T.R., Peterson, R.E., Heideman, W., 2014. Using zebrafish as a model system for studying the transgenerational effects of dioxin. *Toxicol. Sci.* 138, 403–411. <https://doi.org/10.1093/toxsci/kfu006>.
- Balbuena, P., Campbell, J., Clewell, H.J., Clewell, R.A., 2013. Evaluation of a predictive in vitro Leydig cell assay for anti-androgenicity of phthalate esters in the rat. *Toxicol. Vitro* 27, 1711–1718. <https://doi.org/10.1016/j.tiv.2013.03.015>.
- Baumann, L., Holbech, H., Keiter, S., Kinnberg, K.L., Knörr, S., Nagel, T., Braunbeck, T., 2013. The maturity index as a tool to facilitate the interpretation of changes in vitellogenin production and sex ratio in the Fish Sexual Development Test. *Aquat. Toxicol.* 128–129, 34–42. <https://doi.org/10.1016/j.aquatox.2012.11.016>.
- Baumann, L., Knörr, S., Keiter, S., Nagel, T., Rehberger, K., Volz, S., Oberrauch, S., Schiller, V., Fenske, M., Holbech, H., Segner, H., Braunbeck, T., 2014. Persistence of endocrine disruption in zebrafish (*Danio rerio*) after discontinued exposure to the androgen 17 β -trenbolone. *Environ. Toxicol. Chem.* 33, 2488–2496. <https://doi.org/10.1002/etc.2698>.
- Bennetau-Pelissero, C., Breton B, B., Bennetau, B., Corraze, G., Le Menn, F., Davail-Cuisset, B., Helou, C., Kaushik, S.J., 2001. Effect of genistein-enriched diets on the endocrine process of gametogenesis and on reproduction efficiency of the rainbow trout *Oncorhynchus mykiss*. *Gen. Comp. Endocrinol.* 121, 173–187. <https://doi.org/10.1006/gcen.2000.7585>.
- Birben, E., Sahiner, U.M., Sackesen, C., Erzurum, S., Kalayci, O., 2012. Oxidative stress and antioxidant defense. *World Allergy Organ. J.* 5 (1), 9–19. <https://doi.org/10.1097/WOX.0b013e3182439613>.
- Blázquez, M., Medina, P., Crespo, B., Gómez, A., Zanuy, S., 2017. Identification of conserved genes triggering puberty in European sea bass males (*Dicentrarchus labrax*) by microarray expression profiling. *BMC Genom.* 18 (1), 441. <https://doi.org/10.1186/s12864-017-3823-2>.
- Borch, J., Ladefoged, O., Hass, U., Vinggaard, A.M., 2004. Steroidogenesis in fetal male rats is reduced by DEHP and DINP, but endocrine effects of DEHP are not modulated by DEHA in fetal, prepubertal and adult male rats. *Reprod. Toxicol.* 18, 53–61. <https://doi.org/10.1016/j.reprotox.2003.10.011>.
- Bowles, J., Feng, C.W., Knight, D., Smith, C.A., Roeszler, K.N., Bagheri-Fam, S., Harley, V.R., Sinclair, A.H., Koopman, P., 2008. Male-specific expression of *Aldh1a1* in mouse and chicken fetal testes: implications for retinoid balance in gonad development. *Dev. Dynam.* 238, 2073–2080. <https://doi.org/10.1002/dvdy.22024>.
- Bowles, J., Knight, D., Smith, C., Wilhelm, D., Richman, J., Mamiya, S., Yashiro, K., Chawengsaksophak, K., Wilson, M.J., Rossant, J., Hamada, H., Koopman, P., 2006. Retinoid signaling determines germ cell fate in mice. *Science* 80 (312), 596–600. <https://doi.org/10.1126/science.1125691>.
- Braissant, O., Foufelle, F., Scotto, C., Dauça, M., Wahli, W., 1996. Differential expression of peroxisome proliferator-activated receptors (PPARs): tissue distribution of PPAR- α , - β , and - γ in the adult rat. *Endocrinology* 137, 354–366. <https://doi.org/10.1210/endo.137.1.8536636>.
- Buaas, F.W., Val, P., Swain, A., 2009. The transcription co-factor CITED2 functions during sex determination and early gonad development. *Hum. Mol. Genet.* 18, 2989–3001. <https://doi.org/10.1093/hmg/ddp237>.
- Caballero-Gallardo, K., Olivero-Verbel, J., L Freeman, J., 2016. Toxicogenomics to evaluate endocrine disrupting effects of environmental chemicals using the zebrafish model. *Curr. Genom.* 17, 515–527. <https://doi.org/10.2174/1389202917666160513105959>.
- Capel, B., 2017. Vertebrate sex determination: evolutionary plasticity of a fundamental switch. *Nat. Rev. Genet.* 8 (11), 675–689. <https://doi.org/10.1038/nrg.2017.60>.
- Carrillo, M., Espigares, F., Felipe, A., Escobar, S., Molés, G., Rodríguez, R., Alvarado, M.V., Gómez, A., Zanuy, S., 2014. Updating control of puberty in male European sea bass: a holistic approach. *Gen. Comp. Endocrinol.* 221, 42–53. <https://doi.org/10.1016/j.ygcen.2015.06.019>.
- Castañeda-Cortés, D.C., Zhang, J., Boan, A.F., Langlois, V.S., Bernardino, J.I., 2020. High temperature stress response is not sexually dimorphic at the whole-body level and is dependent on androgens to induce sex reversal. *Gen. Comp. Endocrinol.* 299, 113605. <https://doi.org/10.1016/j.ygcen.2020.113605>.
- Castañeda Cortés, D.C., Arias Padilla, L.F., Langlois, V.S., Somoza, G.M., Bernardino, J.I., 2019. The central nervous system acts as a transducer of stress-induced masculinization through corticotropin-releasing hormone B. *Dev* 146 (8), dev172866. <https://doi.org/10.1242/dev.172866>.
- Chen, H., Xin, X., Liu, M., Ma, F., Yu, Y., Huang, J., Dai, H., Li, Z., Ge, R., Shan, 2021. In utero exposure to dipentyl phthalate disrupts fetal and adult Leydig cell development. *Toxicol. Appl. Pharmacol.* 419, 115514. <https://doi.org/10.1016/j.taap.2021.115514>.
- Chen, J., Xiao, Y., Gai, Z., Li, R., Zhu, Z., Bai, C., Tanguay, R.L., Xu, X., Huang, C., Dong, Q., 2015a. Reproductive toxicity of low level bisphenol A exposures in a two-generation zebrafish assay: evidence of male-specific effects. *Aquat. Toxicol.* 169, 204–214. <https://doi.org/10.1016/j.aquatox.2015.10.020>.
- Chen, P., Li, S., Liu, L., Xu, N., 2015b. Long-term effects of binary mixtures of 17 α -ethinyl estradiol and dibutyl phthalate in a partial life-cycle test with zebrafish (*Danio rerio*). *Environ. Toxicol. Chem.* 34, 518–526. <https://doi.org/10.1002/etc.2803>.
- Chen, S.R., Liu, Y.X., 2015. Regulation of spermatogonial stem cell self-renewal and spermatocyte meiosis by Sertoli cell signaling. *Reproduction* 149 (4), R159–R167. <https://doi.org/10.1530/REP-14-0481>.
- Chen, S.X., Bogerd, J., Schoonen, N.E., Martijn, J., De Waal, P.P., Schulz, R.W., 2013. A progestin (17 α ,20 β -dihydroxy-4-pregnen-3-one) stimulates early stages of spermatogenesis in zebrafish. *Gen. Comp. Endocrinol.* 185, 1–9. <https://doi.org/10.1016/j.ygcen.2013.01.005>.
- Chen, Y., Jefferson, W.N., Newbold, R.R., Padilla-Banks, E., Pepling, M.E., 2007. Estradiol, progesterone, and genistein inhibit oocyte nest breakdown and primordial follicle assembly in the neonatal mouse ovary in vitro and in vivo. *Endocrinology* 148, 3580–3590. <https://doi.org/10.1210/en.2007-0088>.
- Chen, Y., Zhou, L., Xu, J., Zhang, L., Li, M., Xie, X., Xie, Y., Luo, D., Zhang, D., Yu, X., Yang, B., Kuang, H., 2017. Maternal exposure to perfluorooctanoic acid inhibits luteal function via oxidative stress and apoptosis in pregnant mice. *Reprod. Toxicol.* 69, 159–166. <https://doi.org/10.1016/j.reprotox.2017.02.010>.
- Clelland, E., Peng, C., 2009. Endocrine/paracrine control of zebrafish ovarian development. *Mol. Cell. Endocrinol.* 312 (1–2), 42–52. <https://doi.org/10.1016/j.mce.2009.04.009>.
- Conley, J.M., Lambright, C.S., Evans, N., McCord, J., Strynar, M.J., Hill, D., Medlock-Kakaley, E., Wilson, V.S., Gray, L.E., 2021. Hexafluoropropylene oxide-dimer acid (HFPO-DA or GenX) alters maternal and fetal glucose and lipid metabolism and produces neonatal mortality, low birthweight, and hepatomegaly in the Sprague-Dawley rat. *Environ. Int.* 146, 106204. <https://doi.org/10.1016/j.envint.2020.106204>.
- Conley, J.M., Lambright, C.S., Evans, N., Strynar, M.J., McCord, J., McIntyre, B.S., Travlos, G.S., Cardon, M.C., Medlock-Kakaley, E., Hartig, P.C., Wilson, V.S., Gray, L.E., 2019. Adverse maternal, fetal, and postnatal effects of hexafluoropropylene oxide dimer acid (GenX) from oral gestational exposure in sprague-dawley rats. *Environ. Health Perspect.* 127, 37008. <https://doi.org/10.1289/EHP4372>.
- Cool, J., Defalco, T., Capel, B., 2012. Testis formation in the fetal mouse: dynamic and complex de novo tubulogenesis. *Wiley Interdiscip. Rev. Dev. Biol.* 1 (6), 847–859. <https://doi.org/10.1002/wdev.62>.
- Cosme, M.M., Lister, A.L., Van Der Kraak, G., 2015. Inhibition of spawning in zebrafish (*Danio rerio*): adverse outcome pathways of quinacrine and ethinylestradiol. *Gen. Comp. Endocrinol.* 219, 89–101. <https://doi.org/10.1016/j.ygcen.2015.01.013>.
- Crain, D.A., Janssen, S.J., Edwards, T.M., Heindel, J., Ho, S. mei, Hunt, P., Iguchi, T., Juul, A., McLachlan, J.A., Schwartz, J., Skakkebaek, N., Soto, A.M., Swan, S., Walker, C., Woodruff, T.K., Woodruff, T.J., Giudice, L.C., Guillette, L.J., 2008. Female reproductive disorders: the roles of endocrine-disrupting compounds and developmental timing. *Fertil. Steril.* 90, 911–940. <https://doi.org/10.1016/j.fertnstert.2008.08.067>.
- Crespo, D., Assis, L.H.C., Furmanek, T., Bogerd, J., Schulz, R.W., 2016. Expression profiling identifies Sertoli and Leydig cell genes as Fsh targets in adult zebrafish testes. *Mol. Cell. Endocrinol.* 437, 237–251. <https://doi.org/10.1016/j.mce.2016.08.033>.
- Culty, M., 2009. Gonocytes, the forgotten cells of the germ cell lineage. *Birth Defects Res. Part C Embryo Today - Rev.* 87 (1), 1–26. <https://doi.org/10.1002/bdrc.20142>.
- Culty, M., Thuillier, R., Li, W., Wang, Y., Martínez-Argüelles, D.B., Benjamin, C.G., Triantafyllou, K.M., Zirkin, B.R., Papadopoulos, V., 2008. In utero exposure to di-(2-ethylhexyl) phthalate exerts both short-term and long-lasting suppressive effects on

- testosterone production in the Rat I. *Biol. Reprod.* 78, 1018–1028. <https://doi.org/10.1095/biolreprod.107.065649>.
- Delbès, G., Duquenne, C., Szenker, J., Taccon, J., Habert, R., Levacher, C., 2007. Developmental changes in testicular sensitivity to estrogens throughout fetal and neonatal life. *Toxicol. Sci.* 99, 234–243. <https://doi.org/10.1093/toxsci/kfm160>.
- Denslow, N.D., Chow, M.C., Kroll, K.J., Green, L., 1999. Vitellogenin as a biomarker of exposure for estrogen or estrogen mimics. *Ecotoxicology* 8, 385–398. <https://doi.org/10.1023/A:1008986522208>.
- Devlin, R.H., Nagahama, Y., 2002. Sex determination and sex differentiation in fish: an overview of genetic, physiological, and environmental influences. *Aquaculture* 208, 191–364. [https://doi.org/10.1016/S0044-8486\(02\)00057-1](https://doi.org/10.1016/S0044-8486(02)00057-1).
- Dewing, P., Bernard, P., Vilain, E., 2002. Disorders of gonadal development. *Semin. Reprod. Med.* 20 (3), 189–198. <https://doi.org/10.1055/s-2002-35383>.
- Ding, N., Harlow, S.D., Randolph, J.F., Loch-Carus, R., Park, S.K., 2020. Perfluoroalkyl and polyfluoroalkyl substances (PFAS) and their effects on the ovary. *Hum. Reprod. Update* 26, 724–752. <https://doi.org/10.1093/humupd/dmaa018>.
- Dong, R., Yang, S., Jiao, J., Wang, T., Shi, H., Zhou, L., Zhang, Y., Wang, D., 2013. Characterization of Stra8 in Southern catfish (*Silurus meridionalis*): evidence for its role in meiotic initiation. *BMC Mol. Biol.* 14, 11. <https://doi.org/10.1186/1471-2199-14-11>.
- Dorman, D.C., Chiu, W., Hales, B.F., Hauser, R., Johnson, K.J., Mantus, E., Martel, S., Robinson, K.A., Rooney, A.A., Rudel, R., Sathyanarayana, S., Schantz, S.L., Waters, K.M., 2018. Systematic reviews and meta-analyses of human and animal evidence of prenatal diethylhexyl phthalate exposure and changes in male anogenital distance. *J. Toxicol. Environ. Health B Crit. Rev.* 21, 207–226. <https://doi.org/10.1080/10937404.2018.1505354>.
- Du, G., Hu, J., Huang, Z., Yu, M., Lu, C., Wang, X., Wu, D., 2019. Neonatal and juvenile exposure to perfluorooctanoate (PFOA) and perfluorooctane sulfonate (PFOS): advance puberty onset and kisspeptin system disturbance in female rats. *Ecotoxicol. Environ. Saf.* 167, 412–421. <https://doi.org/10.1016/j.ecoenv.2018.10.025>.
- Dutta, S., Mark-Kappeler, C.J., Hoyer, P.B., Pepling, M.E., 2014. The steroid hormone environment during primordial follicle formation in perinatal mouse ovaries. *Biol. Reprod.* 91 (3) <https://doi.org/10.1095/biolreprod.114.119214>, 68, 1–12.
- Edson, M.A., Nagaraja, A.K., Matzuk, M.M., 2009. The mammalian ovary from genesis to revelation. *Endocr. Rev.* 30 (6), 624–712. <https://doi.org/10.1210/er.2009-0012>.
- Eladak, S., Moison, D., Guerquin, M.J., Matilionyte, G., Kilcoyne, K., N'Tumba-Byn, T., Messiaen, S., Deceuninck, Y., Poggi-Gaudin, S., Benachi, A., Livera, G., Antignac, J. P., Mitchell, R., Rouiller-Fabre, V., Habert, R., 2018. Effects of environmental Bisphenol A exposures on germ cell development and Leydig cell function in the human fetal testis. *PLoS One* 31 (13), e0191934. <https://doi.org/10.1371/journal.pone.0191934>, 1.
- Enders, G.C., 1993. Sertoli-Sertoli and Sertoli-germ cell communications. *Sertoli. Cell* 447–460.
- Escher, P., Wahli, W., 2000. Peroxisome proliferator-activated receptors: insight into multiple cellular functions. *Mutat. Res. Fund. Mol. Mech. Mutagen* 448, 121–138. [https://doi.org/10.1016/S0027-5107\(99\)00231-6](https://doi.org/10.1016/S0027-5107(99)00231-6).
- Ewen-Campen, B., Schwager, E.E., Extavour, C.G.M., 2010. The molecular machinery of germ line specification. *Mol. Reprod. Dev.* 77 (1), 3–18. <https://doi.org/10.1002/mrd.21091>.
- Faheem, M., Bhandari, R.K., 2021. Detrimental effects of bisphenol compounds on physiology and reproduction in fish: a literature review. *Environ. Toxicol. Pharmacol.* 81, 103497. <https://doi.org/10.1016/j.etap.2020.103497>.
- Fang, M., Webster, T.F., Ferguson, P.L., Stapleton, H.M., 2015. Characterizing the peroxisome proliferator-activated receptor (PPAR γ) ligand binding potential of several major flame retardants, their metabolites, and chemical mixtures in house dust. *Environ. Health Perspect.* 123, 166–172. <https://doi.org/10.1289/ehp.1408522>.
- Feng, X., Cao, X., Zhao, S., Wang, X., Hua, X., Chen, Lin, Chen, Ling, 2017. Exposure of pregnant mice to perfluorobutanesulfonate causes hypothyroxinemia and developmental abnormalities in female offspring. *Toxicol. Sci.* 155, 409–419. <https://doi.org/10.1093/toxsci/kfw219>.
- Feng, X., Wang, X., Cao, X., Xia, Y., Zhou, R., Chen, L., 2015. Chronic exposure of female mice to an environmental level of perfluorooctane sulfonate suppresses estrogen synthesis through reduced histone h3k14 acetylation of the STAR promoter leading to deficits in follicular development and ovulation. *Toxicol. Sci.* 148, 368–379. <https://doi.org/10.1093/toxsci/kfv197>.
- Fent, K., 2015. Progestins as endocrine disrupters in aquatic ecosystems: concentrations, effects and risk assessment. *Environ. Int.* 84, 115–130. <https://doi.org/10.1016/j.envint.2015.06.012>.
- Fick, J., Lindberg, R.H., Parkkonen, J., Arvidsson, B., Tysklind, M., Joakim Larsson, D.G., 2010. Therapeutic levels of levonorgestrel detected in blood plasma of fish: results from screening rainbow trout exposed to treated sewage effluents. *Environ. Sci. Technol.* 44, 2661–2666. <https://doi.org/10.1021/es903440m>.
- Fisher, J.S., Macpherson, S., Marchetti, N., Sharpe, R.M., 2003. Human “testicular dysgenesis syndrome”: a possible model using in-utero exposure of the rat to dibutyl phthalate. *Hum. Reprod.* 18, 1383–1394. <https://doi.org/10.1093/humrep/deg273>.
- Franks, S., Hardy, K., 2018. Androgen action in the ovary. *Front. Endocrinol.* 9, 452. <https://doi.org/10.3389/fendo.2018.00452>.
- García-Moreno, S.A., Plebanek, M.P., Capel, B., 2018. Epigenetic regulation of male fate commitment from an initially bipotential system. *Mol. Cell. Endocrinol.* 468, 19–30. <https://doi.org/10.1016/j.mce.2018.01.009>.
- García-Reyero, N., Kroll, K.J., Liu, L., Orlando, E.F., Watanabe, K.H., Sepúlveda, M.S., Villeneuve, D.L., Perkins, E.J., Ankley, G.T., Denslow, N.D., 2009. Gene expression responses in male fathead minnows exposed to binary mixtures of an estrogen and antiestrogen. *BMC Genom.* 10, 308. <https://doi.org/10.1186/1471-2164-10-308>.
- Gemmell, N.J., Todd, E.V., Goikoetxea, A., Ortega-Recalde, O., Hore, T.A., 2019. Natural Sex Change in Fish, in: *Current Topics in Developmental Biology*. Academic Press Inc., pp. 71–117. <https://doi.org/10.1016/bs.ctdb.2018.12.014>.
- Gillman, M.W., 2005. Developmental origins of health and disease. *N. Engl. J. Med.* 353, 1848–1850. <https://doi.org/10.1056/nejme058187>.
- Goodyer, C.G., Poon, S., Aleksa, K., Hou, L., Atehortua, V., Carnevale, A., Jednak, R., Emil, S., Bagli, D., Dave, S., Hales, B.F., Chevrier, J., 2017. A case-control study of maternal polybrominated diphenyl ether (PBDE) exposure and cryptorchidism in Canadian populations. *Environ. Health Perspect.* 125 (5), 057004 <https://doi.org/10.1289/EHP522>.
- Gray, L.E., Ostby, J., Furr, J., Price, M., Veeramachaneni, D.N.R., Parks, L., 2000. Perinatal exposure to the phthalates DEHP, BBP, and DINP, but not DEP, DMP, or DOTP, alters sexual differentiation of the male rat. *Toxicol. Sci.* 58, 350–365. <https://doi.org/10.1093/toxsci/58.2.350>.
- Greeson, K.W., Fowler, K.L., Estave, P.M., Kate Thompson, S., Wagner, C., Clayton Edenfield, R., Symosko, K.M., Steves, A.N., Marder, E.M., Terrell, M.L., Barton, H., Koval, M., Marcus, M., Easley, C.A., 2020. Detrimental effects of flame retardant, PBB153, exposure on sperm and future generations. *Sci. Rep.* 10 (1), 8567. <https://doi.org/10.1038/s41598-020-65593-x>.
- Gregoraszcuk, E., Rak, A., Kawalec, K., Ropstad, E., 2008. Steroid secretion following exposure of ovarian follicular cells to single congeners and defined mixture of polybrominated dibenzoethers (PBDEs), p,p'-DDT and its metabolite p,p'-DDE. *Toxicol. Lett.* 178, 103–109. <https://doi.org/10.1016/j.toxlet.2008.02.011>.
- Griswold, M.D., Gohar, C.A., Bowles, J., Koopman, P., 2012. Initiating meiosis: the case for retinoic acid. *Biol. Reprod.* 86 (2), 35. <https://doi.org/10.1095/biolreprod.111.096610>.
- Gubbay, J., Collignon, J., Koopman, P., Capel, B., Economou, A., Münsterberg, A., Vivian, N., Goodfellow, P., Lovell-Badge, R., 1990. A gene mapping to the sex-determining region of the mouse Y chromosome is a member of a novel family of embryonically expressed genes. *Nature* 346, 245–250. <https://doi.org/10.1038/346245a0>.
- Guiguen, Y., Fostier, A., Herpin, A., 2018. Sex Determination and Differentiation in Fish, in: *Sex Control in Aquaculture*. John Wiley & Sons, Ltd, pp. 35–63. <https://doi.org/10.1002/9781119127291.ch2>.
- Guyot, R., Odet, F., Leduque, P., Forest, M.G., Magueresse-Battistoni, L., 2004. Diethylstilbestrol inhibits the expression of the Steroidogenic Acute Regulatory protein in mouse fetal testis. *Mol. Cell. Endocrinol.* 220, 67–75. <https://doi.org/10.1016/j.mce.2004.03.008>.
- Habert, R., Muczynski, V., Grisin, T., Moison, D., Messiaen, S., Frydman, R., Benachi, A., Delbes, G., Lambrot, R., Lehraiki, A., N'Tumba-Byn, T., Guerquin, M.J., Levacher, C., Rouiller-Fabre, V., Livera, G., 2014. Concerns about the widespread use of rodent models for human risk assessments of endocrine disruptors. *Reproduction* 147 (4), R119–R129. <https://doi.org/10.1530/REP-13-0497>.
- Habert, R., Picon, R., 1984. Testosterone, dihydrotestosterone and estradiol-17 β levels in maternal and fetal plasma and in fetal testes in the rat. *J. Steroid Biochem.* 21, 193–198. [https://doi.org/10.1016/0022-4731\(84\)90383-2](https://doi.org/10.1016/0022-4731(84)90383-2).
- Hales, B.F., Robaire, B., 2020. Effects of brominated and organophosphate ester flame retardants on male reproduction. *Andrology* 8 (4), 915–923. <https://doi.org/10.1111/andr.12789>.
- Hallmark, N., Walker, M., McKinnell, C., Mahood, I.K., Scott, H., Bayne, R., Coutts, S., Anderson, R.A., Greig, I., Morris, K., Sharpe, R.M., 2007. Effects of monobutyl and Di (n-dutyl) phthalate in vitro on steroidogenesis and Leydig cell aggregation in fetal testis explants from the rat: comparison with effects in vivo in the fetal rat and neonatal marmoset and in vitro in the human. *Environ. Health Perspect.* 115, 390–396. <https://doi.org/10.1289/ehp.9490>.
- Han, X., Wang, Y., Chen, T., Wilson, M.J., Pan, F., Wu, X., Rui, C., Chen, D., Tang, Q., Wu, W., 2019. Inhibition of progesterone biosynthesis induced by deca-brominated diphenyl ether (BDE-209) in mouse Leydig tumor cell (MLTC-1). *Toxicol. Vitro* 60, 383–388. <https://doi.org/10.1016/j.tiv.2019.05.016>.
- Hannon, P.R., Brannick, K.E., Wang, W., Flaws, J.A., 2015. Mono(2-ethylhexyl) phthalate accelerates early folliculogenesis and inhibits steroidogenesis in cultured mouse whole ovaries and antral follicles. *Biol. Reprod.* 92 (5), 120. <https://doi.org/10.1095/biolreprod.115.129148>.
- Hattori, R.S., Castañeda-Cortés, D.C., Arias Padilla, L.F., Strobl-Mazzulla, P.H., Ferdinando, J.I., 2020. Activation of stress response axis as a key process in environment-induced sex plasticity in fish. *Cell. Mol. Life Sci.* 77 (21), 4223–4236. <https://doi.org/10.1007/s00018-020-03532-9>.
- Hattori, R.S., Somoza, G.M., Ferdinando, J.I., Colautti, D.C., Miyoshi, K., Gong, Z., Yamamoto, Y., Strüßmann, C.A., 2019. The duplicated Y-specific amhy gene is conserved and linked to maleness in silversides of the genus *Odontesthes*. *Genes* 10 (9), 679. <https://doi.org/10.3390/genes10090679>.
- Heger, N.E., Hall, S.J., Sandrof, M.A., McDonnell, E.V., Hensley, J.B., McDowell, E.N., Martin, K.A., Gaido, K.W., Johnson, K.J., Boekelheide, K., 2012. Human fetal testis xenografts are resistant to phthalate-induced endocrine disruption. *Environ. Health Perspect.* 120, 1137–1143. <https://doi.org/10.1289/ehp.1104711>.
- Heneweer, M., Van Den Berg, M., Sanderson, J.T., 2004. A comparison of human H295R and rat R2C cell lines as in vitro screening tools for effects on aromatase. *Toxicol. Lett.* 146, 183–194. <https://doi.org/10.1016/j.toxlet.2003.10.002>.
- Hill, P.W.S., Leitch, H.G., Requena, C.E., Sun, Z., Amouroux, R., Roman-Trufero, M., Borkowska, M., Terragni, J., Vaisvila, R., Linnett, S., Bagci, H., Dharmalingham, G., Haberle, V., Lenhard, B., Zheng, Y., Pradhan, S., Hajkova, P., 2018. Epigenetic reprogramming enables the transition from primordial germ cell to gonocyte. *Nature* 555, 392–396. <https://doi.org/10.1038/nature25964>.
- Hill, R.L., Janz, D.M., 2003. Developmental estrogenic exposure in zebrafish (*Danio rerio*): I. Effects on sex ratio and breeding success. *Aquat. Toxicol.* 63, 417–429. [https://doi.org/10.1016/S0166-445X\(02\)00207-2](https://doi.org/10.1016/S0166-445X(02)00207-2).

- Ho, S.M., Cheong, A., Adgent, M.A., Veevers, J., Suen, A.A., Tam, N.N.C., Leung, Y.K., Jefferson, W.N., Williams, C.J., 2017. Environmental factors, epigenetics, and developmental origin of reproductive disorders. *Reprod. Toxicol.* 68, 85–104. <https://doi.org/10.1016/j.reprotox.2016.07.011>.
- Howdeshell, K.L., Rider, C.V., Wilson, V.S., Gray, L.E., 2008. Mechanisms of action of phthalate esters, individually and in combination, to induce abnormal reproductive development in male laboratory rats. *Environ. Res.* 108, 168–176. <https://doi.org/10.1016/j.envres.2008.08.009>.
- Hu, J., Sun, S., Guo, M., Song, H., 2014. Use of antagonists and morpholinos in loss-of-function analyses: estrogen receptor ESR2a mediates the effects of 17 alpha-ethinylestradiol on primordial germ cell distribution in zebrafish. *Reprod. Biol. Endocrinol.* 12, 40. <https://doi.org/10.1186/1477-7827-12-40>.
- Iguchi, T., Fukazawa, Y., Uesugi, Y., Takasugi, N., 1990. Polyovular follicles in mouse ovaries exposed neonatally to diethylstilbestrol in vivo and in vitro. *Biol. Reprod.* 43, 478–484. <https://doi.org/10.1095/biolreprod43.3.478>.
- Iguchi, T., Takasugi, N., 1986. Polyovular follicles in the ovary of immature mice exposed prenatally to diethylstilbestrol. *Anat. Embryol.* 175, 53–55. <https://doi.org/10.1007/BF00315455>.
- Iguchi, T., Takasugi, N., Bern, H.A., Mills, K.T., 1986. Frequent occurrence of polyovular follicles in ovaries of mice exposed neonatally to diethylstilbestrol. *Teratology* 34, 29–35. <https://doi.org/10.1002/tera.1420340105>.
- Iguchi, T., Watanabe, H., Katsu, Y., Mizutani, T., Miyagawa, S., Suzuki, A., Kohno, S., Sone, K., Kato, H., 2002. Developmental toxicity of estrogenic chemicals on rodents and other species. *Congenital. Anom.* 42 (2), 94–105. <https://doi.org/10.1111/j.1741-4520.2002.tb00858.x>.
- Iqbal, K., Tran, D.A., Li, A.X., Warden, C., Bai, A.Y., Singh, P., Wu, X., Pfeifer, G.P., Szabó, P.E., 2015. Deleterious effects of endocrine disruptors are corrected in the mammalian germline by epigenome reprogramming. *Genome Biol.* 16 (1), 59. <https://doi.org/10.1186/s13059-015-0619-z>.
- Irisawa, S., Iguchi, T., 1990. Critical period of induction by tamoxifen of genital organ abnormalities in female mice. *Vivo* 4, 175–179.
- Jensen, T.K., Frederiksen, H., Kyhl, H.B., Lassen, T.H., Swan, S.H., Bornehag, C.-G., Skakkebaek, N.E., Main, K.M., Lind, D.V., Husby, S., Andersson, A.-M., 2016. Prenatal exposure to phthalates and anogenital distance in male infants from a low-exposed Danish cohort (2010–2012). *Environ. Health Perspect.* 124, 1107–1113. <https://doi.org/10.1289/ehp.1509870>.
- Jin, Y., Chen, G., Fu, Z., 2016. Effects of TBEP on the induction of oxidative stress and endocrine disruption in Tm3 Leydig cells. *Environ. Toxicol.* 31, 1276–1286. <https://doi.org/10.1002/tox.22137>.
- Johansson, H.K.L., Damdimopolou, P., van Duursen, M.B.M., Boberg, J., Franssen, D., de Cock, M., Jäger, K., Wagner, M., Velthut-Meikas, A., Xie, Y., Connolly, L., Lelandais, P., Mazaud-Guittot, S., Salumets, A., Draskau, M.K., Filip, P., Fowler, P.A., Christiansen, S., Parent, A.S., Svingen, T., 2020. Putative adverse outcome pathways for female reproductive disorders to improve testing and regulation of chemicals. *Arch. Toxicol.* 94 (10), 3359–3379. <https://doi.org/10.1007/s00204-020-02834-y>.
- Johansson, H.K.L., Svingen, T., Fowler, P.A., Vinggaard, A.M., Boberg, J., 2017. Environmental influences on ovarian dysgenesis-developmental windows sensitive to chemical exposures. *Nat. Rev. Endocrinol.* 13 (7), 400–414. <https://doi.org/10.1038/nrendo.2017.36>.
- Johnson, K.J., Heger, N.E., Boelkeheide, K., 2012. Of mice and men (and rats): phthalate-induced fetal testis endocrine disruption is species-dependent. *Toxicol. Sci.* 129, 235–248. <https://doi.org/10.1093/toxsci/kfs206>.
- Johnson, K.J., McDowell, E.N., Viereck, M.P., Xia, J.Q., 2011. Species-specific dibutyl phthalate fetal testis endocrine disruption correlates with inhibition of SREBP2-dependent gene expression pathways. *Toxicol. Sci.* 120, 460–474. <https://doi.org/10.1093/toxsci/kfr020>.
- Jones, S., Boisvert, A., Duong, T.B., Francois, S., Thrane, P., Culty, M., 2014. Disruption of rat testis development following combined in utero exposure to the phytoestrogen genistein and antiandrogenic plasticizer di-(2-ethylhexyl) phthalate. *Biol. Reprod.* 91 (3), 64. <https://doi.org/10.1095/biolreprod.114.120907>.
- Jones, S., Boisvert, A., Francois, S., Zhang, L., Culty, M., 2015. In utero exposure to di-(2-ethylhexyl) phthalate induces testicular effects in neonatal rats that are antagonized by genistein cotreatment. *Biol. Reprod.* 93 (4), 92. <https://doi.org/10.1095/biolreprod.115.129098>.
- Kagawa, H., 2013. Oogenesis in teleost fish. *Aqua-BioScience Monogr.* 6, 99–127. <https://doi.org/10.5047/absm.2013.00604.0099>.
- Karpeta, A., Gregoraszczuk, E., 2010. Mixture of dominant PBDE congeners (BDE-47, -99, -100 and -209) at levels noted in human blood dramatically enhances progesterone secretion by ovarian follicles. *Endocr. Regul.* 44, 49–55. <https://doi.org/10.4149/endo.2010.02.49>.
- Karpeta, A., Rak-Mardyla, A., Jerzak, J., Gregoraszczuk, E.L., 2011. Congener-specific action of PBDEs on steroid secretion, CYP17, 17β-HSD and CYP19 activity and protein expression in porcine ovarian follicles. *Toxicol. Lett.* 206, 258–263. <https://doi.org/10.1016/j.toxlet.2011.08.005>.
- Kato, T., Miyata, K., Sonobe, M., Yamashita, S., Tamano, M., Miura, K., Kanai, Y., Miyamoto, S., Sakuma, T., Yamamoto, T., Inui, M., Kikusui, T., Asahara, H., Takada, S., 2013. Production of Sry knockout mouse using TALEN via oocyte injection. *Sci. Rep.* 3, 3136. <https://doi.org/10.1038/srep03136>.
- Khalil, A., Parker, M., Brown, S.E., Cevik, S.E., Guo, L.W., Jensen, J., Olmsted, A., Portman, D., Wu, H., Suvorov, A., 2017. Perinatal exposure to 2,2',4,4'-Tetrabromodiphenyl ether induces testicular toxicity in adult rats. *Toxicology* 389, 21–30. <https://doi.org/10.1016/j.tox.2017.07.006>.
- Kinnberg, K., Holbeck, H., Petersen, G.I., Bjerregaard, P., 2007. Effects of the fungicide prochloraz on the sexual development of zebrafish (*Danio rerio*). *Comp. Biochem. Physiol. C Toxicol. Pharmacol.* 145, 165–170. <https://doi.org/10.1016/j.cbpc.2006.11.002>.
- Kiparissis, Y., Balch, G.C., Metcalfe, T.L., Metcalfe, C.D., 2003. Effects of the isoflavones genistein and equol on the gonadal development of Japanese medaka (*Oryzias latipes*). *Environ. Health Perspect.* 111 (9), 1158–1163. <https://doi.org/10.1289/ehp.5928>.
- Kipp, J.L., Kilen, S.M., Bristol-Gould, S., Woodruff, T.K., Mayo, K.E., 2007. Neonatal exposure to estrogens suppresses activin expression and signaling in the mouse ovary. *Endocrinology* 148, 1968–1976. <https://doi.org/10.1210/en.2006-1083>.
- Kirigaya, A., Hayashi, S., Iguchi, T., Sato, T., 2006. Developmental effects of ethinylestradiol on reproductive organs of female mice. *Vivo* 20, 867–873.
- Komar, C.M., 2005. Peroxisome proliferator-activated receptors (PPARs) and ovarian function - implications for regulating steroidogenesis, differentiation, and tissue remodeling. *Reprod. Biol. Endocrinol.* 3, 41. <https://doi.org/10.1186/1477-7827-3-41>.
- Koubova, J., Menke, D.B., Zhou, Q., Cape, B., Griswold, M.D., Page, D.C., 2006. Retinoic acid regulates sex-specific timing of meiotic initiation in mice. *Proc. Natl. Acad. Sci. U. S. A.* 103, 2474–2479. <https://doi.org/10.1073/pnas.0510813103>.
- Kroupova, H.K., Trubiroha, A., Lorenz, C., Contardo-Jara, V., Lutz, I., Grabic, R., Kocour, M., Kloas, W., 2014. The progestin levonorgestrel disrupts gonadotropin expression and sex steroid levels in pubertal roach (*Rutilus rutilus*). *Aquat. Toxicol.* 154, 154–162. <https://doi.org/10.1016/j.aquatox.2014.05.008>.
- Kumar, V., Johnson, A.C., Trubiroha, A., Tumová, J., Ihara, M., Grabic, R., Kloas, W., Tanaka, H., Kroupová, H.K., 2015. The challenge presented by progestins in ecotoxicological research: a critical review. *Environ. Sci. Technol.* 49 (5), 2625–2638. <https://doi.org/10.1021/es5051343>.
- Kurokawa, H., Saito, D., Nakamura, S., Katoh-Fukuy, Y., Ohta, K., Baba, T., Morohashi, K. I., Tanaka, M., 2007. Germ cells are essential for sexual dimorphism in the medaka gonad. *Proc. Natl. Acad. Sci. U. S. A.* 104, 16958–16963. <https://doi.org/10.1073/pnas.0609932104>.
- Lacerda, S.M., dos, S.N., Costa, G.M.J., de França, L.R., 2014. Biology and identity of fish spermatogonial stem cell. *Gen. Comp. Endocrinol.* 207, 56–65. <https://doi.org/10.1016/j.ygcen.2014.06.018>.
- Larney, C., Bailey, T.L., Koopman, P., 2014. Switching on sex: transcriptional regulation of the testis-determining gene Sry. *Dev* 141 (11), 2195–2205. <https://doi.org/10.1242/dev.107052>.
- Lassarguère, J., Livera, G., Habert, R., Jégou, B., 2003. Time- and dose-related effects of estradiol and diethylstilbestrol on the morphology and function of the fetal rat testis in culture. *Toxicol. Sci.* 73, 160–169. <https://doi.org/10.1093/toxsci/kgf065>.
- Lawson, K.A., Dunn, N.R., Roelen, B.A.J., Zeinstra, L.M., Davis, A.M., Wright, C.V.E., Korving, J.P.W.F.M., Hogan, B.L.M., 1999. Bmp 4 is required for the generation of primordial germ cells in the mouse embryo. *Genes Dev.* 13, 424–436. <https://doi.org/10.1101/gad.13.4.424>.
- Leet, J.K., Sassman, S., Amberg, J.J., Olmstead, A.W., Lee, L.S., Ankley, G.T., Sepúlveda, M.S., 2015. Environmental hormones and their impacts on sex differentiation in fathead minnows. *Aquat. Toxicol.* 158, 98–107. <https://doi.org/10.1016/j.aquatox.2014.10.022>.
- Lefevre, P.L.C., Berger, R.G., Ernest, S.R., Gaertner, D.W., Rawn, D.F.K., Wade, M.G., Robaire, B., Hales, B.F., 2016. Exposure of female rats to an environmentally relevant mixture of brominated flame retardants targets the ovary, affecting folliculogenesis and Steroidogenesis 1. *Biol. Reprod.* 94, 1–11. <https://doi.org/10.1095/biolreprod.115.134452>.
- Lefevre, P.L.C., Wade, M., Goodyer, C., Hales, B.F., Robaire, B., 2016. A mixture reflecting polybrominated diphenyl ether (PBDE) profiles detected in human follicular fluid significantly affects steroidogenesis and induces oxidative stress in a female human granulosa cell line. *Endocrinology* 157, 2698–2711. <https://doi.org/10.1210/en.2016-1106>.
- Lessman, C.A., Brantley, N.A., 2020. Puberty visualized: sexual maturation in the transparent Casper zebrafish. *Zygote* 28 (4), 322–332. <https://doi.org/10.1017/S0967199420000180>.
- Li, C., Zou, C., Yan, H., Li, Z., Li, Y., Pan, P., Ma, F., Yu, Y., Wang, Y., Wen, Z., Ge, R.S., 2021. Perfluorodecanoic acid inhibits fetal Leydig cell differentiation after in utero exposure in rats via increasing oxidative stress and autophagy. *Environ. Toxicol.* 36, 1206–1216. <https://doi.org/10.1002/tox.23119>.
- Li, H., Papadopoulos, V., Vidic, B., Dym, M., Culty, M., 1997. Regulation of rat testis gonocyte proliferation by platelet-derived growth factor and estradiol: identification of signaling mechanisms involved. *Endocrinology* 138, 1289–1298. <https://doi.org/10.1210/endo.138.3.5021>.
- Li, L., Bu, T., Su, H., Chen, Z., Liang, Y., Zhang, G., Zhu, D., Shan, Y., Xu, R., Hu, Y., Li, J., Hu, G., Lian, Q., Ge, R.S., 2015. In utero exposure to diisononyl phthalate caused testicular dysgenesis of rat fetal testis. *Toxicol. Lett.* 232, 466–474. <https://doi.org/10.1016/j.toxlet.2014.11.024>.
- Li, L., Liu, J.-C., Lai, F.-N., Liu, H.-Q., Zhang, X.-F., Dyce, P.W., Shen, W., Chen, H., 2016a. Di-(2-ethylhexyl) phthalate exposure impairs growth of antral follicle in mice. *PLoS One* 11, e0148350. <https://doi.org/10.1371/journal.pone.0148350>.
- Li, M., Feng, R., Ma, H., Dong, R., Liu, Z., Jiang, W., Tao, W., Wang, D., 2016b. Retinoic acid triggers meiosis initiation via stra8-dependent pathway in Southern catfish, *Silurus meridionalis*. *Gen. Comp. Endocrinol.* 232, 191–198. <https://doi.org/10.1016/j.ygcen.2016.01.003>.
- Lilienthal, H., Hack, A., Roth-Härer, A., Grande, S.W., Talsness, C.E., 2006. Effects of developmental exposure to 2,2',4,4',5-pentabromodiphenyl ether (PBDE-99) on sex steroids, sexual development, and sexually dimorphic behavior in rats. *Environ. Health Perspect.* 114, 194–201. <https://doi.org/10.1289/ehp.8391>.
- Lin, L.L., Janz, D.M., 2006. Effects of binary mixtures of xenoestrogens on gonadal development and reproduction in zebrafish. *Aquat. Toxicol.* 80, 382–395. <https://doi.org/10.1016/j.aquatox.2006.10.004>.
- Liu, S., Chen, H., Xu, X.R., Hao, Q.W., Zhao, J.L., Ying, G.G., 2017. Three classes of steroids in typical freshwater aquaculture farms: comparison to marine aquaculture

- farms. *Sci. Total Environ.* 609, 942–950. <https://doi.org/10.1016/j.scitotenv.2017.07.207>.
- Liu, S., Yang, G.G., Zhao, J.L., Zhou, L.J., Yang, B., Chen, Z.F., Lai, H.J., 2012. Occurrence and fate of androgens, estrogens, glucocorticoids and progestagens in two different types of municipal wastewater treatment plants. *J. Environ. Monit.* 14, 482–491. <https://doi.org/10.1039/c1em10783f>.
- Liu, X., Matsushima, A., Shimohigashi, M., Shimohigashi, Y., 2014. A characteristic back support structure in the bisphenol A-binding pocket in the human nuclear receptor ERR γ . *PLoS One* 9, e101252. <https://doi.org/10.1371/journal.pone.0101252>.
- Liu, X., Xu, L., Shen, J., Wang, J., Ruan, W., Yu, M., Chen, J., 2016. Involvement of oxidative stress in tri-ortho-cresyl phosphate-induced autophagy of mouse Leydig TM3 cells in vitro. *Reprod. Biol. Endocrinol.* 14, 1–10. <https://doi.org/10.1186/s12958-016-0165-x>.
- Lombó, M., Getino-álvarez, L., Depincé, A., Labbé, C., Herráez, M.P., 2019. Embryonic exposure to bisphenol a impairs primordial germ cell migration without jeopardizing male breeding capacity. *Biomolecules* 9 (8), 307. <https://doi.org/10.3390/biom9080307>.
- López-Arellano, P., López-Arellano, K., Luna, J., Flores, D., Jiménez-Salazar, J., Gavia, G., Teteltitla, M., Rodríguez, J.J., Domínguez, A., Casas, E., Bahena, I., Betancourt, M., González, C., Duclomb, Y., Bonilla, E., 2019. Perfluorooctanoic acid disrupts gap junction intercellular communication and induces reactive oxygen species formation and apoptosis in mouse ovaries. *Environ. Toxicol.* 34, 92–98. <https://doi.org/10.1002/tox.22661>.
- Losa, S.M., Todd, K.L., Sullivan, A.W., Cao, J., Mickens, J.A., Patisaul, H.B., 2011. Neonatal exposure to gestinone adversely impacts the ontogeny of hypothalamic kisspeptin signaling pathways and ovarian development in the peripubertal female rat. *Reprod. Toxicol.* 31, 280–289. <https://doi.org/10.1016/j.reprotox.2010.10.002>.
- Lu, Y., Pan, Y., Sheng, N., Zhao, A.Z., Dai, J., 2016. Perfluorooctanoic acid exposure alters polyunsaturated fatty acid composition, induces oxidative stress and activates the AKT/AMPK pathway in mouse epididymis. *Chemosphere* 158, 143–153. <https://doi.org/10.1016/j.chemosphere.2016.05.071>.
- Luan, M., Liang, H., Yang, F., Yuan, W., Chen, A., Liu, X., Ji, H., Wen, S., Miao, M., 2019. Prenatal polybrominated diphenyl ethers exposure and anogenital distance in boys from a Shanghai birth cohort. *Int. J. Hyg. Environ. Health* 222, 513–523. <https://doi.org/10.1016/j.ijheh.2019.01.008>.
- Lubzens, E., Young, G., Bobe, J., Cerdá, J., 2010. Oogenesis in teleosts: how fish eggs are formed. *Gen. Comp. Endocrinol.* 165, 367–389. <https://doi.org/10.1016/j.ygcen.2009.05.022>.
- Lv, Y., Li, L., Fang, Y., Chen, P., Wu, S., Chen, X., Ni, C., Zhu, Q., Huang, T., Lian, Q., Ge, R.S., 2019. In utero exposure to bisphenol A disrupts fetal testis development in rats. *Environ. Pollut.* 246, 217–224. <https://doi.org/10.1016/j.envpol.2018.12.006>.
- Ly, L., Chan, D., Trasler, J.M., 2015. Developmental windows of susceptibility for epigenetic inheritance through the male germline. *Semin. Cell Dev. Biol.* 43, 96–105. <https://doi.org/10.1016/j.semcdb.2015.07.006>.
- Magre, S., Jost, A., 1984. Dissociation between testicular organogenesis and endocrine cytodifferentiation of Sertoli cells. *Proc. Natl. Acad. Sci. U. S. A.* 81, 7831–7834. <https://doi.org/10.1073/pnas.81.24.7831>.
- Mahood, I.K., Scott, H.M., Brown, R., Hallmark, N., Walker, M., Sharpe, R.M., 2007. In utero exposure to Di(n-butyl) phthalate and testicular dysgenesis: comparison of fetal and adult end points and their dose sensitivity. *Environ. Health Perspect.* 115, 55–61. <https://doi.org/10.1289/ehp.9366>.
- Main, K.M., Kiviranta, H., Virtanen, H.E., Sundqvist, E., Tuomisto, J.T., Tuomisto, J., Vartiainen, T., Skakkebaek, N.E., Toppari, J., 2007. Flame retardants in placenta and breast milk and cryptorchidism in newborn boys. *Environ. Health Perspect.* 115, 1519–1526. <https://doi.org/10.1289/ehp.9924>.
- Maitre, D., Selmoni, O.M., Uppal, A., Marques Da Cunha, L., Wilkins, L.G.E., Roux, J., Mobley, K.B., Castro, I., Knörr, S., Robinson-Rechavi, M., Wedekind, C., 2017. Sex differentiation in grayling (*Salmonidae*) goes through an all-male stage and is delayed in genetic males who instead grow faster. *Sci. Rep.* 7 (1), 15024. <https://doi.org/10.1038/s41598-017-14905-9>.
- Mankidy, R., Ranjan, B., Honaramooz, A., Giesy, J.P., 2014. Effects of novel brominated flame retardants on steroidogenesis in primary porcine testicular cells. *Toxicol. Lett.* 224, 141–146.
- Manku, G., Culty, M., 2015. Mammalian gonocyte and spermatogonia differentiation: recent advances and remaining challenges. *Reproduction* 49 (3), R139–R157. <https://doi.org/10.1530/REP-14-0431>.
- Manku, G., Wang, Y., Merkaoui, V., Boisvert, A., Ye, X., Blonder, J., Culty, M., 2015. Role of retinoic acid and platelet-derived growth factor receptor cross talk in the regulation of neonatal gonocyte and embryonal carcinoma cell differentiation. *Endocrinology* 156, 346–359. <https://doi.org/10.1210/en.2014-1524>.
- Mark-Kappeler, C.J., Hoyer, P.B., Devine, P.J., 2011. Xenobiotic effects on ovarian preantral follicles. *Biol. Reprod.* 85 (5), 871–883. <https://doi.org/10.1095/biolreprod.111.091173>.
- Martínez-Argüelles, D.B., Campioli, E., Culty, M., Zirkín, B.R., Papadopoulos, V., 2013. Fetal origin of endocrine dysfunction in the adult: the phthalate model. *J. Steroid Biochem. Mol. Biol.* 137, 5–17. <https://doi.org/10.1016/j.jsbmb.2013.01.007>.
- Mashayekhi, V., Tehrani, K.H.M.E., Hashemzadeh, M., Tabrizian, K., Shahraiki, J., Hosseini, M.J., 2015. Mechanistic approach for the toxic effects of perfluorooctanoic acid on isolated rat liver and brain mitochondria. *Hum. Exp. Toxicol.* 34, 985–996. <https://doi.org/10.1177/0960327114565492>.
- Matsuda, M., 2005. Sex determination in the teleost medaka, *Oryzias latipes*. *Annu. Rev. Genet.* 39, 293–307. <https://doi.org/10.1146/annurev.genet.39.110304.095800>.
- Mazón, M.J., Gómez, A., Yilmaz, O., Carrillo, M., Zanuy, S., 2014. Administration of follicle-stimulating hormone in vivo triggers testicular recrudescence of juvenile European sea bass (*Dicentrarchus labrax*). *Biol. Reprod.* 90 (1), 6. <https://doi.org/10.1095/biolreprod.113.110569>.
- McMaster, M.E., Van Der Kraak, G.J., Portt, C.B., Munkittrick, K.R., Sibley, P.K., Smith, I. R., Dixon, D.G., 1991. Changes in hepatic mixed-function oxygenase (MFO) activity, plasma steroid levels and age at maturity of a white sucker (*Catostomus commersoni*) population exposed to bleached kraft pulp mill effluent. *Aquat. Toxicol.* 21, 199–217. [https://doi.org/10.1016/0166-445X\(91\)90073-I](https://doi.org/10.1016/0166-445X(91)90073-I).
- Medina, P., Gómez, A., Zanuy, S., Blázquez, M., 2019. Involvement of the retinoic acid signaling pathway in sex differentiation and pubertal development in the European sea bass *Dicentrarchus labrax*. *Heliyon* 5 (2), e01201. <https://doi.org/10.1016/j.heliyon.2019.e01201>.
- Melo, M.C., van Dijk, P., Andersson, E., Nilsen, T.O., Fjelldal, P.G., Male, R., Nijenhuis, W., Bogerd, J., de França, L.R., Taranger, G.L., Schulz, R.W., 2015. Androgens directly stimulate spermatogonial differentiation in juvenile Atlantic salmon (*Salmo salar*). *Gen. Comp. Endocrinol.* 211, 52–61. <https://doi.org/10.1016/j.ygcen.2014.11.015>.
- Mitchell, R.T., Childs, A.J., Anderson, R.A., Van Den Driesche, S., Saunders, P.T.K., McKinnell, C., Wallace, W.H.B., Kelnar, C.J.H., Sharpe, R.M., 2012. Do phthalates affect steroidogenesis by the human fetal testis? Exposure of human fetal testis xenografts to di-n-butyl phthalate. *J. Clin. Endocrinol. Metab.* 97 (3), E341–E348. <https://doi.org/10.1210/jc.2011-2411>.
- Mitchell, R.T., Sharpe, R.M., Anderson, R.A., McKinnell, C., Macpherson, S., Smith, L.B., Wallace, W.H.B., Kelnar, C.J.H., van den Driesche, S., 2013. Diethylstilboestrol exposure does not reduce testosterone production in human fetal testis xenografts. *PLoS One* 8 (4), e61726. <https://doi.org/10.1371/journal.pone.0061726>.
- Miura, C., Higashino, T., Miura, T., 2007. A progestin and an estrogen regulate early stages of oogenesis in Fish 1. *Biol. Reprod.* 77, 822–828. <https://doi.org/10.1095/biolreprod.107.061408>.
- Miura, T., Higuchi, M., Ozaki, Y., Ohta, T., Miura, C., 2006. Progesterin is an essential factor for the initiation of the meiosis in spermatogenic cells of the eel. *Proc. Natl. Acad. Sci. U. S. A.* 103, 7333–7338. <https://doi.org/10.1073/pnas.0508419103>.
- Molés, G., Gómez, A., Carrillo, M., Zanuy, S., 2012. Development of a homologous enzyme-linked immunosorbent assay for European sea bass FSH. Reproductive cycle plasma levels in both sexes and in yearling precocious and non-precocious males. *Gen. Comp. Endocrinol.* 176, 70–78. <https://doi.org/10.1016/j.ygcen.2011.12.029>.
- Morinaga, C., Saito, D., Nakamura, S., Sasaki, T., Asakawa, S., Shimizu, N., Mitani, H., Furutani-Seiki, M., Tanaka, M., Kondoh, H., 2007. The hotel mutation of medaka in the anti-Müllerian hormone receptor causes the dysregulation of germ cell and sexual development. *Proc. Natl. Acad. Sci. U. S. A.* 104, 9691–9696. <https://doi.org/10.1073/pnas.0611379104>.
- Morthorst, J.E., Holbeck, H., Bjerregaard, P., 2010. Trenbolone causes irreversible masculinization of zebrafish at environmentally relevant concentrations. *Aquat. Toxicol.* 98, 336–343. <https://doi.org/10.1016/j.aquatox.2010.03.008>.
- Motta, P., 1997. The ultrastructure of human reproduction. 1. The natural history of the female germ cell: origin, migration and differentiation inside the developing ovary. *Hum. Reprod. Update* 3, 281–297. <https://doi.org/10.1093/humupd/3.3.281>.
- Moyer, B., Hixon, M.L., 2012. Reproductive effects in F1 adult females exposed in utero to moderate to high doses of mono-2-ethylhexylphthalate (MEHP). *Reprod. Toxicol.* 34, 43–50. <https://doi.org/10.1016/j.reprotox.2012.02.006>.
- Mu, X., Liao, X., Chen, X., Li, Y., Wang, M., Shen, C., Zhang, X., Wang, Y., Liu, X., He, J., 2015. DEHP exposure impairs mouse oocyte cyst breakdown and primordial follicle assembly through estrogen receptor-dependent and independent mechanisms. *J. Hazard Mater.* 298, 232–240. <https://doi.org/10.1016/j.jhazmat.2015.05.052>.
- Muczynski, V., Lecureuil, C., Messiaen, S., Guerin, M.J., N'Tumba-Byn, T., Moisson, D., Hodroj, W., Benjelloun, H., Baijer, J., Livera, G., Frydman, R., Benachi, A., Habert, R., Rouiller-Fabre, V., 2012. Cellular and molecular effect of MEHP involving LXR α in human fetal testis and ovary. *PLoS One* 7 (10), e48266. <https://doi.org/10.1371/journal.pone.0048266>.
- Munkittrick, K.R., McMaster, M.E., Portt, C.B., Kraak, G.J., Van Der, Smith, I.R., Dixon, D. G., 1992. Changes in maturity, plasma sex steroid levels, hepatic mixed-function oxygenase activity, and the presence of external lesions in lake whitefish (*Coregonus clupeaformis*) exposed to bleached kraft mill effluent. *Can. J. Fish. Aquat. Sci.* 49, 1560–1569. <https://doi.org/10.1139/f92-173>.
- N'Tumba-Byn, T., Moisson, D., Lacroix, M., Lecureuil, C., Lesage, L., Prud'homme, S.M., Pozzi-Gaudin, S., Frydman, R., Benachi, A., Livera, G., Rouiller-Fabre, V., Habert, R., 2012. Differential effects of bisphenol A and diethylstilbestrol on human, rat and mouse fetal Leydig cell function. *PLoS One* 7 (12), e51579. <https://doi.org/10.1371/journal.pone.0051579>.
- Naciff, J.M., Hess, K.A., Overmann, G.J., Torontali, S.M., Carr, G.J., Tiesman, J.P., Foertsch, L.M., Richardson, B.D., Martinez, J.E., Daston, G.P., 2005. Gene expression changes induced in the testis by transplacental exposure to high and low doses of 17 α -ethynyl estradiol, genistein, or bisphenol A. *Toxicol. Sci.* 86, 396–416. <https://doi.org/10.1093/toxsci/kfi198>.
- Nakamura, S., Kobayashi, K., Nishimura, T., Higashijima, S.I., Tanaka, M., 2010. Identification of germline stem cells in the ovary of the teleost medaka. *Science* (80-328), 1561–1563. <https://doi.org/10.1126/science.1185473>.
- Nardelli, T.C., Albert, O., Lalancette, C., Culty, M., Hales, B.F., Robaire, B., 2017. In utero and lactational exposure study in rats to identify replacements for di(2-ethylhexyl) phthalate. *Sci. Rep.* 7, 1–13. <https://doi.org/10.1038/s41598-017-03979-0>.
- National Toxicology Program, 2003 Apr. NTP-CERHR monograph on the potential human reproductive and developmental effects of di-n-butyl phthalate (DBP). *NTP CERHR MON* (4), i–iii90.
- Nef, S., Shipman, T., Parada, L.F., 2000. A molecular basis for estrogen-induced cryptorchidism. *Dev. Biol.* 224, 354–361. <https://doi.org/10.1006/dbio.2000.9785>.
- Nikula, H., Talonpoika, T., Kaleva, M., Toppari, J., 1999. Inhibition of hCG-stimulated steroidogenesis in cultured mouse Leydig tumor cells by bisphenol A and octylphenols. *Toxicol. Appl. Pharmacol.* 157, 166–173. <https://doi.org/10.1006/taap.1999.8674>.

- Nishimura, T., Nakamura, S., Tanaka, M., 2016. A structurally and functionally common unit in testes and ovaries of medaka (*Oryzias latipes*), a teleost fish. *Sex. Dev.* 10, 159–165. <https://doi.org/10.1159/000447313>.
- Nishimura, T., Sato, T., Yamamoto, Y., Watakabe, I., Ohkawa, Y., Suyama, M., Kobayashi, S., Tanaka, M., 2015. Foxl3 is a germ cell-intrinsic factor involved in sperm-egg fate decision in medaka. *Science* 349 (6245), 328–331. <https://doi.org/10.1126/science.aaa2657>.
- Nishimura, T., Tanaka, M., 2016. The mechanism of germline sex determination in vertebrates. *Biol. Reprod.* 95 <https://doi.org/10.1095/biolreprod.115.138271>, 30–30.
- Nourizadeh-Lillabadi, R., Lyche, J.L., Almaas, C., Stavik, B., Moe, S.J., Aleksandersen, M., Berg, V., Jakobsen, K.S., Stenseth, N.C., Utne Skåre, J., Aleström, P., Ropstad, E., 2009. Transcriptional regulation in liver and testis associated with developmental and reproductive effects in male zebrafish exposed to natural mixtures of persistent organic pollutants (POP). *J. Toxicol. Environ. Health Part A* 72, 112–130. <https://doi.org/10.1080/15287390802537255>.
- Oikawa, S., Kobayashi, S., Miyagawa, S., Iguchi, T., Sato, T., 2019. Diethylstilbestrol alters the expression of actvins in the neonatal mouse ovary in vitro. *Vivo (Brooklyn)* 33, 1095–1102. <https://doi.org/10.21873/invivo.11578>.
- Oktem, O., Urman, B., 2010. Understanding follicle growth in vivo. *Hum. Reprod.* 25 (12), 2944–2954. <https://doi.org/10.1093/humrep/deq275>.
- Okuzawa, K., 2002. Puberty in teleosts. *Fish Physiol. Biochem.* 26, 31–41. <https://doi.org/10.1023/A:1023395025374>.
- Olukole, S.G., Lanipekun, D.O., Ola-Davies, E.O., Oke, B.O., 2019. Maternal exposure to environmentally relevant doses of bisphenol A causes reproductive dysfunction in F1 adult male rats: protective role of melatonin. *Environ. Sci. Pollut. Res.* 26, 28940–28950. <https://doi.org/10.1007/s11356-019-06153-3>.
- Örn, S., Yamani, S., Norrgren, L., 2006. Comparison of vitellogenin induction, sex ratio, and gonad morphology between zebrafish and Japanese medaka after exposure to 17 α -ethinylestradiol and 17 β -trenbolone. *Arch. Environ. Contam. Toxicol.* 51, 237–243. <https://doi.org/10.1007/s00244-005-0103-y>.
- Ortega-Recalde, O., Goikoetxea, A., Hore, T.A., Todd, E.V., Gemmell, N.J., 2020. The genetics and epigenetics of sex change in fish. *Annu. Rev. Anim. Biosci.* 8, 47–69. <https://doi.org/10.1146/annurev-animal-021419-083634>.
- Padilla, L.F.A., Castañeda-Cortés, D.C., Rosa, I.F., Acosta, O.D.M., Hattori, R.S., Nóbrega, R.H., Bernardino, J.I., 2021. Cystic proliferation of germline stem cells is necessary to reproductive success and normal mating behavior in Medaka. *Elife* 10, e62757. <https://doi.org/10.7554/eLife.62757>.
- Parrott, J.A., Skinner, M.K., 1999. Kit-ligand/stem cell factor induces primordial follicle development and initiates folliculogenesis. *Endocrinology* 140, 4262–4271. <https://doi.org/10.1210/endo.140.9.6994>.
- Pasquier, J., Cabau, C., Nguyen, T., Jouanno, E., Severac, D., Braasch, I., Journot, L., Pontarotti, P., Klopp, C., Postlethwait, J.H., Guiguen, Y., Bobe, J., 2016. Gene evolution and gene expression after whole genome duplication in fish: the PhyloFish database. *BMC Genom.* 17, 368. <https://doi.org/10.1186/s12864-016-2709-z>.
- Patel, S., Zhou, C., Rattan, S., Flaws, J.A., 2015. Effects of endocrine-disrupting chemicals on the ovary. *Biol. Reprod.* 93 (1), 20. <https://doi.org/10.1095/biolreprod.115.130336>.
- Pelosi, E., Koopman, P., 2017. Development of the Testis, in: Reference Module in Biomedical Sciences. Elsevier. <https://doi.org/10.1016/b978-0-12-801238-3.99854-4>.
- Peng, C., Wang, Q., Chen, J., Yang, H., Zhang, W., Wang, D., Li, S., Tao, M., Shi, H., Lin, H., Zhao, H., Zhang, Y., 2020. Retinoic acid and androgen influence germ cells development and meiotic initiation in juvenile orange-spotted grouper, *Epinephelus coioides*. *Gen. Comp. Endocrinol.* 289, 113379. <https://doi.org/10.1016/j.ygcn.2019.113379>.
- Pepe, G.J., Billiar, R.B., Albrecht, E.D., 2006. Regulation of baboon fetal ovarian folliculogenesis by estrogen. *Mol. Cell. Endocrinol.* 247 (1–2), 41–46. <https://doi.org/10.1016/j.mce.2005.11.045>.
- Pepling, M.E., 2012. Follicular assembly: mechanisms of action. *Reproduction* 143 (2), 139–149. <https://doi.org/10.1530/REP-11-0299>.
- Pepling, M.E., De Cuevas, M., Spradling, A.C., 1999. Germline cysts: a conserved phase of germ cell development? *Trends Cell Biol.* 9 (7), 257–262. [https://doi.org/10.1016/S0962-8924\(99\)01594-9](https://doi.org/10.1016/S0962-8924(99)01594-9).
- Pepling, M.E., Spradling, A.C., 2001. Mouse ovarian germ cell cysts undergo programmed breakdown to form primordial follicles. *Dev. Biol.* 234, 339–351. <https://doi.org/10.1006/dbio.2001.0269>.
- Perez-Martinez, C., Garcia-Iglesias, M.J., Ferreras-Estrada, M.C., Bravo-Moral, A.M., Espinosa-Alvarez, J., Escudero-Diez, A., 1996. Effects of in-utero exposure to zeronol or diethylstilboestrol on morphological development of the fetal testis in mice. *J. Comp. Pathol.* 114, 407–418. [https://doi.org/10.1016/S0021-9975\(96\)80016-8](https://doi.org/10.1016/S0021-9975(96)80016-8).
- Piprek, R.P., Kloc, M., Kubiak, J.Z., 2016. Early development of the gonads: origin and differentiation of the somatic cells of the genital ridges. *Results Probl. Cell Differ.* 58, 1–22. https://doi.org/10.1007/978-3-319-31973-5_1.
- Prat, F., Sumpter, J.P., Tyler, C.R., 1996. Validation of radioimmunoassays for two salmon gonadotropins (GTH I and GTH II) and their plasma concentrations throughout the reproductive cycle in male and female rainbow trout (*Oncorhynchus mykiss*). *Biol. Reprod.* 54, 1375–1382. <https://doi.org/10.1095/biolreprod54.6.1375>.
- Qian, Y., Ducatman, A., Ward, R., Leonard, S., Bukowski, V., Guo, N.L., Shi, X., Vallyathan, V., Castranova, V., 2010. Perfluorooctane sulfonate (PFOS) induces reactive oxygen species (ROS) production in human microvascular endothelial cells: role in endothelial permeability. *J. Toxicol. Environ. Health - Part A Curr. Issues* 73, 819–836. <https://doi.org/10.1080/15287391003689317>.
- Rattan, S., Beers, H.K., Kannan, A., Ramakrishnan, A., Brehm, E., Bagchi, I., Irudayaraj, J.M.K., Flaws, J.A., 2019. Prenatal and ancestral exposure to di(2-ethylhexyl) phthalate alters gene expression and DNA methylation in mouse ovaries. *Toxicol. Appl. Pharmacol.* 379, 114629. <https://doi.org/10.1016/j.taap.2019.114629>.
- Rattan, S., Brehm, E., Gao, L., Niermann, S., Flaws, J.A., 2018. Prenatal exposure to di(2-ethylhexyl) phthalate disrupts ovarian function in a transgenerational manner in female mice. *Biol. Reprod.* 98, 130–145. <https://doi.org/10.1093/biolre/iox154>.
- Regidor, P.A., 2018. The clinical relevance of progestogens in hormonal contraception: present status and future developments. *Oncotarget* 9 (77), 34628–34638. <https://doi.org/10.18632/oncotarget.26015>.
- Rodríguez-Marí, A., Cañestro, C., BreMiller, R.A., Catchen, J.M., Yan, Y.L., Postlethwait, J.H., 2013. Retinoic acid metabolic genes, meiosis, and gonadal sex differentiation in zebrafish. *PLoS One* 8, 73951. <https://doi.org/10.1371/journal.pone.0073951>.
- Rodríguez-Marí, A., Cañestro, C., BreMiller, R.A., Nguyen-Johnson, A., Asakawa, K., Kawakami, K., Postlethwait, J.H., 2010. Sex reversal in zebrafish fancl mutants is caused by tp53-mediated germ cell apoptosis. *PLoS Genet.* 6, e1001034. <https://doi.org/10.1371/journal.pgen.1001034>.
- Rodríguez, H.A., Santambrosio, N., Santamaría, C.G., Muñoz-de-Toro, M., Luque, E.H., 2010. Neonatal exposure to bisphenol A reduces the pool of primordial follicles in the rat ovary. *Reprod. Toxicol.* 30, 550–557. <https://doi.org/10.1016/j.reprotox.2010.07.008>.
- Rolland, A.D., Lardenois, A., Goupil, A.-S., Lareyre, J.-J., Houlgatte, R., Chalmel, F., Le Gac, F., 2013. Profiling of androgen response in rainbow trout pubertal testis: relevance to male gonad development and spermatogenesis. *PLoS One* 8, e53302. <https://doi.org/10.1371/journal.pone.0053302>.
- Rouiller-Fabre, V., Guerin, M.J., N'Tumba-Byn, T., Muczynski, V., Moison, D., Tourpin, S., Messiaen, S., Habert, R., Livera, G., 2015. Nuclear receptors and endocrine disruptors in fetal and neonatal testes: a gapped landscape. *Front. Endocrinol.* 6, 58. <https://doi.org/10.3389/fendo.2015.00058>.
- Rwigemera, A., El Omri-Charai, R., Lecante, L.L., Delbes, G., 2021. Dynamics in the expression of epigenetic modifiers and histone modifications in perinatal rat germ cells during de novo DNA methylation. *Biol. Reprod.* 104, 361–373. <https://doi.org/10.1093/biolre/iaaa206>.
- Saito, D., Morinaga, C., Aoki, Y., Nakamura, S., Mitani, H., Furutani-Seiki, M., Kondoh, H., Tanaka, M., 2007. Proliferation of germ cells during gonadal sex differentiation in medaka: insights from germ cell-depleted mutant zenzai. *Dev. Biol.* 310, 280–290. <https://doi.org/10.1016/j.ydbio.2007.07.039>.
- Saitou, M., Yamaji, M., 2012. Primordial germ cells in mice. *Cold Spring Harb. Perspect. Biol.* 4, a008375. <https://doi.org/10.1101/cshperspect.a008375>.
- Santos, D., Luzio, A., Coimbra, A.M., 2017. Zebrafish sex differentiation and gonad development: a review on the impact of environmental factors. *Aquat. Toxicol.* 191, 141–163. <https://doi.org/10.1016/j.aquatox.2017.08.005>.
- Sarraj, M.A., Drummond, A.E., 2012. Mammalian foetal ovarian development: consequences for health and disease. *Reproduction*. <https://doi.org/10.1530/REP-11-0247>.
- Schang, G., Robaire, B., Hales, B.F., 2016. Organophosphate flame retardants act as endocrine-disrupting chemicals in MA-10 mouse tumor Leydig cells. *Toxicol. Sci.* 150, 499–509. <https://doi.org/10.1093/toxsci/kfw012>.
- Schulz, R.W., Bogerd, J., Male, R., Ball, J., Fenske, M., Olsen, L.C., Tyler, C.R., 2007. Estrogen-induced alterations in amh and dmrt1 expression signal for disruption in male sexual development in the zebrafish. *Environ. Sci. Technol.* 41, 6305–6310. <https://doi.org/10.1021/es070785+>.
- Schulz, R.W., de França, L.R., Lareyre, J.J., LeGac, F., Chiarini-Garcia, H., Nobrega, R.H., Miura, T., 2010. Spermatogenesis in fish. *Gen. Comp. Endocrinol.* 165, 390–411. <https://doi.org/10.1016/j.ygcn.2009.02.013>.
- Segner, H., 2011. Reproductive and Developmental Toxicity in Fishes, in: Reproductive and Developmental Toxicology. Elsevier Inc., pp. 1145–1166. <https://doi.org/10.1016/B978-0-12-382032-7.10086-4>.
- Sharpe, R.M., 2020. Androgens and the masculinization programming window: human–rodent differences. *Biochem. Soc. Trans.* 48 (4), 1725–1735. <https://doi.org/10.1042/BST20200200>.
- Sharpe, R.M., 2006. Pathways of endocrine disruption during male sexual differentiation and masculinisation. *Best Pract. Res. Clin. Endocrinol. Metab.* 20 (1), 91–110. <https://doi.org/10.1016/j.beem.2005.09.005>.
- Sharpe, R.M., Skakkebaek, N.E., 1993. Are oestrogens involved in falling sperm counts and disorders of the male reproductive tract? *Lancet* 341, 1392–1396. [https://doi.org/10.1016/0140-6736\(93\)90953-E](https://doi.org/10.1016/0140-6736(93)90953-E).
- Shima, Y., Miyabayashi, K., Haraguchi, S., Arakawa, T., Otake, H., Baba, T., Matsuzaki, S., Shishido, Y., Akiyama, H., Tachibana, T., Tsutsui, K., Morohashi, K., 2013. Contribution of Leydig and Sertoli cells to testosterone production in mouse fetal testes. *Mol. Endocrinol.* 27, 63–73. <https://doi.org/10.1210/me.2012-1256>.
- Sinclair, A.H., Berta, P., Palmer, M.S., Hawkins, J.R., Griffiths, B.L., Smith, M.J., Foster, J.W., Frischauf, A.M., Lovell-Badge, R., Goodfellow, P.N., 1990. A gene from the human sex-determining region encodes a protein with homology to a conserved DNA-binding motif. *Nature* 346, 240–244. <https://doi.org/10.1038/346240a0>.
- Skakkebaek, N.E., Rajpert-De Meys, E., Buck Louis, G.M., Toppari, J., Andersson, A.-M. M., Eisenberg, M.L., Jensen, T.K., Jørgensen, N., Swan, S.H., Sapra, K.J., Ziebe, S., Priskorn, L., Juul, A., 2016. Male reproductive disorders and fertility trends: influences of environment and genetic susceptibility. *Physiol. Rev.* 96, 55–97. <https://doi.org/10.1152/physrev.00017.2015>.
- Smith, C.A., Roessler, K.N., Bowles, J., Koopman, P., Sinclair, A.H., 2008. Onset of meiosis in the chicken embryo; evidence of a role for retinoic acid. *BMC Dev. Biol.* 8, 85. <https://doi.org/10.1186/1471-213X-8-85>.
- Song, W., Lu, H., Wu, K., Zhang, Z., Shuk-Wa Lau, E., Ge, W., 2020. Genetic evidence for estrogenicity of bisphenol A in zebrafish gonadal differentiation and its signalling mechanism. *J. Hazard Mater.* 386, 121886. <https://doi.org/10.1016/j.jhazmat.2019.121886>.

- Steves, A.N., Bradner, J.M., Fowler, K.L., Clarkson-Townsend, D., Gill, B.J., Turry, A.C., Caudle, W.M., Miller, G.W., Chan, A.W.S., Easley, C.A., 2018. Ubiquitous flame-retardant toxicants impair spermatogenesis in a human stem cell model. *iScience* 3, 161–176. <https://doi.org/10.1016/j.isci.2018.04.014>.
- Stewart, M.K., Mattiske, D.M., Pask, A.J., 2020. Estrogen suppresses SOX9 and activates markers of female development in a human testis-derived cell line. *BMC Mol. Cell Biol.* 21, 66. <https://doi.org/10.1186/s12860-020-00307-9>.
- Strüssmann, C.A., Nakamura, M., 2002. Morphology, endocrinology, and environmental modulation of gonadal sex differentiation in teleost fishes. *Fish Physiol. Biochem.* 26, 13–29. <https://doi.org/10.1023/A:1023343023556>.
- Suh, K.S., Choi, E.M., Kim, Y.J., Hong, S.M., Park, S.Y., Rhee, S.Y., Oh, S., Kim, S.W., Pak, Y.K., Choe, W., Chon, S., 2017. Perfluorooctanoic acid induces oxidative damage and mitochondrial dysfunction in pancreatic β -cells. *Mol. Med. Rep.* 15, 3871–3878. <https://doi.org/10.3892/mmr.2017.6452>.
- Sumpter, J.P., Jobling, S., 1995. Vitellogenesis as a Biomarker for Estrogen Contamination of the Aquatic Environment, in: *Environmental Health Perspectives. Public Health Services, US Dept of Health and Human Services*, pp. 173–178. <https://doi.org/10.1289/ehp.95103s7173>.
- Sun, M.H., Li, X.H., Xu, Y., Yao, X., Sun, S.C., 2020. Exposure to PBDE47 affects mouse oocyte quality via mitochondria dysfunction-induced oxidative stress and apoptosis. *Ecotoxicol. Environ. Saf.* 198, 110662. <https://doi.org/10.1016/j.ecoenv.2020.110662>.
- Sun, Y.C., Sun, X.F., Dyce, P.W., Shen, W., Chen, H., 2017. The role of germ cell loss during primordial follicle assembly: a review of current advances. *Int. J. Biol. Sci.* 13, 449–457. <https://doi.org/10.7150/ijbs.18836>.
- Suvorov, A., Shershebnv, A., Wu, H., Medvedeva, Y., Sergeev, O., Pilsner, J.R., 2018. Perinatal exposure to low dose 2,2',4,4'-tetrabromodiphenyl ether (BDE47) alters sperm DNA methylation in adult rats. *Reprod. Toxicol.* 75, 136–143. <https://doi.org/10.1016/j.reprotox.2017.10.009>.
- Suzuki, A., Sugihara, A., Uchida, K., Sato, T., Ohta, Y., Katsu, Y., Watanabe, H., Iguchi, T., 2002. Developmental effects of perinatal exposure to bisphenol-A and diethylstilbestrol on reproductive organs in female mice. *Reprod. Toxicol.* 16, 107–116. [https://doi.org/10.1016/S0890-6238\(02\)00005-9](https://doi.org/10.1016/S0890-6238(02)00005-9).
- Svensson, J., Mustafa, A., Fick, J., Schmitz, M., Brunström, B., 2016. Developmental exposure to progesterins causes male bias and precocious puberty in zebrafish (*Danio rerio*). *Aquat. Toxicol.* 177, 316–323. <https://doi.org/10.1016/j.aquatox.2016.06.010>.
- Svingen, T., Koopman, P., 2013. Building the mammalian testis: origins, differentiation, and assembly of the component cell populations. *Genes Dev.* 27 (22), 2409–2426. <https://doi.org/10.1101/gad.228080.113>.
- Swan, S.H., Main, K.M., Liu, F., Stewart, S.L., Kruse, R.L., Calafat, A.M., Mao, C.S., Redmon, J.B., Ternand, C.L., Sullivan, S., Teague, J.L., Drobnis, E.Z., Carter, B.S., Kelly, D., Simmons, T.M., Wang, C., Lumbrears, L., Villanueva, S., Diaz-Romero, M., Lomeli, M.B., Otero-Salazar, E., Hobel, C., Brock, B., Kwong, C., Muehlen, A., Sparks, A., Wolf, A., Whitham, J., Hatterman-Zogg, M., Maifeld, M., 2005. Decrease in anogenital distance among male infants with prenatal phthalate exposure. *Environ. Health Perspect.* 113, 1056–1061. <https://doi.org/10.1289/ehp.8100>.
- Swan, S.H., Sathyanarayana, S., Barrett, E.S., Janssen, S., Liu, F., Nguyen, R.H.N., Redmon, J.B., 2015. First trimester phthalate exposure and anogenital distance in newborns. *Hum. Reprod.* 30, 963–972. <https://doi.org/10.1093/humrep/deu363>.
- Tainaka, H., Takahashi, H., Umezawa, M., Tanaka, H., Nishimune, Y., Oshio, S., Takeda, K., 2012. Evaluation of the testicular toxicity of prenatal exposure to bisphenol A based on microarray analysis combined with MeSH annotation. *J. Toxicol. Sci.* 37, 539–548. <https://doi.org/10.2131/jts.37.539>.
- Takatsu Takatsu, K., Miyaoku, K., Roy, S.R., Murono, Y., Sago, T., Itagaki, H., Nakamura, M., Tokumoto, T., 2013. Induction of female-to-male sex change in adult zebrafish by aromatase inhibitor treatment. *Sci. Rep.* 3, 3400. <https://doi.org/10.1038/srep03400>.
- Talsness, C.E., Kuriyama, S.N., Sterner-Kock, A., Schnitker, P., Grande, S.W., Shakibaei, M., Andrade, A., Grote, K., Chahoud, I., 2008. In Utero and lactational exposures to low doses of polybrominated diphenyl ether-47 alter the reproductive system and thyroid gland of female rat offspring. *Environ. Health Perspect.* 116, 308–314. <https://doi.org/10.1289/ehp.10536>.
- Talsness, C.E., Shakibaei, M., Kuriyama, S.N., Grande, S.W., Sterner-Kock, A., Schnitker, P., De Souza, C., Grote, K., Chahoud, I., 2005. Ultrastructural changes observed in rat ovaries following in utero and lactational exposure to low doses of a polybrominated flame retardant. *Toxicol. Lett.* 157, 189–202. <https://doi.org/10.1016/j.toxlet.2005.02.001>.
- Tanaka, M., 2019. Regulation of Germ Cell Sex Identity in Medaka, in: *Current Topics in Developmental Biology*. Academic Press Inc., pp. 151–165. <https://doi.org/10.1016/bs.ctdb.2019.01.010>.
- Tanaka, M., Nakaya, S., Katayama, M., Leffers, H., Nozawa, S., Nakazawa, R., Iwamoto, T., Kobayashi, S., 2006. Effect of prenatal exposure to bisphenol A on the serum testosterone concentration of rats at birth. *Hum. Exp. Toxicol.* 25, 369–373. <https://doi.org/10.1191/0960327106ht638oa>.
- Taranger, G.L., Carrillo, M., Schulz, R.W., Fontaine, P., Zanuy, S., Felip, A., Weltzien, F. A., Dufour, S., Karlsen, Ø., Norberg, B., Andersson, E., Hansen, T., 2010. Control of puberty in farmed fish. *Gen. Comp. Endocrinol.* 165, 483–515. <https://doi.org/10.1016/j.ygcen.2009.05.004>.
- Tarapore, P., Ouyang, B., 2021. Perfluoroalkyl chemicals and male reproductive health: do pfoa and pfos increase risk for male infertility? *Int. J. Environ. Res. Publ. Health* 18 (7), 3794. <https://doi.org/10.3390/ijerph18073794>.
- Thayer, K.A., Ruhlen, R.L., Howdeshell, K.L., Buchanan, D.L., Cooke, P.S., Preziosi, D., Welshons, W.V., Haseman, J., Vom Saal, F.S., 2001. Altered prostate growth and daily sperm production in male mice exposed prenatally to subclinical doses of 17 α -ethynyl oestradiol. *Hum. Reprod.* 16, 988–996. <https://doi.org/10.1093/humrep/16.5.988>.
- Thuillier, R., Manku, G., Wang, Y., Culty, M., 2009. Changes in MAPK pathway in neonatal and adult testis following fetal estrogen exposure and effects on rat testicular cells. *Microsc. Res. Tech.* 72, 773–786. <https://doi.org/10.1002/jemt.20756>.
- Thuillier, R., Wang, Y., Culty, M., 2003. Prenatal exposure to estrogenic compounds alters the expression pattern of platelet-derived growth factor receptors α and β in neonatal rat testis: identification of gonocytes as targets of estrogen Exposure 1. *Biol. Reprod.* 68, 867–880. <https://doi.org/10.1095/biolreprod.102.009605>.
- Tokumoto, T., Tokumoto, M., Horiguchi, R., Ishikawa, K., Nagahama, Y., 2004. Diethylstilbestrol induces fish oocyte maturation. *Proc. Natl. Acad. Sci. U. S. A.* 101, 3686–3690. <https://doi.org/10.1073/pnas.0400072101>.
- Tokumoto, T., Tokumoto, M., Nagahama, Y., 2005. Induction and inhibition of oocyte maturation by EDCs in zebrafish. *Reprod. Biol. Endocrinol.* 3, 69. <https://doi.org/10.1186/1477-7827-3-69>.
- van den Driesche, S., Kilcoyne, K.R., Wagner, I., Rebourcet, D., Boyle, A., Mitchell, R., McKinnell, C., Macpherson, S., Donat, R., Shukla, C.J., Jorgensen, A., Meyts, E.R. De, Skakkebaek, N.E., Sharpe, R.M., 2017. Experimentally induced testicular dysgenesis syndrome originates in the masculinization programming window. *JCI insight* 2, e91204. <https://doi.org/10.1172/jci.insight.91204>.
- van den Driesche, S., Kolovos, P., Platts, S., Drake, A.J., Sharpe, R.M., 2012. Inter-relationship between testicular dysgenesis and Leydig cell function in the masculinization programming window in the rat. *PLoS One* 7, e30111. <https://doi.org/10.1371/journal.pone.0030111>.
- van den Driesche, S., McKinnell, C., Calarrão, A., Kennedy, L., Hutchison, G.R., Hrabalkova, L., Jobling, M.S., Macpherson, S., Anderson, R.A., Sharpe, R.M., Mitchell, R.T., 2015. Comparative effects of di(*n*-butyl) phthalate exposure on fetal germ cell development in the rat and in human fetal testis xenografts. *Environ. Health Perspect.* 123, 223–230. <https://doi.org/10.1289/ehp.1408248>.
- Walker, C., Garza, S., Papadopoulos, V., Culty, M., 2021. Impact of endocrine-disrupting chemicals on steroidogenesis and consequences on testicular function. *Mol. Cell. Endocrinol.* 527, 112125. <https://doi.org/10.1016/j.mce.2021.112125>.
- Wallacides, A., Chesnel, A., Chardard, D., Flament, S., Dumond, H., 2009. Evidence for a conserved role of retinoic acid in urodele amphibian meiosis onset. *Dev. Dynam.* 238, 1389–1398. <https://doi.org/10.1002/dvdy.21948>.
- Walsh, L.P., McCormick, C., Martin, C., Stocco, D.M., 2000. Roundup inhibits steroidogenesis by disrupting steroidogenic acute regulatory (STAR) protein expression. *Environ. Health Perspect.* 108, 769–776. <https://doi.org/10.1289/ehp.00108769>.
- Wang, C., Zhang, S., Ma, R., Zhang, X., Zhang, C., Li, B., Niu, Q., Chen, J., Xia, T., Li, P., Zhao, Q., Dong, L., Xu, C., Wang, A., 2016. Roles of endoplasmic reticulum stress, apoptosis and autophagy in 2,2',4,4'-tetrabromodiphenyl ether-induced rat ovarian injury. *Reprod. Toxicol.* 65, 187–193. <https://doi.org/10.1016/j.reprotox.2016.07.013>.
- Wang, C., Zhou, B., Xia, G., 2017. Mechanisms controlling germline cyst breakdown and primordial follicle formation. *Cell. Mol. Life Sci.* 74 (14), 2547–2566. <https://doi.org/10.1007/s00018-017-2480-6>.
- Wang, J.J., Tian, Y., Li, M.H., Feng, Y.Q., Kong, L., Zhang, F.L., Shen, W., 2021. Single-cell transcriptome dissection of the toxic impact of Di (2-ethylhexyl) phthalate on primordial follicle assembly. *Theranostics* 11, 4992–5009. <https://doi.org/10.7150/thno.55006>.
- Wang, Y., Culty, M., 2007. Identification and distribution of a novel platelet-derived growth factor receptor β variant: effect of retinoic acid and involvement in cell differentiation. *Endocrinology* 148, 2233–2250. <https://doi.org/10.1210/en.2006-1206>.
- Wei, Y., Han, C., Geng, Y., Cui, Y., Bao, Y., Shi, W., Zhong, X., 2019. Maternal exposure to bisphenol A during pregnancy interferes testis development of F1 male mice. *Environ. Sci. Pollut. Res.* 26, 23491–23504. <https://doi.org/10.1007/s11356-019-05579-z>.
- Weidinger, G., Wolke, U., Köprunner, M., Thisse, C., Thisse, B., Raz, E., 2002. Regulation of zebrafish primordial germ cell migration by attraction towards an intermediate target. *Development* 129, 25–36.
- Welsh, M., Saunders, P.T.K., Finken, M., Scott, H.M., Hutchison, G.R., Smith, L.B., Sharpe, R.M., 2008. Identification in rats of a programming window for reproductive tract masculinization, disruption of which leads to hypospadias and cryptorchidism. *J. Clin. Invest.* 118, 1479–1490. <https://doi.org/10.1172/JCI34241>.
- WHO, 2002. *Global Assessment of the State-Of-The-Science of Endocrine Disruptors*.
- Wielsoe, M., Long, M., Ghisari, M., Bonefeld-Jørgensen, E.C., 2015. Perfluoroalkylated substances (PFAS) affect oxidative stress biomarkers in vitro. *Chemosphere* 129, 239–245. <https://doi.org/10.1016/j.chemosphere.2014.10.014>.
- Williams, C.J., Erickson, G.F., 2012. Morphology and physiology of the ovary. In: Feingold, K.R., Anawalt, B., Boyce, A., Chrousos, G., de Herder, W.W., Dhatariya, K., Dungan, K., Grossman, A., Hershman, J.M., Hofland, J., Kalra, S., Kalsats, G., Koch, G., Kopp, P., Korbonits, M., Kovacs, C.S., Kuohung, W., Laferrère, B., McGee, E. A., McLachlan, R., Morley, J.E., New, M., Purnell, J., Sahay, R., Singer, F., Stratakis, C.A., Trencle, D.L., Wilson, D.P. (Eds.), *Endotext* [Internet]. South Dartmouth (MA): MDText.Com, Inc.; 2000.
- Wohlfahrt-Veje, C., Main, K.M., Skakkebaek, N.E., 2009. Testicular dysgenesis syndrome: foetal origin of adult reproductive problems. *Clin. Endocrinol.* 71 (4), 459–465. <https://doi.org/10.1111/j.1365-2265.2009.03545.x>.
- Wu, H., Hauser, R., Krawetz, S.A., Pilsner, J.R., 2015. Environmental susceptibility of the sperm epigenome during windows of male germ cell development. *Curr. Environ. Heal. reports* 2, 356–366. <https://doi.org/10.1007/s40572-015-0067-7>.

- Wu, N., Yu, A.B., Zhu, H. Bin, Lin, X.K., 2012. Effective silencing of Sry gene with RNA interference in developing mouse embryos resulted in feminization of XY gonad. *J. Biomed. Biotechnol.* 343891. <https://doi.org/10.1155/2012/343891>, 2012.
- Xiong, Y., Wang, S., Gui, J.F., Mei, J., 2020. Artificially induced sex-reversal leads to transition from genetic to temperature-dependent sex determination in fish species. *Sci. China Life Sci.* 63 (1), 157–159. <https://doi.org/10.1007/s11427-019-1568-7>.
- Yamaguchi, T., Yoshinaga, N., Yazawa, T., Gen, K., Kitano, T., 2010. Cortisol is involved in temperature-dependent sex determination in the Japanese flounder. *Endocrinology* 151, 3900–3908. <https://doi.org/10.1210/en.2010-0228>.
- Yamamoto, M., Shirai, M., Sugita, K., Nagai, N., Miura, Y., Mogi, R., Yamamoto, K., Tamura, A., Arishima, K., 2003. Effects of maternal exposure to diethylstilbestrol on the development of the reproductive system and thyroid function in male and female rat offspring. *J. Toxicol. Sci.* 28, 385–394. <https://doi.org/10.2131/jts.28.385>.
- Yamamoto, Y., Zhang, Y., Sarida, M., Hattori, R.S., Strüssmann, C.A., 2014. Coexistence of genotypic and temperature-dependent sex determination in pejerrey *odonates bonariensis*. *PLoS One* 9, e102574. <https://doi.org/10.1371/journal.pone.0102574>.
- Yang, Q., Sui, X., Cao, J., Liu, C., Zheng, S., Bao, M., Huang, Y., Wu, K., 2019. Effects of exposure to bisphenol a during pregnancy on the pup testis function. *Internet J. Endocrinol.* 21, 6785289. <https://doi.org/10.1155/2019/6785289>.
- Yang, Q., Yang, X., Liu, J., Chen, Y., Shen, S., 2018. Effects of exposure to BPF on development and sexual differentiation during early life stages of zebrafish (*Danio rerio*). *Comp. Biochem. Physiol. C Toxicol. Pharmacol.* 210, 44–56. <https://doi.org/10.1016/j.cbpc.2018.05.004>.
- Yasuda, Y., Kihara, T., Tanimura, T., Nishimura, H., 1985. Gonadal dysgenesis induced by prenatal exposure to ethinyl estradiol in mice. *Teratology* 32, 219–227. <https://doi.org/10.1002/tera.1420320210>.
- Yasuda, Y., Ohara, I., Konishi, H., Tanimura, T., 1988. Long-term effects on male reproductive organs of prenatal exposure to ethinyl estradiol. *Am. J. Obstet. Gynecol.* 159, 1246–1250. [https://doi.org/10.1016/0002-9378\(88\)90458-9](https://doi.org/10.1016/0002-9378(88)90458-9).
- Ye, T., Kang, M., Huang, Q., Fang, C., Chen, Y., Shen, H., Dong, S., 2014. Exposure to DEHP and MEHP from hatching to adulthood causes reproductive dysfunction and endocrine disruption in marine medaka (*Oryzias melastigma*). *Aquat. Toxicol.* 146, 115–126. <https://doi.org/10.1016/j.aquatox.2013.10.025>.
- Ying, Y., Liu, X.-M., Marble, A., Lawson, K.A., Zhao, G.-Q., 2000. Requirement of Bmp8b for the generation of primordial germ cells in the mouse. *Mol. Endocrinol.* 14, 1053–1063. <https://doi.org/10.1210/mend.14.7.0479>.
- Yoshizaki, G., Yazawa, R., 2019. Application of surrogate broodstock technology in aquaculture. *Fish. Sci.* 85, 429–437. <https://doi.org/10.1007/s12562-019-01299-y>.
- Zama, A.M., Uzumcu, M., 2010. Epigenetic effects of endocrine-disrupting chemicals on female reproduction: an ovarian perspective. *Front. Neuroendocrinol.* 31 (4), 420–439. <https://doi.org/10.1016/j.yfrne.2010.06.003>.
- Zarean, M., Keikha, M., Feizi, A., Kazemitabae, M., Kelishadi, R., 2019. The role of exposure to phthalates in variations of anogenital distance: a systematic review and meta-analysis. *Environ. Pollut.* 247, 172–179. <https://doi.org/10.1016/j.envpol.2019.01.026>.
- Zawatski, W., Lee, M.M., 2013. Male pubertal development: are endocrine-disrupting compounds shifting the norms? *J. Endocrinol.* 218 (2), R1–R12. <https://doi.org/10.1530/JOE-12-0449>.
- Zhang, X., Tang, S., Qiu, T., Hu, X., Lu, Y., Du, P., Xie, L., Yang, Y., Zhao, F., Zhu, Y., Giesy, J.P., 2020. Investigation of phthalate metabolites in urine and daily phthalate intakes among three age groups in Beijing, China. *Environ. Pollut.* 260, 114005. <https://doi.org/10.1016/j.envpol.2020.114005>.
- Zhang, X.F., Zhang, L.J., Li, L., Feng, Y.N., Chen, B., Ma, J.M., Huynh, E., Shi, Q.H., De Felici, M., Shen, W., 2013. Diethylhexyl phthalate exposure impairs follicular development and affects oocyte maturation in the mouse. *Environ. Mol. Mutagen.* 54, 354–361. <https://doi.org/10.1002/em.21776>.
- Zhang, X.F., Zhang, T., Han, Z., Liu, J.C., Liu, Y.P., Ma, J.Y., Li, L., Shen, W., 2015. Transgenerational inheritance of ovarian development deficiency induced by maternal diethylhexyl phthalate exposure. *Reprod. Fertil. Dev.* 27, 1213–1221. <https://doi.org/10.1071/RD14113>.
- Zhang, Y., Mu, X., Gao, R., Geng, Y., Liu, X., Chen, X., Wang, Yuheng, Ding, Y., Wang, Yingxiong, He, J., 2018. Foetal-neonatal exposure of Di (2-ethylhexyl) phthalate disrupts ovarian development in mice by inducing autophagy. *J. Hazard Mater.* 358, 101–112. <https://doi.org/10.1016/j.jhazmat.2018.06.042>.
- Zhu, Q., Li, H., Wen, Z., Wang, Y., Li, X., Huang, T., Mo, J., Wu, Y., Zhong, Y., Ge, R.S., 2020. Perfluoroalkyl substances cause Leydig cell dysfunction as endocrine disruptors. *Chemosphere* 253, 126764. <https://doi.org/10.1016/j.chemosphere.2020.126764>.