

Effects of atrazine and S-metolachlor on stream periphyton taxonomic and fatty acid compositions

Laura Malbezin^{1*}, Soizic Morin² and Isabelle Lavoie¹

¹Institut national de la recherche scientifique, centre Eau Terre Environnement, 490 rue de la Couronne G1K 9A9, Quebec City, QC, Canada

²INRAE, EABX, 50 avenue de Verdun 33612 Cestas Cedex, France

* Corresponding author: lmalbezin@outlook.fr

ORCID ID:

0000-0002-4544-7952 (Laura Malbezin)

0000-0003-0360-9383 (Soizic Morin)

0000-0002-2918-6297 (Isabelle Lavoie)

Abstract:

Extensive pesticide use for agriculture can diffusely pollute aquatic ecosystems through leaching and runoff events and has the potential to negatively affect non-target organisms. Atrazine and S-metolachlor are two widely used herbicides often detected in high concentrations in rivers that drain nearby agricultural lands. Previous studies focused on concentration-response exposure of algal monospecific cultures, over a short exposure period, with classical descriptors such as cell density, mortality or photosynthetic efficiency as response variables. In this study, we exposed algal biofilms (periphyton) to a concentration gradient of atrazine and S-metolachlor for 14 days. We focused on fatty acid composition as the main concentration-response descriptor, and we also measured chlorophyll a fluorescence. Results showed that atrazine increased cyanobacteria and diatom chlorophyll a fluorescence. Both herbicides caused dissimilarities in fatty acid profiles between control and high exposure concentrations, but S-metolachlor had a stronger effect than atrazine on the observed increase or reduction in saturated fatty acids (SFAs) and very long chain fatty acids (VLCFAs), respectively. Our study demonstrates that two commonly used herbicides, atrazine and S-metolachlor, can negatively affect the taxonomic composition and fatty acid profiles of stream periphyton, thereby altering the nutritional quality of this resource for primary consumers.

Keywords:

Periphyton, Herbicides, Atrazine, S-metolachlor, Fatty acids, Fluorescence

Acknowledgments:

The authors would like to thank Stéphane Moïse from the general laboratory at INRS-ETE for his help on herbicide analysis. We would also like to thank Nolan Pearce for English revisions as well as the Groupe de recherche interuniversitaire en limnologie (GRIL).

37 1. Introduction

38

39 In 2020, worldwide pesticide use in agriculture was estimated at 2.7 million tons (FAO, 2022). The application
40 of these compounds on the landscape has resulted in the detection and persistence of pesticides in aquatic
41 ecosystems. Even at low concentrations, pesticides can interact with other compounds and represent a serious risk
42 to aquatic and terrestrial organisms (Groner and Relyea, 2011; Relyea, 2009). Pesticides that target autotrophs
43 (i.e., herbicides) represent about 48% of the pesticides used globally, and they may comprise an even more
44 substantial proportion, ranging from 63% to upwards of 80% in certain regions of the world such as in the United
45 States of America (USA) (Brain and Anderson, 2019; USEPA, 2017). Atrazine and S-metolachlor are two
46 herbicides commonly applied for grain, legume and cereal crop production. Resultantly, these herbicides are
47 frequently detected in nearby aquatic ecosystems with mean concentrations close to 1 $\mu\text{g.L}^{-1}$ in surface waters in
48 Argentina and in the USA (Bachetti et al., 2021; Hansen et al., 2019). Atrazine daily maximum concentrations
49 reached hundreds $\mu\text{g.L}^{-1}$ in watersheds highly vulnerable to runoff in agricultural regions of the USA (see Perkins
50 et al., 2021 for complete database) and can exceed water quality criteria in Europe (Parlakidis et al., 2022; Székács
51 et al., 2015). S-metolachlor is also commonly applied for corn and soybean production, and can reach
52 concentrations between 5 $\mu\text{g.L}^{-1}$ and 50 $\mu\text{g.L}^{-1}$ in agricultural regions of Europe (Griffini et al., 1997; Kapsi et al.,
53 2019; Roubex et al., 2012; Székács et al., 2015; Vryzas et al., 2011), and up to 100 $\mu\text{g.L}^{-1}$ in agricultural regions
54 of the USA (Battaglin et al., 2003, 2000).

55 Atrazine [2-chloro-4-(ethylamino)-6-(isopropylamino)-s-triazine] is a triazine compound marketed in the late
56 1950s but has been subsequently severely restricted in Europe since 2003-2004 (European Commission, 2004)
57 and banned in certain countries (e.g., France, Sénat de France, 2003, and Germany, LAWA, 2019) due to its
58 presence at concentrations beyond water quality criteria. Although studies and government reports mentioned the
59 potential risk of this molecule on non-target terrestrial and aquatic organisms (de Albuquerque et al., 2020;
60 USEPA, 2016), atrazine is still used in several countries worldwide, including in Canada and in the USA, albeit
61 under increased regulation (e.g., Quebec, see Fortier, 2018). Atrazine is a photosynthesis inhibitor herbicide that
62 binds the D1 protein of photosystem II and blocks electron transport (Valotton et al., 2008). By disrupting electron
63 transport, atrazine leads to the production of reactive oxygen species (ROS), resulting to oxidative stress,
64 peroxidation of membrane lipids and, ultimately, senescence of plant cells (de Albuquerque et al., 2020). When
65 present in aquatic ecosystems, atrazine can be harmful for aquatic plants (Gao et al., 2019), micro-algae (Baxter
66 et al., 2016), as well as non-phototrophic organisms such as bacteria (DeLorenzo et al., 1999). The effects of
67 atrazine on amphibians, in particular, have long been disputed, but the USEPA mentioned a potential chronic risk
68 to amphibians, fish, and aquatic invertebrates in locations where atrazine use is heaviest (USEPA, 2016).

69 S-metolachlor (2-chloro-N-(2-ethyl-6-methylphenyl)-N-[(1S)-2-methoxy-1-methylethyl] acetamide) is an
70 extensively used chloroacetamide herbicide available since the 1990s. S-metolachlor inhibits very long chain fatty
71 acids (VLCFAs) biosynthesis by binding with a synthase involved in fatty acid elongation (HRAC, 2020; WSSA,
72 2021). VLCFAs are an important component for the functioning of biological membranes. For example, Böger et
73 al. (2003) found that S-metolachlor inhibited 68% of VLCFAs biosynthesis in the green algae *Scenedesmus acutus*
74 compared to control. Similarly, Debenest et al. (2009) found that this compound can directly affect cellular density
75 of periphytic diatoms. In addition, S-metolachlor is highly soluble, mobile, can bioaccumulate in non-target

76 organisms (Zemolin et al., 2014), and it is suspected to be an endocrine disruptor for certain fish species (Ou-Yang
77 et al., 2022; Quintaneiro et al., 2017).

78 Freshwater biofilms or periphyton are a heterogeneous assemblage of algae, bacteria, fungi, archaea and
79 viruses as well as micromeiofauna trapped in a matrix of extracellular polymeric substances that develop on
80 various submerged substrates (Wetzel, 1983). Periphyton is an integral part to the function of aquatic ecosystems
81 and provides services in nutrient cycling. In addition, it is the basal resource of aquatic food webs providing
82 essential compounds such as proteins, lipids and fatty acids needed for the growth and metabolism of higher trophic
83 levels (Thompson et al., 2002). Fatty acids (FAs), in particular, are an important compound transferred along the
84 food chain from prey to consumers (Gladyshev et al., 2011). Polyunsaturated fatty acids (PUFAs) are involved in
85 physiological processes and maintain membrane structure (Huggins et al., 2004). While vegetal cells can
86 synthesize PUFAs *de novo*, consumers must obtain them through dietary pathways (Brett and Müller-Navarra,
87 1997). In particular, certain essential FAs such as linoleic acid (LIN; C18:2n6) and α -linoleic acid (ALA; C18:3n3)
88 are almost exclusively produced by vegetal cells; therefore, algae represent an essential source of these molecules
89 for animal consumers (Brett and Müller-Navarra, 1997). In aquatic ecosystems, long-chain PUFAs (LCPUFAs)
90 such as arachidonic acid (ARA; C20:4n6), eicosapentanoic acid (EPA; C20:5n3) and docosahexanoic acid (DHA;
91 C22:6n3) are also mainly produced by microalgae (Li et al., 2014) and are transferred to consumers with high
92 efficiency (Gladyshev et al., 2011). There is some evidence that herbicides may affect the FA composition of
93 microalgae by interfering with vegetal lipid metabolism (Demailly et al., 2019; Gonçalves et al., 2021). Herbicides
94 may also induce changes in microorganism community structure of periphyton by selecting for more tolerant
95 species that differ in FA composition (Konschak et al., 2021). For example, diatoms are known to be rich in EPA,
96 while green algae are characterised by high content of ALA and bacteria by C18:1n9, C16:0 and C18:0. Thus,
97 there is considerable risk that herbicides reaching aquatic ecosystems may affect the structure of periphyton
98 assemblages and consequently alter the nutritional quality of this basal resource to higher consumers (Konschak
99 et al., 2021). Indeed, it has been shown that food quality affects growth and development of consumers (Da Costa
100 et al., 2023; Müller-Navarra et al., 2000; Rossoll et al., 2012).

101 This study investigated the effects of two herbicides frequently detected in aquatic ecosystems on complex
102 biofilm communities. Fatty acid composition was used as the main response variable due to the key role FAs play
103 in food webs. Most studies adopting a concentration-response exposure design have been carried out on mono-
104 specific cultures and over timescales of a few hours to a few days, with conventional descriptors such as cell
105 numbers, mortality, or photosynthetic capacity as response variables. To our knowledge, this study is one of the
106 first to adopt a concentration-response design with complex microorganism matrices (biofilms) in a chronic
107 context (7 and 14 days of exposure) and focusing on fatty acids as a response variable to pesticide contamination.
108 The primary aim was to provide information on the long-term effects of the tested herbicides. More specifically,
109 we conducted a laboratory experiment to (1) determine the effects of atrazine and S-metolachlor on periphyton FA
110 composition and to (2) relate possible modifications in FA profiles to changes in the community structure of
111 autotrophic organisms monitored by chlorophyll a fluorescence measurements. For this purpose, we exposed
112 cultured periphyton in microcosms to either atrazine or S-metolachlor along an environmentally relevant
113 concentration gradient.

114 2. Materials and methods

115 2.1. Experimental setup and periphyton sampling

116

117 Periphyton inoculum was collected in a stream (watershed= 82 km²) with low to moderate anthropogenic
118 activities (agricultural and urban) located a few kilometers west of Quebec City (Quebec, Canada; lat:
119 46°45'48.8"N, long: 71°21'24.0"W). The inoculum was acclimated in the laboratory in aquaria for two months
120 under experimental conditions (temperature = 20-22°C, natural photoperiod). Before the start of the experiment,
121 acclimated periphyton was evenly transferred in suspension into 23 microcosms (dimensions: 30 x 15 x 20 cm)
122 filled with 7.5 L of dechlorinated tap water enriched with nutrients (temperature = 20°C, photoperiod= 16h day/8h
123 night, average light flux= 54 $\mu\text{mol photons}\cdot\text{m}^{-2}\cdot\text{s}^{-1}$, nutrients summarised in Tab.S1) and equipped with an aeration
124 pump. Each microcosm contained six glass slides (double-sided for a total of 141 cm²) to increase the surface area
125 available for periphyton colonisation. After a one-month colonization period in the microcosms, periphyton were
126 exposed to a gradient of atrazine and S-metolachlor concentrations (PESTANAL, analytical standard, Sigma
127 Aldrich). The nominal concentrations of both herbicides tested were: 0, 5, 10, 50, 100, 500 and 1000 $\mu\text{g}\cdot\text{L}^{-1}$.
128 Treatments henceforth will be referred to by the first letter of the herbicide (A for Atrazine; S for S-metolachlor),
129 followed by the nominal concentration (e.g., A5, A10, S5, S10, etc.) and the treatment that did not receive herbicide
130 will be referred to as the control. We conducted a gradient study design where we chose to increase the number
131 of treatments to the detriment of replication (Larras et al., 2018). However, experimental replicates were
132 incorporated for the control (n = 4), A10 (n = 3), S10 (n = 3), and S100 (n = 3) treatments that we considered
133 environmentally relevant concentrations. Increasing the number of treatments tested instead of testing replication
134 is suggested to be an advantageous strategy (Green et al., 2018). A limited number of replications may result in
135 increased inter-treatment variability; however, this could be reduced by the number of measurements taken within
136 each replication. Within each microcosm, samples were collected on three occasions, before exposure (day 0), and
137 after 7 and 14 days of exposure. To ensure a homogeneous and representative sample, periphyton were scrapped
138 from randomly collected glass slides as well as from the walls of the microcosms to make one composite sample
139 per treatment which was preserved at -80°C for FA analyses.

140 Chlorophyll a fluorescence of green algae, diatoms and cyanobacteria composing the periphyton was measured
141 with the fluorometer probe Benthotorch (bbe Benthotorch, Moldaenke, Germany) that uses the excitation-
142 emission responses at several wavelengths (470 nm, 525 nm and 610 nm) to determine chlorophyll a concentrations
143 of attached autotrophic organisms. At each sampling time, six measurements were randomly taken per microcosm
144 by placing the instrument directly onto the glass slides that were delicately and temporarily removed from the
145 water. Biofilm samples were also collected and fixed with formaldehyde (3% from a stock formalin 37%) in order
146 to qualitatively compare the relative composition of the main algal groups with fluorescence data provided by the
147 Benthotorch. This comparison was conducted only for six samples as the objective was simply to verify the
148 fluorescence data. Despite the fact that green algae were observed under the microscope but were not very
149 abundant as measured with the probe (leading to an underestimation by the Benthotorch) their relative increase
150 in the biofilm during the course of the experiment was measured by the probe and qualitatively verified by
151 microscopy.

152 Throughout the experiment, pH=8.2 ± 0.1, conductivity=318.3 ± 25.6 μS.cm⁻¹ and water
 153 temperature=18.9 ± 0.3°C (n=66) were stable. Herbicide concentrations were determined by liquid
 154 chromatography (Finnigan Surveyor) with tandem mass spectrometry (TSQ Quantum Access; Thermo Scientific)
 155 (LC-MS/MS) (Limit of detection= 0.1 μg.L⁻¹, analytical standards: Atrazine-D5 and Metolachlor-D6). Herbicide
 156 concentrations were re-adjusted as needed over the course of the experiment. To determine any abiotic loss of
 157 atrazine and S-metolachlor in microcosms, three microcosms without periphyton were contaminated with atrazine
 158 at a nominal concentration of 500 μg.L⁻¹ and three additional microcosms were contaminated with 50 μg.L⁻¹ of S-
 159 metolachlor. Water was sampled after 7 days and analysed by LC-MS/MS following the same method as described
 160 above. Measured atrazine concentrations were close to nominal concentration in biotic microcosms, while S-
 161 metolachlor concentrations were below targeted values (S2 for details). Despite the fact that measured
 162 concentrations deviated from the targeted nominal concentrations, a concentration gradient was observed for both
 163 herbicides as seen in Tab.1.

164 Tab.1: Measured concentrations of atrazine and S-metolachlor (mean± standard deviation when treatment was replicated) at
 165 day 0, 7 and 14 before concentration adjustment. Italics: concentrations in abiotic microcosms

Nominal concentration (μg.L ⁻¹)	Measured concentration of atrazine (μg.L ⁻¹)			Measured concentration of S- metolachlor (μg.L ⁻¹)		
	0	7	14	0	7	14
Day						
Control (n=4)	<LOD	<LOD	<LOD	<LOD	<LOD	<LOD
5	1.0	4.9	5.3	3.4	1.9	2.2
10 (n=3)	8.7± 0.4	8.8± 1.1	8.9± 0.9	3.0± 0.8	3.7± 2.4	4.5± 1.3
50	47.5	46.4	37.9	26.5	28.1	29.4
50 abiotic condition (n=3)				38.6 ± 5.0	20.0 ± 1.8	
100 (n=3 for S-metolachlor)	19.1	80.6	73.8	109.5 ± 13.3	73.7 ± 15.8	74.6 ± 9.1
500	551.3	517.7	459.9	542.7	238.1	261.9
500 abiotic condition (n=3)	520.1 ± 5.0	428.9 ± 27.2				
1000	1259.3	1231.8	1266.1	998.2	272.5	567.8

166

167

2.2. Fatty acid analysis

Fatty acid extraction and analysis were performed according to Fadhlaoui et al., (2020), where a 40 mg subsample of periphyton was homogenized in 8.4 mL of chloroform/methanol (2v/1v) solution for 1 minute using a Homogenizer 850 (Fisherbrand™). A volume of 20 µL of trycosilic acid (C23:0) was added as an internal standard and the samples were then sonicated for 5 min using a Sonifier® (Branson). A 2 mL solution of NaCl (0.73%) was then added followed by centrifugation of the sample for 15 min at 3000 tr/min at 4 °C allowing for lipid separation in the lower phase. Lipids were recovered from this lower phase and evaporated using a TurboVap® (Caliper Life Sciences TurboVap II) for 15 min at 40 °C before being transferred to screw-capped tubes with 3 mL of BF₃ (boron trifluoride-methanol solution 14% in methanol). The BF₃ is used to esterified fatty acids and to facilitate analysis by gas chromatography. After one hour of incubation at 75 °C, fatty acid methyl esters (FAMES) were extracted by adding 3 mL of ultra-pure water and 3 mL of petroleum ether. This step was repeated two more times to improve FAMES recovery. The top fraction of petroleum ether was recovered and dried using the TurboVap® for 15 min at 40 °C. Finally, FAMES were dissolved in 240 µL of hexane and then transferred into screw-capped vials to be analyzed by gas chromatography with a flame ionization detector (Agilent Technologies; 7890D GC system) equipped with a fused silica capillary column (DB-FATWAX from Agilent Technologies: 30m [length], 0.250 mm [inner diameter], 0.25 µm [film thickness]). Injection was conducted at a constant pressure, and helium was used as the carrier gas. Temperature programming was as follows: initial temperature of 140 °C increased to 170 °C at a rate of 6.5 °C.min⁻¹, then to 200 °C at a rate of 2.75 °C.min⁻¹ for 14 min, and finally to 230 °C at a rate of 3 °C.min⁻¹ for 12 min. Because the periphyton is highly heterogeneous, five subsamples from the one composite sample collected in each microcosm were analysed (pseudo-replicates) to ensure a proper representation of fatty acid profiles within each microcosm.

2.3. Statistical analysis

Statistical analyses were performed in RStudio (R version 4.2.2). Water chemistry and chlorophyll a fluorescence data (µg of chlorophyll a.cm⁻²) were expressed as mean ± standard deviation. Due to inter-microcosm variability prior to exposure, photoautotroph fluorescence and FA composition changes (i.e., deltas Δ) between day 0 and the two sampling times (7 and 14) were used. Delta values were then used to perform linear regressions. For all statistical analyses, results were considered significant when the p-value was less than 0.05 and marginally significant where p-value was between 0.05 and 0.1. For photoautotroph chlorophyll a fluorescence, one-way ANOVAs were performed on raw data and only for replicated conditions. Pairwise t-tests with Bonferroni adjustment were used for post hoc comparisons.

Principal Component Analyses (PCA) were conducted on fatty acid data (including fatty acids with proportions >5% in at least one sample) from pseudo-replicates, allowing the representation of intra-condition variability. The “FactoMineR” and “factoextra” packages were used to explore patterns in FA profiles as a function of exposure concentrations. A PERMutational ANalysis Of VARIance (PERMANOVA) on dissimilarity matrix was performed on replicated conditions and was followed by a pairwise comparison to test for differences in FA profiles between conditions using the ‘adonis2’ (method=‘gower’) and ‘pairwise.adonis2’ functions from the ‘vegan’ package.

208 3. Results

209

210

211 3.1. Community structure of the autotrophic organisms

212

213 *Effect of atrazine on chlorophyll a fluorescence*


214

215 Diatoms and cyanobacteria were the two main groups of photoautotroph organisms in periphyton with green
216 algae having a lower relative chlorophyll a fluorescence (see Fig.S1 for raw data). Chlorophyll a fluorescence data
217 suggested inter-microcosms variability before contamination, in particular between controls and S100, A100,
218 A500 and A1000. The fluorescence probe detected green algae in certain microcosms and diatoms and
219 cyanobacteria exhibited lower levels under control condition. The data were standardized to reduce the effect of
220 pre-exposure variability where data for day 7 and day 14 were normalized using the data from day 0.

221 Under the control condition, the total chlorophyll a fluorescence significantly decreased between day 0 and
222 day 14 (Df=2, F=9.01, p-value=0.01). Especially, diatom specific fluorescence marginally decreased between 7
223 days and 14 days of exposure (Df=2, F=3.67, p-value=0.05). Linear regressions showed some effect of atrazine
224 on photoautotroph chlorophyll a fluorescence (Fig.1). Specifically, cyanobacteria and diatoms specific
225 fluorescence increased with atrazine concentration after 7 days (Df=10, F=26.44, R²=0.73, p-value<0.001 and
226 Df=10, F=28.98, R²=0.75, p-value<0.001, respectively) and 14 days of exposure (Df=10, F=20.36, R²=0.67, p-
227 value=0.001 and Df=10, F=25.83, R²=0.72, p-value<0.001, respectively). Green algae were a minor autotrophic
228 group based on chlorophyll a fluorescence, and atrazine did not appear to affect its chlorophyll a fluorescence as
229 it remained stable between exposure concentrations and over time.

230 For all photoautotrophic groups, no significant differences were observed in Δ Chla-fluorescence between
231 control and 10 $\mu\text{g.L}^{-1}$ conditions after 7 days (Df=5, with F=0.68, p-value=0.45 for cyanobacteria; F=3.25, p-
232 value=0.13 for diatoms and F=5.19, p-value=0.07 for green algae) and 14 days of exposure (Df=5, with F=0, p-
233 value=0.99 for cyanobacteria; F=0.15, p-value=0.72 for diatoms and F=0.98, p-value=0.37 for green algae).
234 Atrazine then appeared to have a significant effect on chlorophyll a fluorescence at concentrations higher than 10
235 $\mu\text{g.L}^{-1}$.

236



237

238 **Fig.1** Chlorophyll a fluorescence variation (Δ) at day 7 and day 14 compared to day 0 (initial time of the
 239 experiment), based on chlorophyll a fluorescence of photoautotrophic groups as a function of measured atrazine
 240 concentrations ($\mu\text{g.L}^{-1}$) (linear regression Df=10)

241


242 *Effect of S-metolachlor on chlorophyll a fluorescence*

243

244 As observed for atrazine, green algae remained the minor photosynthetic group. In contrast to atrazine,
245 S-metolachlor had no effect on photoautotroph chlorophyll a fluorescence (Fig.2).

246 The one-way ANOVA showed no effect of S-metolachlor at $10 \mu\text{g.L}^{-1}$ and $100 \mu\text{g.L}^{-1}$ on photoautotrophic
247 group fluorescence compared to the control condition after 7 days (Df=2, with $F=1.30$, p-value=0.33 for
248 cyanobacteria; $F=3.56$, p-value=0.09 for diatoms and $F=1.23$, p-value=0.35 for green algae) and 14 days of
249 exposure (Df=2, with $F=1.06$, p-value=0.40 for cyanobacteria; $F=2.02$, p-value=0.20 for diatoms and $F=0.69$, p-
250 value=0.54 for green algae).

251



252

253 **Fig.2** Chlorophyll a fluorescence variation (Δ) at day 7 and day 14 compared to day 0 (initial time of the
 254 experiment), based on chlorophyll a fluorescence of photoautotrophic groups as a function of measured S-
 255 metolachlor concentrations ($\mu\text{g.L}^{-1}$) (linear regression Df=12)

256
257


3.2. Effects of herbicides on periphyton fatty acid composition

258 *Effect of atrazine on fatty acids*

259 A total of 27 fatty acids were identified in the total lipid fraction of the periphyton. Average (+/- standard
260 deviation) proportions of each FA are presented as supplementary information (Tab.S3). Unsaturated fatty acids
261 (UFAs) were generally the predominant FA group in all treatments with a relative percentage of up to 62.9%
262 comprised mostly of mono-unsaturated fatty acids (MUFAs; 23.0% to 41.1%) followed by poly-unsaturated fatty
263 acids (PUFAs; 15.2% to 26.1%). Saturated fatty acids (SFAs) represented up to 54.3% of total lipid content in the
264 periphyton samples.

265 PCA of periphyton FA relative percentages 14 days after atrazine exposure explained 66.9% of the
266 variance in FA composition on two axes (dim1=42.9% and dim2=24%; Fig.3). PCAs of FA composition on day
267 0 and day 7 are presented in the supplementary information (Fig.S2). The A5 and A100 treatments clustered
268 together on the top left of the ordination and had higher proportions of MUFAs, in particular C16:1n7, compared
269 to the A50, A500 and A1000 treatments that clustered in the lower portion of the PCA and were more characterized
270 by SFAs and C18:0. A large dispersion of fatty acid data was observed, especially for the control and A10
271 conditions which overlapped all treatment groups.

272



273

274 **Fig.3** Principal component analysis (PCA) of fatty acid profiles at day 14 for the different atrazine conditions. The
275 left panel is the graph of individual FA and on the right panel corresponds to the circle of correlations. Ellipses
276 have been plotted with a confidence level of 80%. All pseudo-replicates were considered to better represent the
277 intra-condition variability


278

279

280 The PERMANOVA and pairwise comparisons performed on replicated conditions (control and A10)
281 revealed a significant difference in FA composition (Df=1, F=3.92; p-value=0.02) after 14 days under atrazine
282 exposure. PERMANOVA were also conducted at day 0 and day 7 and showed that differences were already present
283 at day 7 (Df=1, F=4.88; p-value=0.008) but not prior to exposure (Df=1, F=2.33, p-value=0.08).

284 Linear regressions showed that atrazine concentration did not markedly affect the main FA groups (Fig.4;
285 linear regressions for individual FA are shown in Fig.S3). Only a regression marginally significant was observed
286 for SFAs at 14 days (Df=10, R²=0.27; p-value=0.08).

287



289

290 **Fig.4** Linear regressions based on differences in proportions ($\Delta\%$) for the main fatty acid groups as a function of
 291 measured concentrations of atrazine ($\mu\text{g.L}^{-1}$) after 7 days and 14 days of exposure

292 *Effect of S-metolachlor on fatty acids*

293


294 Mean (+/- standard deviation) proportions of each FA are presented in the supplementary information
295 (Tab.S4). Unsaturated fatty acids (UFAs) comprised up to 62.5% of the total FA content of periphyton among
296 treatments, while mono-unsaturated fatty acids (MUFAs) varied from 27.3% to 38.8% and poly-unsaturated fatty
297 acids (PUFAs) varied from 16.9% to 33.4%. Saturated fatty acids (SFAs) represented up to 54.3% of total FA
298 content in the periphyton samples.

299 A PCA was performed to assess the effect of S-metolachlor after 14 days of exposure (Fig.3) (See Fig.S4
300 for 0 and 7 days). The first two dimensions explained 62% of the variance (dim1=42.1% and dim2=19.9%). The
301 two highest concentrations clustered on the left side of the ordination, while S10, S100 clustered on the right side.
302 The S5 and S50 conditions clustered on the top portion of the ordination (dimension 2) and S100 clustered on the
303 lower half of the ordination. The control condition clustered in the middle and showed high dispersion. High S-
304 metolachlor concentrations (S500 and S1000) were more associated with SFAs such as C18:1n9 and C18:0, while
305 lower S-metolachlor concentrations were rather characterized by PUFAs such as ALA, EPA and C20:4n6.

306

307

308




309 **Fig.5** Principal component analysis (PCA) of fatty acid profiles at day 14 for the different S-metolachlor
310 conditions. The left panel is the graph of individual FA and on the right panel corresponds to the circle of
311 correlations. Ellipses have been plotted with a confidence level of 80%. All pseudo-replicates were considered to
312 better represent the intra-condition variability
313

315 At day 14, the PERMANOVA (performed only on replicated conditions; control, S10 and S100) showed
316 a significant effect of S-metolachlor concentrations on the fatty acid composition of the periphyton. Indeed, there
317 was a strong dissimilarity between the control and S100 (Df=2, F=3.79, p-value=0.001). PERMANOVA
318 conducted at day 0 and day 7 also revealed differences in FA profiles. Especially, the S10 condition (Df=1, F=4.70,
319 p-value=0.008) was already different from the control at day 0, while S100 had different FA composition from the
320 control (Df=1, F=7.35, p-value=0.003) after 7 days of S-metolachlor exposure. These results suggest that S-
321 metolachlor affected the fatty acid profile of periphyton after only 7 days of exposure.

322 Linear regressions showed an effect of S-metolachlor contamination on the FA composition of the
323 periphyton (Fig.6; linear regressions for some specific FAs are shown in Fig.S5). Specifically, SFAs increased
324 along the S-metolachlor gradient (Df=12, F=18.70, R²=0.60; p-value<0.001) after 14 days of exposure. MUFAs
325 did not vary with exposure concentrations, while PUFAs marginally decreased with increasing S-metolachlor
326 concentrations after 7 days of exposure (Df=12, F=3.23, R²=0.24; p-value=0.08) and then significantly decreased
327 after 14 days (Df=12, F=5.05, R²=0.30; p-value=0.04). Finally, VLCFAs decreased with increasing herbicide
328 concentration after 7 days (Df=12, F=5.85, R²=0.33; p-value=0.03). This relationship was stronger after 14 days
329 (Df=12, F=19.61, R²=0.62; p-value<0.001), where a delta of 10% between the highest concentration and the
330 control was observed.

331



333

334 **Fig.6** Linear regressions based on differences in proportions ($\Delta\%$) for the main fatty acid groups as a function of
 335 measured concentrations of S-metolachlor ($\mu\text{g.L}^{-1}$) after 7 days and 14 days of exposure

336 4. Discussion

337

338 Despite high variability observed in photoautotroph community structure and FA composition, results
339 showed some effects of herbicide exposure. Photoautotroph related chlorophyll a fluorescence tended to increase
340 with the increase of atrazine concentration after 7 days of exposure. In contrast, S-metolachlor did not clearly
341 affect periphyton fluorescence. As periphytic biofilms are very heterogeneous, fluorescence and fatty acid data
342 showed large intra-condition variability. Despite marked variability, results showed that the two herbicides, in
343 particular S-metolachlor, affected fatty acid profiles. S-metolachlor had a stronger effect than atrazine, with a
344 greater effect after 14 days of exposure compared to 7 days. In particular, the S500 condition showed a 30%
345 increase in SFAs and a 34% decrease in VLCFAs proportions compared to the control condition.

346 4.1. Effects of herbicides on photoautotrophs chlorophyll a fluorescence

347 S-metolachlor exposure did not clearly affect periphyton biomass as measured by chlorophyll a
348 fluorescence. Indeed, total chlorophyll a fluorescence and specific diatom fluorescence decreased over time under
349 all concentrations including the control condition. Our finding of no significant herbicide effect is in contrast with
350 several studies that showed chloroacetamide herbicides to decrease photoautotroph growth and chlorophyll a
351 fluorescence. For example, Thakkar et al. (2013) showed that the exposure of the marine chlorophyte *Dunaliella*
352 *tertiolecta* to a high metolachlor concentration (1 mg.L⁻¹) led to a decrease in chlorophyll a and b fluorescence and
353 inhibited cell growth. Likewise, Coquillé et al. (2015) showed a decrease in chlorophyll a content in a freshwater
354 diatom culture (*Gomphonema gracile*) after 7 days of exposure to 100 µg.L⁻¹ of S-metolachlor. The limited effect
355 of S-metolachlor on photoautotrophic groups could be linked to the periphyton matrix that is composed by
356 extracellular polymeric substances (EPS) which may represent up to 90% of the dry mass (Flemming and
357 Wingender, 2010). The EPS matrix has several functional groups allowing for the sorption of nutrients and
358 xenobiotics, but can also form a protective layer for the biofilm cells against substances such as pesticides (Melo
359 et al., 2022). The overall decrease in periphyton chlorophyll a fluorescence (i.e biomass) that we observed over
360 time is likely due to the age of the periphyton in our experiment. The colonization time of periphyton varies
361 between 2 and 4 weeks (Cattaneo and Amireault, 1992), and is followed by a biomass loss phase after 4 to 5 weeks
362 (Trbojević et al., 2017). In order to have sufficient biomass for fatty acid analyses, periphyton was contaminated
363 after 4 weeks of colonization and growth. As the experiment lasted an additional 14 days, the periphyton may have
364 started a senescence phase, with potential detachment of biomass under all treatment conditions (Boulêtreau et al.,
365 2006).

366 In contrast, atrazine increased diatoms and cyanobacteria biomass measured by chlorophyll a
367 fluorescence. The increase in chlorophyll fluorescence can be linked to the mode of action of atrazine. When
368 photosynthesis is proceeding normally, several steps contribute to the creation of an electron flow between
369 different elements of the thylakoid membranes where the inhibition of photosystem II (PSII) by atrazine takes
370 place. Atrazine competes with plastoquinone for the quinone binding site on the D1 protein (QB site) in PSII,
371 interrupting the electron flow from plastoquinone QA to QB (Rea et al., 2009) leading to the re-emission of
372 excitation energy as fluorescence (Muller et al., 2008) which is then captured by our measuring device. The
373 increase in chlorophyll a fluorescence could also be linked to an increase of chlorophyll cell content. When

374 exposed to atrazine, autotrophs within the periphyton may physiologically adapt to stress by increasing chlorophyll
375 a content per cell (Pannard et al., 2009) to increase the number of photosystems. This “shade-adaptation” response
376 may be a strategy to compensate for the inhibition of photosynthesis and has previously been documented to occur
377 in response to other PSII inhibitor herbicides (e.g., diuron; Chesworth et al., 2004; Proia et al., 2011; Ricart et al.,
378 2009). Given the mode of action of atrazine, the increase in chlorophyll a fluorescence could be taken as evidence
379 of an atrazine effect on the periphyton.

380 In addition to affecting the fluorescence of photosynthetic organisms, the presence of herbicides may
381 select for more resistant/tolerant taxa (Murdock et al., 2013), thus modifying the community structure of the
382 periphyton (Schmitt-Jansen and Altenburger, 2005). We found that atrazine exposure increased the chlorophyll a
383 fluorescence of cyanobacteria in periphyton. Cyanobacteria may be more tolerant to atrazine and more competitive
384 than diatoms and green algae as they have the potential to adapt to photosynthesis inhibition by the use of
385 alternative carbon fixation pathways (Egorova and Bukhov, 2006). This is consistent with Pannard et al. (2009)
386 who showed that chronic exposure to atrazine (0.1, 1 and 10 $\mu\text{g}\cdot\text{L}^{-1}$ for 7 weeks) led to a change in microalgal
387 populations with the selection of opportunistic resistant species, some of which were cyanobacteria. Herbicides
388 could also decrease competition for nutrients or increase labile carbon released after cell death further stimulating
389 bacterial production (Downing et al., 2004).

390 The response of periphyton to herbicide contamination may be reversible as recovery from herbicide
391 exposure as been observed (King et al., 2016; Morin et al., 2010; Prosser et al., 2013). For example, Laviale et al.,
392 (2011) showed that a 7h exposure to atrazine affects the effective and optimal quantum yields of PSII
393 photochemistry of periphyton, but that these effects were reversible within 12 hours. However, Proia et al., (2011)
394 mentioned that although the biofilm has a high recovery potential, pulses of contamination longer than a few hours
395 could result in more persistent effects on the biofilm.

396 **4.2. Effects of herbicides on periphyton fatty acid composition**

397 Herbicide exposure caused different changes in the FA composition of periphyton with atrazine having
398 little effect on FA composition compared to S-metolachlor. In particular, periphyton SFAs increased with S-
399 metolachlor concentrations, while PUFAs and VLCFAs decreased. The decrease in VLCFAs proportion with the
400 increase in S-metolachlor concentration exposure is consistent with previous results from Böger (2003), who
401 showed a 68% inhibition in VLCFAs of *Scenedesmus acutus* (green algae) after exposure to 283 $\mu\text{g}\cdot\text{L}^{-1}$ of S-
402 metolachlor. VLCFAs ($C\geq 20$) have a structural role in membranes (Bach et al., 2011; Vallotton et al., 2008).
403 S-metolachlor binds to the fatty acid elongation synthase (FAE1-synthase) and inhibits the formation of VLCFAs,
404 which can then affect the rigidity and permeability of cell membranes, resulting in increased cell size and impaired
405 cell division (Matthes and Böger, 2002; Thakkar et al., 2013). Even at lower concentrations of exposure (10 $\mu\text{g}\cdot\text{L}^{-1}$),
406 Demailly et al. (2019) experimentally showed that S-metolachlor significantly increased the saturated fatty acid
407 C16:0 and decreased PUFAs including C18:4n3 and C20:4n6 of the diatom *Gomphonema gracile* after one week
408 of exposure. The loss of PUFAs observed here and in past studies may be due to the ability of S-metolachlor to
409 increase ROS production (e.g., singlet oxygen $^1\text{O}_2$) resulting in the peroxidation of unsaturated fatty acids in lipid
410 membranes. More specifically, these ROS remove hydrogen from the unsaturated chain of PUFAs constituting the
411 lipids, leading to the loss of membrane integrity (Maronić et al., 2018), in turn jeopardizing the functioning of the

412 cell (Garg and Manchanda, 2009). In response to stress, algae often produce triacylglycerols (TAGs) (Nakamura
413 and Li-Beisson, 2016; Shanta et al., 2021). TAGs are considered as carbon and energy storage products (Morales
414 et al., 2021) and are used to maintain bioenergetic stability in the cell. SFAs and MUFAs such as C16:0, C18:0
415 and C18:1 are among the main components of triacylglycerols (TAGs). The increase in SFAs (e.g., C18:0) and the
416 decrease in some long-chain UFAs that we observed in our experiment could therefore suggest a protective
417 response of the cells against membrane S-metolachlor damages (Kabra et al., 2014).

418 Herbicides can also have an indirect effect on the FA composition of periphyton by altering the taxonomic
419 composition of periphyton communities. Indeed, different taxonomic groups in the periphyton complex have
420 different fatty acid profiles. For example, diatoms are particularly rich in EPA (C20:5n3) (Drerup and Vis, 2016)
421 and green algae are rich in ALA (C18:3n3) (Genter and Lehman, 2000), while the SFA C16:0 (palmitic acid) is
422 important for the structure of phospholipid membranes in prokaryotes (Rock, 2008). Changes in the proportion of
423 fatty acids may thus reflect herbicide-induced changes in the composition of the periphyton communities. More
424 specifically, it is possible that the increase in SFAs with atrazine exposure may be due to an increase in bacteria
425 resulting from reduced competition with photosynthetic organisms impacted by the contaminant as we observed
426 increased cyanobacteria biomass (measured by chlorophyll a fluorescence and expressed in $\mu\text{gchl}a.\text{cm}^{-2}$) by this
427 contaminant (Figure S3). Nevertheless, this increase in cyanobacteria was hardly detectable in the FA profiles,
428 where no significant changes in C18:2n6 and C18:3n3 were observed despite the fact that cyanobacteria are
429 generally rich in these C18 PUFAs (Desvillettes et al., 1997). At present, it is still unclear what level of organization
430 (i.e., from the cellular level to subtle changes at the community level) is responsible for the changes in the FA
431 composition of periphyton highlighted by our experiment. It would then be useful to carry out further studies to
432 use more endpoints such as specific composition and the number of cells per autotrophic group.

433 As previously mentioned, some studies suggest a recovery of periphyton structure and function after a
434 post-exposure recovery phase to herbicides. Most studies used endpoints such as photosynthetic parameters,
435 biomass, or taxonomic composition. However, to our knowledge, there are no studies using biofilm fatty acid
436 composition to monitor recovery nor focusing on how recovery time may affect consumer organisms. In
437 agricultural streams, pesticides contamination generally occurs by pulse, via surface runoff processes. Pulse
438 exposure can occur in various scenarios, depending on its intensity (pulse height), duration (pulse width) and
439 frequency/recovery time (Chèvre and Vallotton, 2013). As a result, reproducing realistic pulse exposure scenarios
440 in a laboratory context may be complicated by logistical constraints. As only few studies if any have examined the
441 influence of atrazine and S-metolachlor on fatty acid profiles in biofilms, using a continuous exposure approach
442 provides a simpler way of controlling experimental conditions. However, to reproduce environmentally realistic
443 conditions, it would be interesting to consider chronic multi-pulse exposure experiments including biofilm
444 recovery phases (Giddings et al., 2018; King et al., 2016).

445 **5. Conclusion**

446

447 Periphyton plays a key role in the structure and function of aquatic ecosystems. The nutritional quality of
448 periphyton is essential for the development of primary consumers and can be used as indicator of ecosystem health
449 (Desvillettes et al., 1997). Fatty acids are key nutritional compounds transferred through trophic interactions that
450 are sensitive to various environmental contaminants. We investigated the effects of two commonly used
451 agricultural herbicides, atrazine and S-metolachlor on periphyton and found that the two herbicides acted
452 differently on the periphyton photoautotroph chlorophyll a fluorescence and fatty acid composition suggesting that
453 there is no standard pattern of herbicide effects on stream periphytic communities. Fluorescence measurements
454 provided information on changes in the relative proportion of the photoautotrophic groups (i.e., green algae,
455 diatoms and cyanobacteria) within periphyton, however, we were limited in our quantification of heterotrophs.
456 Considering that bacteria account for a large amount of biofilm mass (Ricart et al., 2009), are involved in nutrient
457 cycles and can affect the fate of herbicides in water and within the biofilm, future studies should investigate the
458 heterotrophic compartment of the biofilm, especially by DNA sequencing or the study of the metabolism of
459 bacteria. The widespread presence of these two herbicides in rivers raises the question of their toxicity to non-
460 target aquatic organisms and their interaction with the many other molecules present in water (i.e., antagonist,
461 additive or synergistic effects) (Glinski et al., 2018). This study supports the interest to use fatty acids as
462 biomarkers (Gugger, 2002; Lang et al., 2011; Maltsev and Maltseva, 2021; Shen et al., 2016) in the context of
463 pesticide effect assessment (Filimonova et al., 2016; Gonçalves et al., 2021) but also as a tool for water quality
464 biomonitoring (George et al., 2016).

465 References

- 466
- 467 Bach, L., Gissot, L., Marion, J., Tellier, F., Moreau, P., Satiat-Jeunemaître, B., Palauqui, J.-C., Napier,
468 J.A., Faure, J.-D., 2011. Very-long-chain fatty acids are required for cell plate formation during
469 cytokinesis in *Arabidopsis thaliana*. *Journal of Cell Science* 124, 3223–3234.
470 <https://doi.org/10.1242/jcs.074575>
- 471 Bachetti, R.A., Urseler, N., Morgante, V., Damilano, G., Porporatto, C., Agostini, E., Morgante, C.,
472 2021. Monitoring of Atrazine Pollution and its Spatial-Seasonal Variation on Surface Water
473 Sources of an Agricultural River Basin. *Bull Environ Contam Toxicol* 106, 929–935.
474 <https://doi.org/10.1007/s00128-021-03264-x>
- 475 Battaglin, W.A., Furlong, E.T., Burkhardt, M.R., Peter, C.J., 2000. Occurrence of sulfonylurea,
476 sulfonamide, imidazolinone, and other herbicides in rivers, reservoirs and ground water in
477 the Midwestern United States, 1998. *Science of The Total Environment* 248, 123–133.
478 [https://doi.org/10.1016/S0048-9697\(99\)00536-7](https://doi.org/10.1016/S0048-9697(99)00536-7)
- 479 Battaglin, W.A., Thurman, E.M., Kalkhoff, S.J., Porter, S.D., 2003. Herbicides and transformation
480 products in surface waters of the midwestern united states. *J Am Water Resources Assoc* 39,
481 743–756. <https://doi.org/10.1111/j.1752-1688.2003.tb04402.x>
- 482 Baxter, L., Brain, R.A., Lissemore, L., Solomon, K.R., Hanson, M.L., Prosser, R.S., 2016. Influence of
483 light, nutrients, and temperature on the toxicity of atrazine to the algal species *Raphidocelis*
484 *subcapitata*: Implications for the risk assessment of herbicides. *Ecotoxicology and*
485 *Environmental Safety* 132, 250–259. <https://doi.org/10.1016/j.ecoenv.2016.06.022>
- 486 Böger, P., 2003. Mode of Action for Chloroacetamides and Functionally Related Compounds. *J. Pestic.*
487 *Sci.* 28, 324–329. <https://doi.org/10.1584/jpestics.28.324>
- 488 Boulêtreau, S., Garabetian, F., Sauvage, S., Sanchez-Perez, J.-M., 2006. Assessing the importance of a
489 self-generated detachment process in river biofilm models. *Freshwater Biol* 51, 901–912.
490 <https://doi.org/10.1111/j.1365-2427.2006.01541.x>
- 491 Brain, R.A., Anderson, J.C., 2019. The agro-enabled urban revolution, pesticides, politics, and popular
492 culture: a case study of land use, birds, and insecticides in the USA. *Environ Sci Pollut Res* 26,
493 21717–21735. <https://doi.org/10.1007/s11356-019-05305-9>
- 494 Brett, M., Müller-Navarra, D., 1997. The role of highly unsaturated fatty acids in aquatic foodweb
495 processes. *Freshwater Biology* 38, 483–499. <https://doi.org/10.1046/j.1365-2427.1997.00220.x>
- 496
- 497 Cattaneo, A., Amireault, M.C., 1992. How Artificial Are Artificial Substrata for Periphyton? *Journal of*
498 *the North American Benthological Society* 11, 244–256. <https://doi.org/10.2307/1467389>
- 499 Chesworth, J.C., Donkin, M.E., Brown, M.T., 2004. The interactive effects of the antifouling herbicides
500 Irgarol 1051 and Diuron on the seagrass *Zostera marina* (L.). *Aquatic Toxicology* 66, 293–305.
501 <https://doi.org/10.1016/j.aquatox.2003.10.002>
- 502 Chèvre, N., Vallotton, N., 2013. Pulse Exposure in Ecotoxicology, in: Féraud, J.-F., Blaise, C. (Eds.),
503 *Encyclopedia of Aquatic Ecotoxicology*. Springer Netherlands, Dordrecht, pp. 917–926.
504 https://doi.org/10.1007/978-94-007-5704-2_84
- 505 Coquillé, N., Jan, G., Moreira, A., Morin, S., 2015. Use of diatom motility features as endpoints of
506 metolachlor toxicity. *Aquatic Toxicology* 158, 202–210.
507 <https://doi.org/10.1016/j.aquatox.2014.11.021>
- 508 Da Costa, F., González-Araya, R., Robert, R., 2023. Using combinations of microalgae to condition
509 European flat oyster (*Ostrea edulis*) broodstock and feed the larvae: Effects on reproduction,
510 larval production and development. *Aquaculture* 568, 739302.
511 <https://doi.org/10.1016/j.aquaculture.2023.739302>
- 512 de Albuquerque, F.P., de Oliveira, J.L., Moschini-Carlos, V., Fraceto, L.F., 2020. An overview of the
513 potential impacts of atrazine in aquatic environments: Perspectives for tailored solutions
514 based on nanotechnology. *Science of The Total Environment* 700, 134868.
515 <https://doi.org/10.1016/j.scitotenv.2019.134868>

516 Debenest, T., Pinelli, E., Coste, M., Silvestre, J., Mazzella, N., Madigou, C., Delmas, F., 2009. Sensitivity
517 of freshwater periphytic diatoms to agricultural herbicides. *Aquatic Toxicology* 93, 11–17.
518 <https://doi.org/10.1016/j.aquatox.2009.02.014>

519 DeLorenzo, M.E., Lauth, J., Pennington, P.L., Scott, G.I., Ross, P.E., 1999. Atrazine effects on the
520 microbial food web in tidal creek mesocosms. *Aquatic Toxicology* 46, 241–251.
521 [https://doi.org/10.1016/S0166-445X\(98\)00132-5](https://doi.org/10.1016/S0166-445X(98)00132-5)

522 Demailly, F., Elfeky, I., Malbezin, L., Le Guédard, M., Eon, M., Bessoule, J.-J., Feurtet-Mazel, A.,
523 Delmas, F., Mazzella, N., Gonzalez, P., Morin, S., 2019. Impact of diuron and S-metolachlor on
524 the freshwater diatom *Gomphonema gracile*: Complementarity between fatty acid profiles
525 and different kinds of ecotoxicological impact-endpoints. *Science of The Total Environment*
526 688, 960–969. <https://doi.org/10.1016/j.scitotenv.2019.06.347>

527 Desvillettes, Ch., Bourdier, G., Amblard, Ch., Barth, B., 1997. Use of fatty acids for the assessment of
528 zooplankton grazing on bacteria, protozoans and microalgae. *Freshwater Biology* 38, 629–
529 637. <https://doi.org/10.1046/j.1365-2427.1997.00241.x>

530 Downing, H.F., Delorenzo, M.E., Fulton, M.H., Scott, G.I., Madden, C.J., Kucklick, J.R., 2004. Effects of
531 the Agricultural Pesticides Atrazine, Chlorothalonil, and Endosulfan on South Florida
532 Microbial Assemblages. *Ecotoxicology* 13, 245–260.
533 <https://doi.org/10.1023/B:ECTX.0000023569.46544.9f>

534 Drerup, S.A., Vis, M.L., 2016. Responses of Stream Biofilm Phospholipid Fatty Acid Profiles to Acid
535 Mine Drainage Impairment and Remediation. *Water Air Soil Pollut* 227, 159.
536 <https://doi.org/10.1007/s11270-016-2856-5>

537 Egorova, E.A., Bukhov, N.G., 2006. Mechanisms and functions of photosystem I-related alternative
538 electron transport pathways in chloroplasts. *Russ J Plant Physiol* 53, 571–582.
539 <https://doi.org/10.1134/S1021443706050013>

540 European Commission, 2004. Commission Regulation (EC) No 775/2004 of 26 April 2004 amending
541 Annex I to Regulation (EC) No 304/2003 of the European Parliament and of the Council
542 concerning the export and import of dangerous chemicals (Text with EEA relevance), CE.

543 Fadhlouli, M., Laderriere, V., Lavoie, I., Fortin, C., 2020. Influence of Temperature and Nickel on Algal
544 Biofilm Fatty Acid Composition. *Environ Toxicol Chem* 39, 1566–1577.
545 <https://doi.org/10.1002/etc.4741>

546 FAO, 2022. Pesticides use, pesticides trade and pesticides indicators. FAO.
547 <https://doi.org/10.4060/cc0918en>

548 Filimonova, V., Gonçalves, F., Marques, J.C., De Troch, M., Gonçalves, A.M.M., 2016. Fatty acid
549 profiling as bioindicator of chemical stress in marine organisms: A review. *Ecological*
550 *Indicators* 67, 657–672. <https://doi.org/10.1016/j.ecolind.2016.03.044>

551 Flemming, H.-C., Wingender, J., 2010. The biofilm matrix. *Nat Rev Microbiol* 8, 623–633.
552 <https://doi.org/10.1038/nrmicro2415>

553 Fortier, A., 2018. Règlement modifiant le Code de gestion des pesticides 5.

554 Gao, Y., Fang, Jianguang, Li, W., Wang, X., Li, F., Du, M., Fang, Jinghui, Lin, F., Jiang, W., Jiang, Z., 2019.
555 Effects of atrazine on the physiology, sexual reproduction, and metabolism of eelgrass
556 (*Zostera marina* L.). *Aquatic Botany* 153, 8–14.
557 <https://doi.org/10.1016/j.aquabot.2018.10.002>

558 Garg, N., Manchanda, G., 2009. ROS generation in plants: Boon or bane? *Plant Biosystems - An*
559 *International Journal Dealing with all Aspects of Plant Biology* 143, 81–96.
560 <https://doi.org/10.1080/11263500802633626>

561 Genter, R.B., Lehman, R.M., 2000. Metal toxicity inferred from algal population density,
562 heterotrophic substrate use, and fatty acid profile in a small stream. *Environ Toxicol Chem*
563 19, 869–878. <https://doi.org/10.1002/etc.5620190413>

564 George, S.D., Ernst, A.G., Baldigo, B.P., Honeyfield, D.C., 2016. Response of periphyton fatty acid
565 composition to supplemental flows in the upper Esopus Creek, Catskill Mountains, New York:
566 U.S. Geological Survey Scientific Investigations Report (Scientific Investigations Report). U.S.
567 Geological Survey, Reston, Virginia.

568 Giddings, J.M., Campana, D., Nair, S., Brain, R., 2018. Data quality scoring system for microcosm and
569 mesocosm studies used to derive a level of concern for atrazine: Atrazine Microcosm and
570 Mesocosm Data Quality Scoring. *Integr Environ Assess Manag* 14, 489–497.
571 <https://doi.org/10.1002/ieam.4050>

572 Gladyshev, M.I., Sushchik, N.N., Anishchenko, O.V., Makhutova, O.N., Kolmakov, V.I., Kalachova, G.S.,
573 Kolmakova, A.A., Dubovskaya, O.P., 2011. Efficiency of transfer of essential polyunsaturated
574 fatty acids versus organic carbon from producers to consumers in a eutrophic reservoir.
575 *Oecologia* 165, 521–531. <https://doi.org/10.1007/s00442-010-1843-6>

576 Glinski, D.A., Purucker, S.T., Van Meter, R.J., Black, M.C., Henderson, W.M., 2018. Analysis of
577 pesticides in surface water, stemflow, and throughfall in an agricultural area in South
578 Georgia, USA. *Chemosphere* 209, 496–507.
579 <https://doi.org/10.1016/j.chemosphere.2018.06.116>

580 Gonçalves, A.M.M., Rocha, C.P., Marques, J.C., Gonçalves, F.J.M., 2021. Fatty acids as suitable
581 biomarkers to assess pesticide impacts in freshwater biological scales – A review. *Ecological*
582 *Indicators* 122, 107299. <https://doi.org/10.1016/j.ecolind.2020.107299>

583 Green, J.W., Springer, T.A., Holbech, H., 2018. *Statistical analysis of ecotoxicity studies*, First edition.
584 ed. John Wiley & Sons, Hoboken, NJ

585 Griffini, O., Bao, M.L., Barbieri, D., Pantani, F., 1997. Occurrence of Pesticides in the Arno River and in
586 Potable Water - A Survey of the Period 1992-1995 8.

587 Groner, M.L., Relyea, R.A., 2011. A tale of two pesticides: how common insecticides affect aquatic
588 communities: A tale of two pesticides. *Freshwater Biology* 56, 2391–2404.
589 <https://doi.org/10.1111/j.1365-2427.2011.02667.x>

590 Gugger, M., 2002. Cellular fatty acids as chemotaxonomic markers of the genera *Anabaena*,
591 *Aphanizomenon*, *Microcystis*, *Nostoc* and *Planktothrix* (cyanobacteria). *International journal*
592 *of systematic and evolutionary microbiology* 52, 1007–1015.
593 <https://doi.org/10.1099/ijs.0.01917-0>

594 Guo, F., Kainz, M.J., Sheldon, F., Bunn, S.E., 2016a. Effects of light and nutrients on periphyton and
595 the fatty acid composition and somatic growth of invertebrate grazers in subtropical streams.
596 *Oecologia* 181, 449–462. <https://doi.org/10.1007/s00442-016-3573-x>

597 Guo, F., Kainz, M.J., Sheldon, F., Bunn, S.E., 2016b. The importance of high-quality algal food sources
598 in stream food webs - current status and future perspectives. *Freshw Biol* 61, 815–831.
599 <https://doi.org/10.1111/fwb.12755>

600 Hansen, S.P., Messer, T.L., Mittelstet, A.R., 2019. Mitigating the risk of atrazine exposure: Identifying
601 hot spots and hot times in surface waters across Nebraska, USA. *Journal of Environmental*
602 *Management* 250, 109424. <https://doi.org/10.1016/j.jenvman.2019.109424>

603 HRAC, 2020. Global herbicide classification lookup. Herbicide resistance action committee.

604 Huggins, K., Frenette, J.-J., Arts, M.T., 2004. Nutritional quality of biofilms with respect to light regime
605 in Lake Saint-Pierre (Quebec, Canada). *Freshwater Biol* 49, 945–959.
606 <https://doi.org/10.1111/j.1365-2427.2004.01236.x>

607 Kabra, A.N., Ji, M.-K., Choi, J., Kim, J.R., Govindwar, S.P., Jeon, B.-H., 2014. Toxicity of atrazine and its
608 bioaccumulation and biodegradation in a green microalga, *Chlamydomonas mexicana*.
609 *Environ Sci Pollut Res* 21, 12270–12278. <https://doi.org/10.1007/s11356-014-3157-4>

610 Kapsi, M., Tsoutsis, C., Paschalidou, A., Albanis, T., 2019. Environmental monitoring and risk
611 assessment of pesticide residues in surface waters of the Louros River (N.W. Greece). *Science*
612 *of The Total Environment* 650, 2188–2198. <https://doi.org/10.1016/j.scitotenv.2018.09.185>

613 King, R.S., Brain, R.A., Back, J.A., Becker, C., Wright, M.V., Toteu Djomte, V., Scott, W.C., Virgil, S.R.,
614 Brooks, B.W., Hosmer, A.J., Chambliss, C.K., 2016. Effects of pulsed atrazine exposures on
615 autotrophic community structure, biomass, and production in field-based stream
616 mesocosms: Pulsed atrazine exposures. *Environ Toxicol Chem* 35, 660–675.
617 <https://doi.org/10.1002/etc.3213>

618 Korschak, M., Zubrod, J.P., Duque Acosta, T.S., Bouchez, A., Kroll, A., Feckler, A., Röder, N., Baudy, P.,
619 Schulz, R., Bundschuh, M., 2021. Herbicide-Induced Shifts in the Periphyton Community

620 Composition Indirectly Affect Feeding Activity and Physiology of the Gastropod Grazer
621 *Physella acuta*. Environ. Sci. Technol. 55, 14699–14709.
622 <https://doi.org/10.1021/acs.est.1c01819>

623 Lang, I., Hodac, L., Friedl, T., Feussner, I., 2011. Fatty acid profiles and their distribution patterns in
624 microalgae: a comprehensive analysis of more than 2000 strains from the SAG culture
625 collection. BMC Plant Biol 11, 124. <https://doi.org/10.1186/1471-2229-11-124>

626 Larras, F., Billoir, E., Baillard, V., Siberchicot, A., Scholz, S., Wubet, T., Tarkka, M., Schmitt-Jansen, M.,
627 Delignette-Muller, M.-L., 2018. DRomics: A Turnkey Tool to Support the Use of the Dose–
628 Response Framework for Omics Data in Ecological Risk Assessment. Environ. Sci. Technol. 52,
629 14461–14468. <https://doi.org/10.1021/acs.est.8b04752>

630 Laviale, M., Morin, S., Créach, A., 2011. Short term recovery of periphyton photosynthesis after pulse
631 exposition to the photosystem II inhibitors atrazine and isoproturon. Chemosphere 84, 731–
632 734. <https://doi.org/10.1016/j.chemosphere.2011.03.035>

633 LAWA, 2019. Rapport sur la qualité des eaux souterraines - produits phytosanitaires.
634 Länderarbeitsgemeinschaft Wasser (LAWA) - Sous-comité Produits phytosanitaires dans les
635 eaux souterraines.

636 Li, H.-Y., Lu, Y., Zheng, J.-W., Yang, W.-D., Liu, J.-S., 2014. Biochemical and Genetic Engineering of
637 Diatoms for Polyunsaturated Fatty Acid Biosynthesis. Marine Drugs 12, 153–166.
638 <https://doi.org/10.3390/md12010153>

639 Maltsev, Y., Maltseva, K., 2021. Fatty acids of microalgae: diversity and applications. Rev Environ Sci
640 Biotechnol 20, 515–547. <https://doi.org/10.1007/s11157-021-09571-3>

641 Matthes, B., Böger, P., 2002. Chloroacetamides Affect the Plasma Membrane. Zeitschrift für
642 Naturforschung C 57, 843–852. <https://doi.org/10.1515/znc-2002-9-1015>

643 Melo, A., Quintelas, C., Ferreira, E.C., Mesquita, D.P., 2022. The Role of Extracellular Polymeric
644 Substances in Micropollutant Removal. Front. Chem. Eng. 4, 778469.
645 <https://doi.org/10.3389/fceng.2022.778469>

646 Morales, M., Aflalo, C., Bernard, O., 2021. Microalgal lipids: A review of lipids potential and
647 quantification for 95 phytoplankton species. Biomass and Bioenergy 150, 106108.
648 <https://doi.org/10.1016/j.biombioe.2021.106108>

649 Morin, S., Pesce, S., Tlili, A., Coste, M., Montuelle, B., 2010. Recovery potential of periphytic
650 communities in a river impacted by a vineyard watershed. Ecological Indicators 10, 419–426.
651 <https://doi.org/10.1016/j.ecolind.2009.07.008>

652 Muller, R., Schreiber, U., Escher, B.I., Quayle, P., Bengtson Nash, S.M., Mueller, J.F., 2008. Rapid
653 exposure assessment of PSII herbicides in surface water using a novel chlorophyll a
654 fluorescence imaging assay. Science of The Total Environment 401, 51–59.
655 <https://doi.org/10.1016/j.scitotenv.2008.02.062>

656 Müller-Navarra, D.C., Brett, M.T., Liston, A.M., Goldman, C.R., 2000. A highly unsaturated fatty acid
657 predicts carbon transfer between primary producers and consumers. Nature 403, 74–77.
658 <https://doi.org/10.1038/47469>

659 Murdock, J.N., Shields, F.D., Lizotte, R.E., 2013. Periphyton responses to nutrient and atrazine
660 mixtures introduced through agricultural runoff. Ecotoxicology 22, 215–230.
661 <https://doi.org/10.1007/s10646-012-1018-9>

662 Nakamura, Y., Li-Beisson, Y. (Eds.), 2016. Lipids in Plant and Algae Development, Subcellular
663 Biochemistry. Springer International Publishing, Cham. <https://doi.org/10.1007/978-3-319-25979-6>

664

665 Ou-Yang, K., Feng, T., Han, Y., Li, G., Li, J., Ma, H., 2022. Bioaccumulation, metabolism and endocrine-
666 reproductive effects of metolachlor and its S-enantiomer in adult zebrafish (*Danio rerio*).
667 Science of The Total Environment 802, 149826.
668 <https://doi.org/10.1016/j.scitotenv.2021.149826>

669 Pannard, A., Le Rouzic, B., Binet, F., 2009. Response of Phytoplankton Community to Low-Dose
670 Atrazine Exposure Combined with Phosphorus Fluctuations. Arch Environ Contam Toxicol 57,
671 50–59. <https://doi.org/10.1007/s00244-008-9245-z>

672 Parlakidis, P., Rodriguez, M.S., Gikas, G.D., Alexoudis, C., Perez-Rojas, G., Perez-Villanueva, M.,
673 Carrera, A.P., Fernández-Cirelli, A., Vryzas, Z., 2022. Occurrence of Banned and Currently
674 Used Herbicides, in Groundwater of Northern Greece: A Human Health Risk Assessment
675 Approach. *IJERPH* 19, 8877. <https://doi.org/10.3390/ijerph19148877>

676 Perkins, D.B., Chen, W., Jacobson, A., Stone, Z., White, M., Christensen, B., Ghebremichael, L., Brain,
677 R., 2021. Development of a mixed-source, single pesticide database for use in ecological risk
678 assessment: quality control and data standardization practices. *Environ Monit Assess* 193,
679 827. <https://doi.org/10.1007/s10661-021-09596-9>

680 Proia, L., Morin, S., Peipoch, M., Romani, A.M., Sabater, S., 2011. Resistance and recovery of river
681 biofilms receiving short pulses of Triclosan and Diuron. *Science of The Total Environment*
682 409, 3129–3137. <https://doi.org/10.1016/j.scitotenv.2011.05.013>

683 Prosser, R.S., Brain, R.A., Hosmer, A.J., Solomon, K.R., Hanson, M.L., 2013. Assessing sensitivity and
684 recovery of field-collected periphyton acutely exposed to atrazine using PSII inhibition under
685 laboratory conditions. *Ecotoxicology* 22, 1367–1383. <https://doi.org/10.1007/s10646-013-1123-4>

687 Quintaneiro, C., Patrício, D., Novais, S.C., Soares, A.M.V.M., Monteiro, M.S., 2017. Endocrine and
688 physiological effects of linuron and S-metolachlor in zebrafish developing embryos. *Science*
689 *of The Total Environment* 586, 390–400. <https://doi.org/10.1016/j.scitotenv.2016.11.153>

690 Rea, G., Polticelli, F., Antonacci, A., Scognamiglio, V., Katiyar, P., Kulkarni, S.A., Johanningmeier, U.,
691 Giardi, M.T., 2009. Structure-based design of novel *Chlamydomonas reinhardtii* D1-D2
692 photosynthetic proteins for herbicide monitoring. *Protein Science* 18, 2139–2151.
693 <https://doi.org/10.1002/pro.228>

694 Relyea, R.A., 2009. A cocktail of contaminants: how mixtures of pesticides at low concentrations
695 affect aquatic communities. *Oecologia* 159, 363–376. <https://doi.org/10.1007/s00442-008-1213-9>

697 Ricart, M., Barceló, D., Geiszinger, A., Guasch, H., Alda, M.L. de, Romani, A.M., Vidal, G., Villagrasa,
698 M., Sabater, S., 2009. Effects of low concentrations of the phenylurea herbicide diuron on
699 biofilm algae and bacteria. *Chemosphere* 76, 1392–1401.
700 <https://doi.org/10.1016/j.chemosphere.2009.06.017>

701 Rock, C.O., 2008. Fatty acid and phospholipid metabolism in prokaryotes, in: *Biochemistry of Lipids,*
702 *Lipoproteins and Membranes.* Elsevier, pp. 59–96. <https://doi.org/10.1016/B978-044453219-0.50005-2>

704 Rossoll, D., Bermúdez, R., Hauss, H., Schulz, K.G., Riebesell, U., Sommer, U., Winder, M., 2012. Ocean
705 Acidification-Induced Food Quality Deterioration Constrains Trophic Transfer. *PLoS ONE* 7,
706 e34737. <https://doi.org/10.1371/journal.pone.0034737>

707 Roubex, V., Fauvelle, V., Tison-Rosebery, J., Mazzella, N., Coste, M., Delmas, F., 2012. Assessing the
708 impact of chloroacetanilide herbicides and their metabolites on periphyton in the Leyre River
709 (SW France) via short term growth inhibition tests on autochthonous diatoms. *J. Environ.*
710 *Monit.* 14, 1655. <https://doi.org/10.1039/c2em10887a>

711 Schmitt-Jansen, M., Altenburger, R., 2005. Predicting and observing responses of algal communities
712 to photosystem ii–herbicide exposure using pollution-induced community tolerance and
713 species-sensitivity distributions. *Environ Toxicol Chem* 24, 304. <https://doi.org/10.1897/03-647.1>

715 Sénat de France, 2003. Annexe 47 (Atrazine) du rapport d’office parlementaire sur la qualité de l’eau
716 et assainissement en France.

717 Shanta, P.V., Li, B., Stuart, D.D., Cheng, Q., 2021. Lipidomic Profiling of Algae with Microarray MALDI-
718 MS toward Ecotoxicological Monitoring of Herbicide Exposure. *Environ. Sci. Technol.* 55,
719 10558–10568. <https://doi.org/10.1021/acs.est.1c01138>

720 Shen, P.-L., Wang, H.-T., Pan, Y.-F., Meng, Y.-Y., Wu, P.-C., Xue, S., 2016. Identification of
721 Characteristic Fatty Acids to Quantify Triacylglycerols in Microalgae. *Front. Plant Sci.* 7.
722 <https://doi.org/10.3389/fpls.2016.00162>

723 Špoljarić Maronić, D., Štolfa Čamagajevac, I., Horvatić, J., Žuna Pfeiffer, T., Stević, F., Žarković, N.,
724 Waeg, G., Jaganjac, M., 2018. S-metolachlor promotes oxidative stress in green microalga
725 *Parachlorella kessleri* - A potential environmental and health risk for higher organisms.
726 *Science of The Total Environment* 637–638, 41–49.
727 <https://doi.org/10.1016/j.scitotenv.2018.04.433>

728 Székács, A., Mörtl, M., Darvas, B., 2015. Monitoring Pesticide Residues in Surface and Ground Water
729 in Hungary: Surveys in 1990–2015. *Journal of Chemistry* 2015, 1–15.
730 <https://doi.org/10.1155/2015/717948>

731 Thakkar, M., Randhawa, V., Wei, L., 2013. Comparative responses of two species of marine
732 phytoplankton to metolachlor exposure. *Aquatic Toxicology* 126, 198–206.
733 <https://doi.org/10.1016/j.aquatox.2012.10.002>

734 Thompson, F.L., Abreu, P.C., Wasielesky, W., 2002. Importance of biofilm for water quality and
735 nourishment in intensive shrimp culture. *Aquaculture* 203, 263–278.
736 [https://doi.org/10.1016/S0044-8486\(01\)00642-1](https://doi.org/10.1016/S0044-8486(01)00642-1)

737 Trbojević, I., Jovanović, J., Kostić, D., Popović, S., Krizmanić, J., Karadžić, V., Subakov Simić, G., 2017.
738 Structure and succession of periphyton in an urban reservoir: artificial substrate specificity.
739 *Oceanological and Hydrobiological Studies* 46, 379–392. <https://doi.org/10.1515/ohs-2017-0038>

740

741 USEPA, 2017. Pesticides industry sales and usage, 2008-2012 Market estimates. United States
742 Environmental Protection Agency.

743 USEPA, 2016. Refined Ecological Risk Assessment for Atrazine. United States Environmental
744 Protection Agency.

745 Vallotton, N., Moser, D., Eggen, R.I.L., Junghans, M., Chèvre, N., 2008. S-metolachlor pulse exposure
746 on the alga *Scenedesmus vacuolatus*: Effects during exposure and the subsequent recovery.
747 *Chemosphere* 73, 395–400. <https://doi.org/10.1016/j.chemosphere.2008.05.039>

748 Vryzas, Z., Alexoudis, C., Vassiliou, G., Galanis, K., Papadopoulou-Mourkidou, E., 2011. Determination
749 and aquatic risk assessment of pesticide residues in riparian drainage canals in northeastern
750 Greece. *Ecotoxicology and Environmental Safety* 74, 174–181.
751 <https://doi.org/10.1016/j.ecoenv.2010.04.011>

752 Wetzel, R.G. (Ed.), 1983. *Periphyton of Freshwater Ecosystems: Proceedings of the First International*
753 *Workshop on Periphyton of Freshwater Ecosystems held in Växjö, Sweden, 14–17 September*
754 *1982*. Springer Netherlands, Dordrecht. <https://doi.org/10.1007/978-94-009-7293-3>

755 WSSA, 2021. WSSA-Herbicide Site of Action (SOA) Classification List. *Weed Science Society of*
756 *America*.

757 Zemolin, C.R., Avila, L.A., Cassol, G.V., Massey, J.H., Camargo, E.R., 2014. Environmental fate of S-
758 Metolachlor: a review. *Planta daninha* 32, 655–664. <https://doi.org/10.1590/S0100-83582014000300022>

759

760

761 **Statements and Declarations**

762 **Funding**

763 We would like to thank the Fonds de recherche du Québec (FRQNT) for a grant to I. Lavoie (FRQNT Relève
764 professorale; 2021-NC-285440) and the Centre de recherche en écotoxicologie du Québec (EcotoQ) for funding
765 to L. Malbezin.

766 **Conflict of interests**

767 The authors declare that they have no conflict of interest.
768

769 **Competing Interests**

770 None.

771

772 **Author contributions**

773 All authors made substantial contributions to this paper. L.M. was in charge of experimental conceptualization,
774 laboratory experiments, sample collection, data analysis and writing. S.M. was involved in experimental
775 conceptualization, project management, reviewing and editing. I.L. was responsible for funding acquisition,
776 project administration and was involved in the project conception, experimental conceptualization, reviewing and
777 editing.