

Supplemental Table.
Sex-specific associations between BMI and lung cancer, Montreal, Canada, 1996-2002

	Men			Women		
	N _{ca}	N _{co}	OR ¹ (95% CI)	N _{ca}	N _{co}	OR ¹ (95% CI)
BMI_{age 20}²						
Underweight	28	66	0.60 (0.35, 1.04)	86	126	0.72 (0.47, 1.11)
Normal	473	629	-	326	412	-
Overweight	118	138	1.22 (0.88, 1.70)	27	35	0.96 (0.46, 2.01)
Obese	9	16	0.52 (0.18, 1.52)	4	9	0.85 (0.19, 3.85)
BMI_{2 yrs ago}²						
Underweight	25	9	2.64 (1.07, 6.51)	38	19	2.41 (1.06, 5.45)
Normal	299	343	-	222	282	-
Overweight	220	391	0.75 (0.57, 0.98)	126	197	0.99 (0.65, 1.49)
Obese	84	106	0.94 (0.64, 1.39)	57	84	0.88 (0.51, 1.52)
BMI change						
Decrease by ≥ 1 category	43	22	2.78 (1.44, 5.38)	28	11	2.73 (1.01, 7.44)
No change	341	393	-	198	252	-
Increase by 1 category	189	356	0.70 (0.53, 0.92)	156	218	0.80 (0.53, 1.20)
Increase by > 1 category	55	78	0.97 (0.62, 1.52)	61	101	0.76 (0.45, 1.29)

¹ Adjusted for age, education, respondent status, ancestry, fruit and vegetables consumption, occupational exposures, recreational physical activity, alcohol, and CSI.

² Underweight (<18.5 kg/m²), normal (18.5-24.9 kg/m²), overweight (25.0-29.9 kg/m²), and obese (≥30 kg/m²).