

IL2R β -dependent signals drive terminal exhaustion and suppress memory development during chronic viral infection

Jean-Christophe Beltra^{a,b}, Sara Bourbonnais^a, Nathalie Bédard^c, Tania Charpentier^d, Moana Boulangé^{a,b},
Eva Michaud^{a,b}, Ines Boufaied^e, Julie Bruneau^{c,f}, Naglaa H. Shoukry^{c,g}, Alain Lamarre^d, and Hélène Decaluwe^{a,b,h,1}

^aCytokines and Adaptive Immunity Laboratory, Sainte-Justine University Hospital Research Center, Montreal, QC, Canada H3T 1C5; ^bDepartment of Microbiology and Immunology, Faculty of Medicine, University of Montreal, Montreal, QC, Canada H3C 3J7; ^cUniversity of Montreal Hospital Research Center, Montreal, QC, Canada H2X 0A9; ^dImmunovirology Laboratory, Institut National de la Recherche Scientifique-Institut Armand-Frappier, Laval, QC, Canada H7V 1B7; ^eFlow Cytometry Platform, Sainte-Justine University Hospital Research Center, QC, Canada H3T 1C5; ^fDepartment of Family Medicine and Emergency Medicine, Faculty of Medicine, University of Montreal, Montreal, QC, Canada H2X 0A9; ^gDepartment of Medicine, Faculty of Medicine, University of Montreal, Montreal, QC, Canada H3T 1J4; and ^hImmunology and Rheumatology Division, Department of Pediatrics, Faculty of Medicine, University of Montreal, Montreal, QC, Canada H3T 1C5

Edited by Michael B. A. Oldstone, The Scripps Research Institute, La Jolla, CA, and approved July 26, 2016 (received for review March 14, 2016)