

[Click here to view linked References](#)

1 **Title: Towards a multibioassay-based index for toxicity assessment of fluvial waters**

2
3
4 Lalit K. Pandey^{a,b}, Isabelle Lavoie^c, Soizic Morin^d, Stephen Depuydt^e, Lyu Jie^f, Hojun Lee^g, Jinho Jung^h,
5 Dong-Hyuk Yumⁱ, Taejun Han^{g,j}, Jihae Park^{e,*}

6
7 ^a*Institute of Green Environmental Research Center, 169, Gaetbeol-ro, Yeonsu-gu, Incheon 21999, Korea.*

8 ^b*Department of Plant Science, Faculty of Applied Sciences, MJP Rohilkhand University, Bareilly, 243006,*
9 *India.*

10 ^c*Institut national de la recherche scientifique, centre Eau Terre Environnement, 490 rue de la Couronne,*
11 *Québec City, Québec, G1K 9A9, Canada.*

12 ^d*Irstea, UR EABX, 50 avenue de Verdun 33612 Cestas cedex, France.*

13 ^e*Lab of Plant Growth Analysis, Ghent University Global Campus, 119, Songdomunwha-ro, Yeonsu-gu,*
14 *Incheon 21985, Korea*

15 ^f*Department of Life Sciences, Jilin Normal University, Siping City, Jilin Province, China*

16 ^g*Department of Marine Science, Incheon National University, 119, Academy-ro, Yeonsu-gu, Incheon*
17 *22012, Korea.*

18 ^h*Division of Environmental Science & Ecological Engineering, Korea University, Seoul 02841, Korea*

19 ⁱ*Ecotoxicology Team, Korea Institute of Toxicology, Daejeon 34114, Korea*

20 ^j*Ghent University Global Campus, 119, Songdomunwha-ro, Yeonsu-gu, Incheon 21985, Korea.*

21
22 *Corresponding author at:

23 Lab of Plant Growth Analysis, Ghent University Global Campus, 119, Songdomunwha-ro, Yeonsu-gu,
24 Incheon 21985, Republic of Korea. E-mail address: jihae.park@ghent.ac.kr / Telephone number: +82 32
25 626 4305 / ORCID: 0000-0002-1873-1480 (J. Park).

26
27
28
29 **Highlights**

- 30
31 - An ecotoxicological assessment of natural waters based on a multi-organism trial was conducted
32 in four South Korean rivers
33 - The tested organisms showed distinct levels of performance in their response to natural waters
34 - A scoring system is proposed to integrate biological responses into an overall toxicity category
35 - This bioassay approach identified more sites as potentially degraded as did water chemistry
36 measurements alone
37

38 **Abstract**

39 Despite their proven reliability for revealing “acceptable” degrees of toxicity in waste-
40 and reclaimed waters, bioassays are rarely used to assess the toxicity of hazardous
41 contaminants present in natural waters. In this study, we used organisms from different
42 trophic levels to assess the toxicity of water samples collected from four different South
43 Korean rivers. The main objective was to develop a multi-descriptor index of toxicity for
44 undiluted river water. The responses of six test organisms
45 (*Aliivibrio fischeri*, *Pseudokirchneriella subcapitata*, *Heterocypris incongruens*, *Moina m*
46 *acrocopa*, *Danio rerio*, and *Lemna minor*) after laboratory exposure to water samples
47 were considered for this index, as well as the frequency of teratologies in diatom
48 assemblages. Each individual test was attributed a toxicity class and score (three levels;
49 no toxicity = 0, low toxicity = 1, confirmed toxicity = 2) based on the organism’s
50 response after exposure and a total score was calculated. The proposed index also
51 considers the number of test organisms that received the highest toxicity score (value =2).
52 An overall toxicity category was then attributed to the water sample based on those two
53 metrics: A = no toxicity, B = slight toxicity, C = moderate toxicity; D = toxicity, E = high
54 toxicity. The susceptibility of the test organisms varied greatly and the sensitivity of their
55 response also differed among bioassays. The combined responses of organisms from
56 different trophic levels and with different life strategies provided multi-level diagnostic
57 information about the intensity and the nature of contamination.

58

59 **Keywords:** Aquatic plants; Bioassay; Biological indicators; Microorganisms; Multi-descriptor index;
60 Multiple endpoints; Receiving water

61

62 **1. Introduction**

63 Fluvial ecosystems experience multiple anthropogenic disturbances such as industrial, municipal
64 and agricultural effluents leading to eutrophication and chemical contamination, acidification, hydrological
65 and hydro-morphological alterations, and invasion by non-native species (Li et al. 2010). In particular,
66 wastewater effluents derived from industrial and municipal sources contain a mixture of chemicals which,
67 once released into the receiving water bodies, can have deleterious effects on the biological integrity of the
68 flora and fauna (Kim et al. 2015). These impacts are observable at different levels of biological
69 organisation (e.g., molecular, individual, population, and community; Nedeau et al. 2003; Ntengwe and
70 Maseka 2006; Tabrez and Ahmad 2012; Hassan et al. 2015) and may ultimately alter ecosystem function.
71 The most commonly used approach for assessing water quality is the measurement of chemical substances
72 and their metabolites. However, it is now generally accepted that there are potential limitations when
73 relying solely on chemical profiling to evaluate ecosystem health status, regulate acceptable loads of
74 wastewater effluents, and conduct risk assessments (Wolska et al. 2007). Routine chemical monitoring does
75 not account for the bioavailability of chemicals and nutrients, the temporal changes in exposure, or the
76 additive and synergistic effects of contaminants (Ahlf et al. 2002; Chu and Chow 2002). Most importantly,
77 chemistry-based monitoring does not provide information regarding the effect of contaminants on the biota.
78 Ecological surveillance of running waters based on an integrated assessment of the biological, chemical and
79 physical properties of a system contributes to better water resource protection and conservation and helps
80 water managers to plan rehabilitation.

81 Numerous water quality monitoring approaches based on organisms such as algae (mostly
82 diatoms) (e.g., Ponader et al. 2007; Coste et al. 2009; Kelly 2013; Lavoie et al. 2014), macrophytes (Small
83 et al. 1996; Thiebaut et al. 2002), invertebrates (e.g., Reynoldson et al. 1997; Lento et al. 2008; Canesi and
84 Corsi 2016) and fish (e.g., Joy and Death 2001; Oberdorff et al. 2002) are commonly used worldwide.
85 These indices are generally based on metrics such as species assemblage structure, diversity, % tolerant
86 species, life-forms, traits, size distribution, etc. However, these monitoring approaches have mostly been
87 developed to assess overall biological integrity, and often mostly reflect nutrient and organic matter
88 enrichment and habitat degradation, not toxicity. Biological assessment of potential water toxicity from

89 various types of inorganic and organic contaminants is usually performed using single and multi-species
90 toxicity tests which can effectively demonstrate causal relationships between the presence of contaminants
91 and adverse effects on the biota. Bioassays are widely used to assess the toxicity of a substance or a
92 combination of compounds in aquatic and terrestrial environments. As a general trend, this ecotoxicity
93 assessment approach relies on stepwise dilutions of samples to determine effective or lethal concentrations
94 (EC and LC, for example, the concentration of a substance or mixture giving half-maximal response of the
95 test organism is the EC₅₀), with the objective to establish acceptable degrees of toxicity before wastes can
96 be discharged in the environment or reclaimed. The information can then be used for monitoring and
97 predicting the effects of chemical discharges and for deriving chemical-specific water quality guidelines
98 (Ankley et al. 1992). Bioassays, such as bioluminescence in the bacteria *Vibrio fischeri*, the cell count of
99 microalgae such as *Pseudokirchneriella subcapitata*, photosynthetic activity of microalgal communities
100 (Kim Tiam et al. 2016), mortality and growth of small invertebrates such as ostracods and cladocerans, and
101 survival of fish such as *Danio rerio* (zebrafish) are internationally standardized test methods commonly
102 used in ecotoxicology. Bioassays can also be used to monitor potential toxicity of natural waters, although
103 this approach is not as conventional and necessitates a different method for the determination of the
104 contamination level than the EC or LC generally used for wastewaters.

105 In this paper, assessment of natural water toxicity is proposed using undiluted river samples and a
106 battery of biological indicators. Including various organisms in toxicity evaluation ensure a better-
107 integrated response by providing information at different levels (e.g., biochemical function, cellular growth,
108 mortality, etc.). The various test organisms have different sensitivities to the suite of contaminants, which
109 in turn maximises chances to detect a response after exposure to a sample with unknown chemical
110 composition. The goal of our research was to develop a multi-species toxicity test procedure using six
111 commonly used test organisms (*Aliivibrio fischeri* (bacterium), *Pseudokirchneriella subcapitata* (green
112 microalga), *Heterocypris incongruens* (ostracod), *Moina macrocopa* (cladocera), *Danio rerio* (fish), *Lemna*
113 *minor* (aquatic macrophyte)) for ecotoxicological assessments of river waters collected at different sites
114 distributed among four watersheds impacted by industrial and municipal effluents in South Korea. In
115 addition, the frequency of teratologies (abnormalities) in diatoms (siliceous brown microalga) collected *in*
116 *situ* was also included in this multi-species toxicity assessment.

117 This study proposes an approach for a rapid screening of flowing waters with suspected toxic
118 contamination, and provides valuable knowledge for further development of a bioassay-based index for
119 future implementation in the national river water quality management program in South Korea.

120 **2. Materials and methods**

121 **2.1. Water sample collection and on-site field measurements**

122 Sites were selected based on an existing micropollutant dataset from an eight-year extensive
123 monitoring program, conducted by the South Korean government, to assess river water quality at 159 sites
124 receiving effluents from 34 industrial complexes (Cho et al. 2014) (Tables S1 and S2). The South Korean
125 Ministry of Environment (MOE) also performs regular water quality assessments of streams and rivers,
126 including measurements of metals (As, Ag, Cd, Cr, Cu, Hg, Ni, Pb, and Zn), total phosphorus (TP) and
127 total nitrogen (TN), biological oxygen demand (BOD₅), total dissolved solids (TDS), pH, conductivity, and
128 total coliforms. Based on the available information on river water quality, 16 sites were chosen to cover
129 various sub-watersheds across South Korea. Additional information on the sampling sites selected and on
130 the four watersheds is provided in supplementary tables (Tables S1 and S2).

131 Surface water samples were collected in plastic bottles (2 litres) at the 16 selected sites distributed
132 along four major South Korean rivers in September 2015 (Fig. 1). The samples were kept cold during
133 sampling and transportation and were stored at 4°C upon arrival at the laboratory. Temperature, pH, and
134 dissolved oxygen were measured directly in the field with a multi-parameter display system (YSI 650,
135 USA). Conductivity was also measured on-site using a portable metre (Milwaukee, USA), and total
136 dissolved solids measurements were performed with a field amperometric graphite electrode (Hanna
137 HI98301 DiST® 1, USA).

138 **2.2. Chemical analyses**

139 Biochemical oxygen demand was measured by dark and light bottle incubation for 24 h at 20°C in
140 controlled light conditions, and Colilert-18 tests (IDEXX) were carried out to estimate the total number of
141 coliforms. Water samples were filtered through 0.45 µm syringe filters prior to analyses of nutrients and
142 metals. An automatic water analyser (Skalar/ Netherlands, SAN ++) was used to measure TN and TP.
143 Analyses of metals were conducted by inductively coupled plasma-optical emission spectrometry (ICP-
144 OES; Varian Vista PRO, CA, USA). Standard solutions were prepared fresh and calibration curves ($r^2 >$

145 0.995) were generated daily. Standard solutions were analysed after every 10 samples to verify their
146 concentrations. Measurement precision ranged from 94 to 107%, and detection limits were calculated based
147 on the standard deviations of blanks triplicates (range: 4 to 14 $\mu\text{g L}^{-1}$). Organic compounds were analysed
148 by the South Korean Ministry of Environment, following the methods presented by Cho *et al.* (2014).

149 **2.3. Toxicity assessment**

150 **2.3.1. Microtox bioassay (bacteria)**

151 The Microtox[®] (Newark, DE, USA) bacterial acute toxicity assay was used to determine inhibition
152 in the metabolism of *Aliivibrio fischeri* when exposed for 30 minutes to each of the 16 surface water
153 samples. The cultures were maintained at 15°C throughout the experiment. The control treatment consisted
154 of 2% NaCl water. Bacteria were exposed in triplicates to each treatment, according to the standard
155 Microtox procedure (Tarkpea and Hansson 1989). Bioluminescence of the bacteria was measured with a
156 Microtox photometer (Model 500). Results were expressed as % inhibition in bioluminescence between
157 pre-exposure to the water samples and after 30 minutes of exposure, taking into account the temporal
158 changes in luminescence occurring in control (ISO 11348-1:2009; Parvez et al. 2006). Confidence intervals
159 (CI) were calculated for α -value = 0.05.

160 **2.3.2. Microalgal bioassay**

161 The test performed in triplicate on *Pseudokirchneriella subcapitata* was conducted using an
162 Algaltoxkit (Microbiotests, Belgium). The methods followed standard operational procedures provided by
163 the manufacturer and in accordance with the OECD Test Guideline 201 (OECD 1984). Microplates were
164 filled with 900 μL of test water (for the control, 900 μL of MBL medium (Nichols 1973) was used) and 100
165 μL of an algal-inoculum solution, with the initial number of algal cells being adjusted to 10^5 cells mL^{-1} . The
166 exposure tests were performed for 72 h at 25°C under light conditions of 60-80 $\mu\text{mol photons m}^{-2} \text{s}^{-1}$.
167 Determination of algal growth was conducted using absorbance values measured at a 670 nm wavelength
168 using a spectrophotometer (Scinco, S-3100, Korea). Optical densities of the blank and samples were taken
169 at 670 nm and then converted into cell densities (cells mL^{-1}). The growth inhibition (% calculated as the
170 percentage reduction in average specific growth rate compared to the control, following OECD 1984) was
171 used as an indicator of the toxicity of the 16 river water samples. Confidence intervals ($\alpha = 0.05$) were
172 calculated.

173 **2.3.3. Ostracod bioassay**

174 Hatching of ostracod (*Heterocypris incongruens*) cysts was initiated 48 h prior to the start of the
175 toxicity test, and freshly hatched ostracods were pre-fed (with Spirulina powder from Spirulina
176 International, Tilburg, Holland) for 4h. The toxicity test followed the procedure described in the
177 International Standard ISO 14371:2012. Pore sediment water was collected at each site and sand was used
178 as reference sediment following the protocol of Chial and Persoone (2002a). Multi-well plates were filled
179 with 100 µL of sediment and 2 ml of overlying water (distilled water for the control). *Pseudokirchneriella*
180 *subcapitata* was provided as food source. Ten ostracod eggs per treatment were placed in the dark at 25°C
181 for six days. Water was not replaced during the experiment—that is, the experiment was static. Each of the
182 16 river water treatments and the control was tested in triplicate. Ostracods were recovered from the multi-
183 well plates at the end of the six-day exposure to determine % mortality and growth length inhibition and
184 were compared to the controls. For both endpoints, confidence intervals ($\alpha = 0.05$) were calculated.

185 **2.3.4. Cladocera bioassay**

186 Toxicity effects of the 16 river water samples on *Moina macrocopa* (≤ 24 h old) were tested after
187 adaptation of the OECD Test Guideline 211 (OECD 2012) for *Daphnia magna* to better reflect conditions
188 appropriate to *M. macrocopa* intrinsic population dynamics. Each toxicity test was conducted on 10
189 neonate female cladocerans, with one animal per well. Elendt M4 medium (Elendt 1990) was used as the
190 control water. Exposure was performed for nine days at $20 \pm 2^\circ\text{C}$ with 16 h light and 8 h dark photoperiods.
191 The test medium (river water) was renewed every three days and the cladocerans were fed daily *ad libitum*
192 with green algae (*Chlorella vulgaris*). Offspring production per female was recorded and expressed as % of
193 control values (CI: $\alpha = 0.05$). Growth inhibition induced by the treatment, compared to the control, was
194 determined from individual length measurements. Mortality was determined as the percentage of dead
195 individuals at the end of the exposure.

196 **2.3.5. Fish embryo test**

197 The fish embryo toxicity test (FET; OECD Test Guideline 236 2013) was conducted with *Danio*
198 *rerio*. A total of 20 fertilised eggs for each treatment were incubated at 20°C for 96 h. Percent mortality
199 was evaluated based on the numbers of coagulated embryos or lack of heartbeat at the end of the

200 experiment. Criteria to establish the validity of the test were reached, with > 90% survival in the negative
201 controls, and 100% mortality in the positive control (3,4-dichloroaniline, 4 mg L⁻¹).

202 **2.3.6. *Lemna* root bioassay**

203 A *Lemna* root bioassay was adapted from Park et al. (2017). Fresh green fronds of *Lemna minor*,
204 consisting of two fronds, were selected as the test material. Roots were excised, and rootless plants were
205 placed in a well plate with 3.0 ml of test water (3.0 ml of Steinberg medium for the control; Steinberg
206 1946). The experiment was conducted using 20 fronds per treatment. Toxicity tests were carried out in a
207 growth chamber for 72 h at 25°C under continuous light (90-100 μmol photons m⁻² s⁻¹). Plants were
208 examined after the 72 h incubation period and the root lengths were measured using an imaging analyser
209 (Moticam 2500, Ted Pella Inc., USA). Root elongation inhibition was then expressed as the % reduction in
210 the root length of exposed *Lemna* (measured in mm), compared to the control (CI: α = 0.05).

211 **2.3.7. Deformity assessment in diatom frustules**

212 The occurrence of deformities was assessed using on-site diatom assemblages chronically exposed
213 to river water impacted with industrial discharges. The methods for biofilm collection, sample preparation,
214 slide mounting, and deformity evaluation are explained in Pandey and Bergey (2016), Cerisier et al. (2018)
215 and Pandey et al. (2018). Briefly, biofilm samples were collected at each site by scraping hard substrates
216 such as stones and concrete walls (~ 25 cm²) using a blade and a brush. Diatom valve deformities were
217 enumerated using digested material mounted onto permanent microscope slides and observed at 1000x
218 magnification under an oil immersion microscope (Carl Zeiss, Axiostar plus, Germany). The proportion of
219 deformed diatom valves at each site was estimated based on 500 valve counts and expressed as %
220 deformity (CI: α = 0.05). For comparison purposes, biofilms were also sampled in least-impacted areas
221 upstream of the major sources of contamination (three sites and four samples per river), and these
222 “reference” diatom assemblages were examined for deformities. The average % deformity observed in the
223 upstream site samples was used as the control.

224 **2.3.8. Scoring of bioassay results**

225 For each test, three toxicity classes were established to estimate the level of contamination: no
226 toxicity, moderate toxicity, and high toxicity (Table 1). For all of the bioassays, > 50% response (i.e., above
227 the EC₅₀ or LC₅₀) was considered as the lower boundary for “high toxicity” of the water samples, in line

228 with classical toxicity criteria. This 50% effect threshold is commonly reported as a significant acute
229 toxicity value in the bioluminescence assay (Niemiryecz et al. 2007; ISO 11348:2009), algal growth
230 inhibition test (OECD 1984), *Lemna* root regrowth test (Park et al. 2017), invertebrate mortality bioassays
231 (ISO-14371 2012; OECD 2012), and fish embryo toxicity test (OECD 2013). Samples resulting in a 20-
232 50% toxicity response were assigned to the “moderate toxicity” class, in agreement with the test validity
233 criteria defined for *A. fischeri* bioluminescence (Niemiryecz et al. 2007) or for ostracod and cladocera
234 mortality (ISO-14371 2012; OECD 2012). For the purpose of the present study, a 20% effect boundary was
235 used for the *Lemna* re-growth tests for which validity criteria have not yet been established. This criterion
236 ensures consistency with the results from the multiple-organisms trial and is in line with the hazard
237 classification system proposed by Persoone et al. (2003), which recommends the use of a 20% effect level
238 as the lowest significant toxic impact value. However, this threshold was lowered to 10% in the *Danio*
239 embryo tests, as per OECD (2013). The criteria for abnormal diatom valves (% teratologies) included three
240 categories with narrow ranges. The upper limit for the “no toxicity” class was set at < 0.5%, which is
241 regarded as the baseline in natural conditions, as suggested by Morin et al. (2008a) and Arini et al. (2012).
242 Teratology percentages ranging between 0.5-1% were assigned to the category “moderate toxicity” based
243 on previous observations (Lavoie et al. 2012; Morin et al. 2012; Pandey et al. 2014, 2015; Pandey and
244 Bergey 2016), while values above 1% were assigned to the category “high toxicity”. The toxicity classes
245 for diatom teratologies are subjected to modifications due to insufficient information and data regarding the
246 presence and type of abnormalities as a response to stress along a contamination gradient (Lavoie et al.
247 2017).

248 **2.4 Overall toxicity assessment**

249 A score was assigned to each toxicity class (“no toxicity” = 0, “moderate toxicity” = 1, and “high
250 toxicity” = 2), allowing for the calculation of a total score integrating the response of all organisms tested.
251 Where multiple endpoints were tested on an organism (reproduction, growth, and mortality), it was the
252 worst toxicity class observed that was considered for the calculation of the final score. Here, the maximum
253 total score possible is 14, based on the seven biological descriptors included in the present study. However,
254 this value will differ as a function of the number of bioassays tested in future bioassessments. The total
255 toxicity value is then reported as a percentage. Low values indicate no or low toxicity. Finally, each sample

256 was assigned an overall toxicity category by considering both their toxicity class and the number of tests
257 (expressed as %) that obtained a score of 2, “high toxicity” (Table 2). For example, a sample which would
258 obtain the following scores for each of the seven tests: 0, 2, 1, 0, 1, 0, and 2, would have a total score of 6
259 out of a maximum of 14 ($6/14 \times 100 = 43\%$). Two out of the seven tests would have been classified as “high
260 toxicity” (score value = 2), which represents 29% of the cases ($2/7 \times 100 = 29\%$). Following the criteria
261 proposed in Table 2, this sample would be categorised as “toxic” (D).

262 **3. Results and Discussion**

263 **3.1. Physico-chemical assessment of water quality**

264 The purpose of this study was to evaluate the potential of various toxicity bioassays for testing
265 water toxicity in natural freshwaters and to develop a scoring system allowing for an integrated assessment
266 of aquatic ecosystems health. An exhaustive presentation of the chemical and physical properties of the
267 tested waters is thus beyond the scope of this paper. Briefly, analysis of the physicochemical characteristics
268 of the 16 sites sampled (Table 3) revealed that all waters were nutrient-enriched. Particularly elevated
269 nitrogen concentrations and high conductivities were found at N2 and N3 (Nakdong River), as well as at
270 H1 and H2 (Han River). Water quality data for these four sites (N2 and N3; H1 and H2) represented the
271 most severely impacted conditions with particularly high suspended solids and elevated metal
272 concentrations (zinc at H1 and H2, Han River; copper and nickel at N2 and N3, Nakdong River). The
273 remaining sites were less impacted, based on the data available. However, it is worth highlighting the high
274 TP values measured in the samples collected on the Geum River, in particular, G1, G2, G3, and G4 (TP =
275 $0.28 \pm 0.03 \text{ mg L}^{-1}$). Organic compound analyses were also conducted during this survey, and
276 concentrations were below detection limits (data not shown). The fact that organic contaminants were not
277 detected is surprising because these ecosystems receive wastewater effluents from 34 industrial complexes
278 distributed along the four main rivers studied. Moreover, Cho et al. (2014) published the results from an
279 extensive survey conducted by the South Korean government and reported the presence of numerous toxic
280 chemicals (mainly organochlorine and organophosphate pesticides, volatile organic compounds, solvents,
281 and plasticisers) in these environments. As stated previously, traditional water chemistry measurements are
282 not always reliable to fully characterise environmental conditions because they do not integrate fluctuations
283 in water quality, and they are susceptible to missing intermittent contamination unless samples are collected

284 during wastewater effluent release or high-resolution analyses are performed. It is also possible that the low
285 concentrations of organic contaminants observed in the present study, compared with the values reported in
286 Cho et al. (2014), reflect the efforts of the South Korean government to reduce contaminant inputs from
287 industrial effluents.

288 **3.2. Bioassay-based toxicity assessment**

289 According to a report on a survey and evaluation of aquatic ecosystem health in South Korea
290 (Hwang et al. 2011), some rivers and streams are considerably contaminated in the country and are in
291 worse biological condition than predicted by conventional chemical analysis data. As a result, the country
292 recently adopted the use of bioindicators to improve water quality evaluation as part of a more integrated
293 concept of ecological status assessment. The use of biological descriptors is now part of the surface water
294 management policy in South Korea with the creation of the “Nationwide Aquatic Ecological Monitoring
295 Program (NAEMP)”. This new monitoring program is similar to the Water Framework Directive
296 established in the European Union (2000/60/EC), where the use of organisms from multiple biological
297 compartments (different trophic levels and different life strategies) is recommended in addition to
298 chemical, physical, and bacteriological parameters for examining the ecological status of fluvial
299 ecosystems (Geiszinger et al. 2009). Data gathered from the multiple bioassays used in this study provide
300 preliminary considerations for the development of a biological index of river water toxicity and serve as a
301 foundation for future optimisation of the approach. The results from each test are presented in Table 4,
302 along with CI and F-statistics from analysis of variance (ANOVA). The overall toxicity index values for
303 the 16 tested waters are reported in Table 5, along with individual scores obtained for each test organism.
304 The toxicity categories were attributed based on the criteria presented in Tables 1 and 2. The results for all
305 biological descriptors, as well as the final toxicity categories, are presented in detail and discussed in the
306 following paragraphs.

307 *3.2.1. Inhibition of bioluminescence*

308 Bacterial bioluminescence inhibition assays are commonly used to evaluate the toxicity of
309 contaminants released by wastewater effluents (e.g., Rodrigues and Umbuzeiro 2011). Here, inhibition of
310 bioluminescence in *Aliivibrio fischeri*, after exposure to the 16 river water samples, was used to evaluate its
311 potential in determining the toxicity of natural river waters. Significant luminescence inhibitions were only

312 revealed for the N2 and G2 sites, with changes of -28% (CI = 3%) and -16% (CI = 3%), respectively (Table
313 4). Based on the criteria established for this bioassay (Table 1 and Table 2), almost all sites were
314 categorised as “no toxicity”, except N2, which exhibited “moderate toxicity” (Table 5). The higher metal
315 concentrations at N2 were probably responsible for the observed toxicity, potentially in combination with
316 other toxic organic compounds suspected to contaminate these rivers (according to Cho et al. (2014)).
317 Because the presence and concentrations of the various contaminants may vary greatly, depending on
318 various factors such as water level and timing of wastewater effluent release in the river water, it is possible
319 that the other water samples were not as toxic to bacteria as was previously observed (MOE 2007). It is
320 also possible that *Aliivibrio fischeri* is simply not very sensitive to the type and level of contamination
321 tested in this study, as was observed by Macken et al. (2009) with the same bacterium shown to be the least
322 sensitive indicator tested for Cd contamination. Stimulation of bacterial metabolism was also observed in
323 the present study, with a significant increase in bioluminescence for the water sample collected at H2 (19%,
324 CI = 15%). This increased bioluminescence, compared to the control, may result from interferences caused
325 by the presence of volatile or insoluble substances in the waters (ISO 11348-1), or from a biological
326 response. Indeed, the presence of higher (but non-lethal) concentrations of metals, nutrients, and organic
327 matter in the waters tested compared to the control may, in turn, favour bacterial metabolism. For example,
328 bioluminescence stimulation of *Aliivibrio* species was also observed under metal exposure in other studies
329 (Fulladosa et al. 2005, 2007; Shen et al. 2009) and was attributed to hormesis (Calabrese 2005).

330 Based on the lack of a clear response of the bacteria to the test waters in this study, i.e., no strong
331 inhibition observed and stimulation of bioluminescence noted for numerous samples, this bioassay does not
332 seem to be sensitive enough nor appropriate when used alone for routine biological assessments of flowing
333 water toxicity. Becouze-Lareure et al. (2016) came to the same conclusion when assessing water and
334 sediment quality of a peri-urban river subjected to combined sewer overflow, which is also supported by
335 others (Angerville 2009; Gonzalez-Merchan et al. 2014a, b). This acute bioluminescence test is usually
336 carried out with an exposure time of 30 minutes or less, which does not provide information on potential
337 adverse effects that may happen later (e.g., Backhaus et al. 1997). For example, Hsieh et al. (2004)
338 observed that chronic exposure (22 h) of *Aliivibrio fischeri* to seven priority pollutant metals showed a

339 toxic response at concentrations many-fold lower than for acute exposure (5 or 15 min). This suggests that
340 toxicity assessments based on bioluminescence could be better evaluated with longer exposure times.

341 3.2.2. Algal cell density

342 The *Pseudokirchneriella subcapitata* bioassay was selected for toxicity assessment in the present
343 study for various reasons. This species is easily available from culture collections and easily maintained
344 under laboratory conditions. It is among the most widely recommended species for freshwater toxicity
345 testing, with standard guidelines having been established (OECD 1984; Environment Canada 1992;
346 USEPA 1994), and this bioassay is currently being endorsed for regulatory purposes. Moreover, the
347 responses of *P. subcapitata* to a variety of contaminants and its relative sensitivity compared with other test
348 organisms have been studied extensively (e.g., Radix et al. 2000; Weyers et al. 2000).

349 Algal cells exposed to the 16 test water samples showed significant growth inhibition (Table 4)
350 when exposed to the water samples from H1 (-30%, CI = 8%) and N1 (-11%, CI = 5%), compared to the
351 controls (growth rate: 0.035 div h⁻¹). Based on the selected toxicity thresholds, only H1 fell into the
352 “moderate toxicity” class (Table 5). These results are in contrast with the numerous reports of good
353 sensitivity of this test organism. For example, Katsumata et al. (2006) reported lower cell counts in *P.*
354 *subcapitata* when exposed to two herbicides (simazine and 3,5-dichlorophenol). However, it is most likely
355 that those pesticides have a more noticeable effect on algae (as their mode of action directly targets plant
356 functions) than do contaminants of largely industrial and municipal origin, as in the present study. Moreira-
357 Santos et al. (2004) reported a negative effect of a mine effluent (low pH, metals, and turbidity), with
358 growth inhibition reaching 98%, also suggesting good sensitivity of algal bioassays. The metal
359 concentrations in their study were much higher than the values observed in our selected South Korean
360 rivers which could explain why the magnitude of the effects we observed was lower. Moreover, acid mine
361 drainage (pH = 3) also likely inhibited algal growth in their study. The water samples from the Yeongsan
362 and Geum Rivers significantly stimulated algal growth ($p < 0.005$), which can most likely be attributed to
363 their nutrient-enriched waters (e.g., higher TP in the Geum River, see Table 3). This result suggests that the
364 potential toxicity of the samples may be overshadowed by the positive effect of nutrients on algae.
365 Although this algal bioassay does not seem to be very sensitive to the toxicity level and type of
366 contamination characterising the tested waters in the present study, it is worth noting that an increase in cell

367 density may provide information on the eutrophication potential of the waters. In other words, the fact that
368 most of the tested waters scored “no toxicity” does not guarantee good overall water quality.

369 3.2.3. Toxicity to Ostracoda

370 Freshwater ostracods are reported to be excellent bioindicators of surrounding physico-chemical
371 conditions and anthropogenic stressors (pesticides, hydrocarbons, and metals) (Ruiz et al. 2013). Mortality
372 in *Heterocypris incongruens* is a widely-used endpoint in ecotoxicology and constitutes a standardised
373 bioassay recognised by the International Organization for Standardization (ISO 2012). Ostracods in
374 bioassays are generally used to assess toxicity of solid phase matrices such as sediments or storm water
375 run-off particles (e.g., Chial and Persoone 2002a, b; Watanabe et al. 2008, 2011; Angerville et al. 2013)
376 because this organism spends most of its life in contact with sediments. However, this test has also been
377 used to assess the toxicity of various chemicals and to evaluate the toxicity potential of natural waters
378 (Toumi et al. 2015).

379 After six days of exposure to the tested waters, *Heterocypris incongruens* showed a mortality rate
380 higher than the 20% of the control. Mortality was significantly higher at H1 (with 100% mortality on the
381 date of recovery, $p < 0.0001$) and lower at G5 and G6 ($\leq 10\%$, $p < 0.05$). Concerning growth, cell
382 elongation in control was higher than expected, with final ostracod lengths 2.8 (± 0.2)-fold the initial sizes.
383 With exposure to the tested waters, growth inhibition compared to the controls was highlighted at G2 (44%,
384 CI = 19%), G3 (45%, CI = 37%), and Y1 (52%, CI = 19%). Based on the criteria presented in Table 1, H1
385 and Y1 received the “high toxicity” score, while G3 and G2 were classified in the “moderate toxicity”
386 category. The remaining sites were classified as “no toxicity”. Compared to the bacterial and algal
387 bioassays, the response of the ostracods suggests that they have a higher sensitivity to the toxicity of the
388 samples, with four sites highlighting toxicity, including two with the highest score (Table 5). In a study on
389 the effects of metals on *H. incongruens*, Sevilla et al. (2014) observed that aquatic exposure to different
390 concentrations of dissolved Cd (3.2-339 $\mu\text{g L}^{-1}$) and Cu (260-2600 $\mu\text{g L}^{-1}$) resulted in high mortalities (57-
391 100% and 95-100%, respectively). Watanabe et al. (2011) also reported high mortality in ostracods exposed
392 to elevated concentrations of metals (Cu, Zn, Ni, As, Cd, and Pb). Based on water chemistry measurements
393 available for this study, metal concentrations were much lower than the above-mentioned values, which
394 could partly explain the lower mortality rates observed. Moreover, the toxicity found using the ostracod

395 bioassay in this study may also result from exposure to other substances that were not measured by the
396 targeted micropollutant analyses, or it may reflect an integrated response to a cocktail of multiple
397 chemicals.

398 3.2.4. Toxicity to Cladocera

399 The *Daphnia magna* acute toxicity test is a common standard protocol to help regulate wastewater
400 effluents. However, this organism has not been found in aquatic systems in Korea. Thus, it has been
401 suggested that a domestic species, rather than the international standard species, be used for the Korean
402 whole effluent toxicity criteria. *Moina macrocopa*, which has been widely used in ecotoxicology
403 applications, is one of the most promising domestic species for this purpose. Kim et al. (2012) found that
404 effluents discharged from wastewater treatment plants in Korea induce multi-level toxicity, including acute
405 toxicity, feeding rate inhibition, and oxidative stress in *M. macrocopa*. Yi et al. (2010) found that *M.*
406 *macrocopa* was more sensitive to toxins than *Daphnia magna*, based on an acute toxicity test with
407 industrial effluents. Ji et al. (2008) noted significantly higher % mortality (50-100%) in *M. macrocopa*
408 when exposed to PFOS (perfluorooctane sulfonic acid) and PFOA (perfluorooctanoic acid) than in the
409 control treatment. These results show the sensitivity of this cladoceran species when exposed to high
410 concentrations of contaminants or when directly incubated with wastewaters. However, in the context of
411 the present study, the contamination levels of the tested waters were undoubtedly lower, which could
412 explain the fact that most samples were categorised as “no toxicity” (Table 5). Mortality assessment was
413 invalidated by the 22% mortality in the controls. Growth was significantly affected by exposure to the
414 tested waters ($F = 4.49; p \leq 0.001$), with significant growth inhibition at N1, H2, Y3, Y1, N3, G5, G6, and
415 Y4, compared to the control. However, despite significant differences with the control, growth inhibition
416 was always lower than 20% and therefore did not suggest toxicity. Contrastingly, reproduction expressed as
417 offspring production per female was inhibited by 44% (CI = 8%) at Y4, falling into the “moderate toxicity”
418 category (Table 5). Nine-day exposure to this water sample also led to 50% mortality (although this
419 endpoint was not considered due to >20% mortality in control). Interestingly, significant increases in
420 offspring production were observed in G1, G3, and H1 ($p \leq 0.05$), where survival was 100%. The same
421 pattern, although not significant, occurred at N3. This suggests that *Moina macrocopa* was indifferent to, or
422 even stimulated by the substances present in these waters. It must be noted, however, that the test was not

423 replicated and was invalidated due to high mortality under control condition. Based on the reproduction of
424 *M. macrocopa*, all the natural waters tested reflected a non-toxic environment, except site Y4 in the
425 Yeongsan River.

426 3.2.5. Fish bioassay

427 Worldwide, zebrafish has proven to be popular for toxicity evaluations because they are easy to
428 keep and to breed in the laboratory, are free of pathogenic microorganisms, and deliver eggs of high quality
429 independent of the season (Bresch 1991). For the purpose of this study, fish embryos were incubated with
430 river waters and toxicity was estimated based on % mortality. The results suggest that the tested waters are
431 generally of “moderate toxicity”, based on the selected criteria (Table 5), except for the samples from N1,
432 Y4, G1, G4, and G6, which exhibited < 10% mortality (Table 4). It must be noted that this exposure test
433 was not replicated. One hundred percent of surviving eggs hatched within 96 hours, irrespective of the
434 treatment.

435 In line with numerous studies showing the sensitive response of zebrafish embryos to wastewaters
436 of differing compositions (e.g., Şişman *et al.* 2008; Vincze *et al.* 2014), the results obtained here from
437 exposure to natural waters showed some toxicity for more than 50% of the samples. This finding may
438 reflect the sensitivity of fish embryos to moderate contamination and/or to a wide variety of substances.
439 However, it is worth underscoring the fact that, following the OECD (2013) specifications, we used a more
440 conservative effect value (10%) as the lower boundary for the “moderate toxicity” class.

441 3.2.6. Lemna root re-growth

442 The small size, structural simplicity, and rapid growth of *Lemna* are some of the characteristics
443 that make it advantageous for use in laboratory toxicity tests (Park *et al.* 2017). Moreover, this plant is an
444 essential primary producer and has a wide geographical distribution. Bioassays using *Lemna* as a test
445 organism are traditionally based on endpoints such as the number of fronds, their growth rate and their wet
446 or dry biomass, which require standard exposure durations of at least seven days to detect toxicity (Park *et al.*
447 2017). Roots of *Lemna minor* are highly sensitive to environmental stressors (Panda and Upadhyay
448 2003). However, there have been few studies incorporating root elongation as a test endpoint, partly due to
449 the fragility of the roots, which introduces difficulties into measurements (Davis 1981), and to the difficulty
450 of obtaining a sufficient number of test specimens with identical root lengths for exposure experiments.

451 The use of *Lemna* in ecotoxicology assessments has thus been re-evaluated (Gopalapillai et al. 2014; Park
452 et al. 2017) using root re-growth (after cutting them at the base) as an endpoint. This method has been
453 shown to be sensitive, precise, and ecologically significant in comparison with more traditional
454 measurements such as frond growth and biomass.

455 Root elongation under control conditions was 34 ± 4 mm. Compared to the controls, exposure of
456 *L. minor* to the 16 water samples collected in South Korean rivers showed significant toxicity for six sites
457 (Table 4). Samples from the Han River (H1 and H2) fell into the category “high toxicity”, and “moderate
458 toxicity” was noted for N2, N3, Y4, and G1; while the remaining 10 sites were considered “non-toxic”
459 (Table 5). The observed root re-growth inhibition may be a response to elevated zinc concentration in the
460 Han River, as this bioassay was shown to respond to metal contamination. For example, Park et al. (2013)
461 reported root elongation in three *Lemna* spp. exposed to Ag, Cd, Cr, Cu, and Hg to be a sensitive endpoint
462 to assess metal toxicity. However, in the sites where “moderate toxicity” was observed, metals do not seem
463 to represent a contamination concern, at least not based on measurements of punctual water samples taken
464 for these experiments. We cannot exclude a contribution of organic contaminants (and additive/synergistic
465 effects) to the observed toxicity, which may not have been detected owing to concentrations being below
466 the detection limit and/or because they were not targeted in the analysis (unknown compounds such as
467 degradation products).

468 3.2.7. *Diatom teratologies*

469 Percent deformed frustules were examined from periphytic diatom samples collected at the 16
470 sites (Fig. 2). Diatom deformities were found at high abundance at H2 ($5.9 \pm 0.7\%$) and H1 ($4.9 \pm 0.4\%$).
471 N2, Y1, and Y3 also showed potential toxicity, with deformities present at $> 1\%$. Based on the categories
472 tentatively suggested for this study, these teratology occurrences would place these five samples into the
473 category “high toxicity”. A marginal increase in diatom teratologies was also noticed at N1 ($0.8 \pm 0.7\%$)
474 and N3 ($0.7 \pm 0.1\%$), which were thus categorised as being of “moderate toxicity” (Table 5).

475 The approach based on diatom teratologies differs markedly from the six bioassays presented
476 above because it is conducted with organisms collected *in situ*, and therefore reflects an integrated response
477 to environmental fluctuations over a longer period of time. Moreover, the assessment of toxicity is not
478 based on the level of inhibition or mortality compared with a control. Rather, control conditions are

479 estimated to be the % of teratologies generally encountered in natural systems with minimal disturbance
480 (observed to be $\leq 0.5\%$, based on Morin et al. 2012). In this study, the % deformities averaged for three
481 sites (four replicates per site) sampled upstream of contamination was $0.3 \pm 0.1\%$. One particular advantage
482 of using teratologies as indicators of contamination is that no exposure experiment is required because
483 samples are collected *in situ* and examined, which allows for sampling of multiple sites in a short period of
484 time. Moreover, analyses can be performed later if samples are properly stored and preserved.

485 The ecological status of the same 16 sites was also assessed based on a suite of diatom
486 descriptors (cell size, frustule health, lipid bodies, frustules deformities, etc.) in previous studies (Pandey
487 and Bergey 2016; Pandey et al. 2018). Overall, the authors found that water quality assessment based on
488 diatom assemblages and diatom-based metrics had a good fit with the available physicochemical data (least
489 versus most impacted sites). However, as was observed in the present study, a greater number of sites
490 showed signs of degradation based on the diatom metrics used (biological descriptors) compared with the
491 available physicochemical data. This suggests that the use of biotic indicators provides useful
492 complementary information on ecosystem health status at the selected sites, and that *in situ* diatom
493 assemblages are good assessment tools for monitoring rivers, alone or in combination with bioassays.

494 **3.3. Bioassay scores and overall index values**

495 *3.3.1. Toxicity assessment of water samples: complementarity of the bioassays*

496 The scores obtained for the six bioassays after exposure to the 16 test water samples as well as for
497 the diatom teratology assessment are presented in Table 5, along with the overall index classes. The test
498 organisms used covered different trophic levels, from decomposers (*A. fischeri*) and primary producers (*P.*
499 *subcapitata*, *L. minor*, diatoms), to primary consumers (*H. incongruens* and *M. macrocopa*) and, ultimately,
500 secondary consumers (*D. rerio*). The scores obtained for each test organism varied greatly, with certain
501 bioassays indicating high toxicity while others showed no response. The bacterial and microalgal bioassays
502 were the least sensitive because only one of the water samples was characterised as being of moderate
503 toxicity. The fish embryo test was the most sensitive to the types of contamination present in the test waters
504 because it showed a response in almost all samples. Different levels of responses to toxicity were observed
505 in the present study, with only ostracods, duckweed, and diatoms evidencing “high toxicity” for certain
506 samples. This heterogeneity in response sensitivity using multiple organisms was also noted in other

507 bioassay-based assessments (e.g., Persoone et al. 2003; Pandard et al. 2006; Mankiewicz-Boczek et al.
508 2008). Organisms from various trophic levels have also shown different responses to perturbations using
509 biological indices based on assemblage structure and other biological descriptors (e.g., Marzin et al. 2012;
510 Lainé et al. 2014). In fact, this combination of trophic levels allows for a better assessment of the overall
511 contamination of waters by substances with different modes of action as well as biological targets.
512 When contamination was high, such as at H1, most bioassays (five out of seven) highlighted potential
513 toxicity. This was also the case, to a lower extent, at N2 and H2. The advantages of this multi-bioassay
514 approach are more striking in the case of subtle water contamination. In the latter case, the type of
515 organisms affected, or the functions impaired, could provide valuable information to identify the potential
516 nature of contamination. For example, despite the low sensitivity of the *P. subcapitata* bioassay to water
517 toxicity, microalgal growth was significantly enhanced at 10 sites, where higher nutrient loads were found.
518 Analysing this response in light of the response of the other test organisms, two scenarios can be
519 hypothesised. First, growth could have been stimulated by higher nutrient availability at sites with low
520 concentrations of toxicants. This scenario may be valid at sites Y2, G2, G3, G5, G7, and G6, where no
521 toxicity was observed on other plant organisms tested (macrophytes and diatoms). In contrast, and given
522 the high sensitivity of the root re-growth assay to metal contamination (Park et al. 2013), and of diatom
523 teratologies (Morin et al. 2012 Lavoie et al. 2017), a second scenario likely occurred at Y1, Y3, Y4, G1,
524 and G4: the stimulating effects of nutrients on microalgae may have masked the potential toxicity of zinc
525 present in the test waters. Microalgal growth stimulation by nutrients in metal- (Cd/Zn) contaminated sites
526 was also observed in periphytic biofilms (Morin et al. 2008b). The first scenario, however, only discards
527 the presence of plant-targeting substances; indeed, some toxicity was observed in the growth of freshly
528 hatched ostracods and/or in fish embryo survival at G2, G3, G5, and G7. This suggests contamination by
529 compounds specifically affecting the first stages of animal cell development. In contrast, at N2 and H1,
530 moderate to high toxicity was found through diverse bioassays, covering different trophic levels; this may
531 be the consequence of contamination by narcotic toxicants (i.e., non-specific acting) or by a mixture of
532 dissimilarly acting compounds. The results from the present study underscore the significant benefit of
533 using a multi-organism approach allowing for a better integration of water quality testing, which in turn
534 results in a more complete assessment of complex environmental stresses. Moreover, this approach is of

535 particular significance in the goal to characterise the overall water toxicity of the freshwater ecosystem.
536 Organisms have the major advantage of reflecting the toxic potential of waters, which cannot be
537 sufficiently highlighted by chemical measurements alone due to the limitations stated below.

538 *3.3.2. Toxicity scores versus water chemical analyses*

539 Based solely on available water chemistry data, N2, N3, H1, and H2 are the sampling sites
540 suggesting more severely degraded conditions, mostly due to high metal concentrations, total nitrogen,
541 coliforms, BOD, and conductivity. No toxic contamination (metals or organic compounds) was found at the
542 other sampling sites based on water chemistry (Pandey et al. 2018). The results from the present multi-
543 bioassay study identified H1 as “highly toxic” to living organisms (class D). N2, H2, Y1, and Y3 were
544 classified into the “moderate toxicity” class (C), and N3 into the “slight toxicity” class (B). Some sites,
545 although exhibiting some impact (e.g., two assays indicating some toxicity at Y4, G2, and G3), fell into the
546 “no toxicity” category. All samples from the Geum River, as well as N1, Y2, and Y4, were considered not
547 toxic to aquatic life. The fact that additional sampling sites suggested some toxicity when using biological
548 descriptors compared to chemistry alone underscores that (i) chemical analyses cannot be exhaustive (some
549 compounds with toxicity may not have been targeted), (ii) quantification limits may be higher than the
550 toxic concentrations, and (iii) mixture effects are not considered. In fact, toxicity often results from the
551 effects of a cocktail of compounds and their degradation products (Kim Tiam et al. 2016), which renders
552 the task of water quality assessment even more challenging due to additive, synergistic, and antagonistic
553 interactions.

554 *3.3.3. Scoring approach: preliminary considerations*

555 Classification based on the “harmfulness potential” of natural waters is relatively uncommon as
556 bioassays are generally used to derive EC_{50} for particular chemicals or for wastewaters. The most similar
557 approach found for comparison is the scoring system presented by Persoone et al. (2003) and also used by
558 Mankiewicz-Boczek et al. (2008), where the scores obtained from the bioassays are expressed as an overall
559 degree of hazard or toxicity. Although the toxicity classes differ as well as the overall method for indexing
560 hazardous potential of the sample, the approach proposed in the present study is generally comparable to
561 that of Persoone et al. (2003). As mentioned in the methods, the criteria used to establish the toxicity
562 categories and scores might be subjected to change in the future as the use of this approach becomes more

563 popular and experience is gained. For example, the criteria were set using similar boundaries between
564 biotests for simplicity, but this may be refined as similar studies multiply. The final overall toxicity classes
565 may also have to be refined in the future, but as of now, this classification system seems to be appropriate
566 to adequately qualify the tested waters. This type of investigation is still in its infancy and it is presently
567 difficult to identify the most sensitive and reliable test organisms. As an example, bioluminescence does
568 not seem to be an appropriate bioassay based on the results from this study. However, this does not
569 necessarily mean that bacteria-based bioassays are inefficient. The potential toxic effects of natural waters
570 on bioluminescence need to be tested using a larger array of water samples covering a range of different
571 types of contamination. This statement is also valid for the other organisms tested. With experience, it
572 should become easier to select the suite of organisms that is most appropriate for the type of water tested,
573 based on the nature of the suspected contamination (when this applies). The main objective of this
574 preliminary investigation was to lay the foundation for this biomonitoring approach in South Korean rivers,
575 and the results are promising in the way that the test organisms used complemented each other and
576 supplemented traditional chemistry measurements.

577 **Conclusion**

578 South Korean rivers, as with many rivers worldwide, receive a great deal of various chemicals
579 from urban, industrial, and agricultural activities, which makes the measurement of all substances
580 practically impossible (e.g., Vörösmarty et al. 2010). Moreover, new chemical compounds are constantly
581 detected in surface waters, rendering the task of water toxicity assessment even more challenging and
582 costly. The toxicity index proposed in this study is a valuable tool for preliminary screening of water
583 contamination as an alternative or in addition to traditional chemistry-based assessments. Based on the 16
584 water samples collected, the various bioassays tested provided complementary information to chemical
585 analyses by flagging additional sites as being potentially degraded. The variability in the response of each
586 organism to water exposure underscores the need for testing toxicity based on a multi-organism approach,
587 which can possibly highlight the kind of toxic substance that is most responsible for water degradation. As
588 it would be utopic to recommend using all organisms in routine monitoring, suggesting to at least test water
589 samples using representative organisms from different trophic levels seems appropriate at this time.

590

591 **Acknowledgement**

592 This work was partly supported by Post-Doctor Research Program (2016) through Incheon National
593 University and Industrial Strategic Technology Development Program (Grant No. 10079956) funded by the
594 Ministry of Trade, Industry & Energy. We are grateful to Dr J.C. Taylor (North-West University, South
595 Africa) for his generous donation of Pleurax and to Emilie Saulnier-Talbot for valuable comments on the
596 manuscript and for English revision.

597
598 **References**

- 599
600 Ahlf, W., Hollert, H., Neumann-Hensel, H., & Ricking, M. (2002). A guidance for the assessment and
601 evaluation of sediment quality: a German approach based on ecotoxicological and chemical
602 measurements. *Journal of Soils and Sediments*, 2, 37–42.
- 603 Angerville, R. (2009). Evaluation des risques écotoxicologiques liés au déversement de rejets urbains par
604 temps de pluie (RUTP) dans les cours d'eau : Application à une ville française et à une ville
605 haïtienne. Chemical Sciences. INSA de Lyon, in French.
- 606 Angerville, R., Perrodin, Y., Bazin, C., & Emmanuel, E. (2013). Evaluation of ecotoxicological risks
607 related to the discharge of combined sewer overflows (CSOs) in a periurban river. *International
608 Journal of Environmental Research and Public Health*, 10, 2670-2687.
- 609 Ankley, G.T., Schubauer-Berigan, M.K., & Hoke, R.A. (1992). Use of toxicity identification evaluation
610 techniques to identify dredged material disposal options: a proposed approach. *Environmental
611 Management*, 16, 1–6.
- 612 Arini, A., Feurtet-Mazel, A., Maury-Brachet, R., Pokrovsky, O. S., Coste, M., & Delmas, F. (2012).
613 Recovery potential of periphytic biofilms translocated in artificial streams after industrial
614 contamination (Cd and Zn). *Ecotoxicology*, 21, 1403-1414.
- 615 Backhaus, T., Froehner, K., Altenburger, R., & Grimme, L. H. (1997). Toxicity testing with *Vibrio fischeri*:
616 A comparison between the long term (24 h) and the short term (30 min) bioassay. *Chemosphere*,
617 35, 2925-2938.
- 618 Becouze-Lareure, C., Thiebaud, L., Bazin, C., Namour, P., Breil, P., & Perrodin, Y. (2016). Dynamics of
619 toxicity within different compartments of a peri-urban river subject to combined sewer overflow
620 discharges. *Science of the Total Environment*, 539, 503–514.
- 621 Bresch, H. (1991). Early life-stage in zebrafish versus a growth test in rainbow trout to evaluate toxic
622 effects. *Bulletin of Environmental Contamination and Toxicology*, 46, 641–648.
- 623 Calabrese, E.J. (2005). Paradigm lost, paradigm found: The re-emergence of hormesis as a fundamental
624 dose response model in the toxicological sciences. *Environ. Pollut*, 138, 378-411.
- 625 Canesi, L., & Corsi, I. (2016). Effects of nanomaterials on marine invertebrates. *Science of The Total
626 Environment*, 565, 933-940.
- 627 Cerisier, A., Vedrenne, J., Lavoie, I., & Morin, S. (2018). Assessing the severity of diatom deformities
628 using geometric morphometry. *Botany Letters*, 1-9.
- 629 Chial, Z. B., & Persoone, G. (2002a). Cyst-based toxicity tests XII - Development of a short chronic
630 sediment toxicity test with the ostracod crustacean *Heterocypris incongruens*: selection of test
631 parameters. *Environmental Toxicology*, 17, 520-527.
- 632 Chial, Z. B., & Persoone, G. (2002b). Cyst-based toxicity tests XIV - Application of the ostracod solid-
633 phase microbiotest for toxicity monitoring of river sediments in Flanders (Belgium).
634 *Environmental Toxicology*, 17, 533-537.
- 635 Cho, E., Khim, J., Chung, S., Seo, D., & Son, Y. (2014). Occurrence of micropollutants in four major rivers
636 in Korea. *Science of the Total Environment*, 491-492, 138-147.
- 637 Chu, K. W., & Chow, K. L. (2002). Synergistic toxicity of multiple heavy metals is revealed by a
638 biological assay using a nematode and its transgenic derivative. *Aquatic Toxicology*, 6, 53–64.

- 639 Coste, M., Boutry, S., Tison-Rosebery, J., & Delmas, F. (2009). Improvements of the Biological Diatom
640 Index (BDI): Description and efficiency of the new version (BDI-2006). *Ecological Indicators*, 9,
641 621-650.
- 642 Davis, J. A. (1981). Comparison of static-replacement and flow through bioassays using duckweed, *Lemna*
643 *gibba* G3.EPA 560/6-81-003.United States Environmental Protection Agency, Washington, DC,
644 USA.
- 645 Elendt, B.-P. (1990). Selenium deficiency in Crustacea; An ultrastructural approach to antennal damage in
646 *Daphnia magna* Straus. *Protoplasma*, 154, 25-33.
- 647 Environment Canada. (1992). Biological Test method: Growth Inhibition Test Using the Freshwater Alga
648 *Selenastrum capricornutum*. Report EPS 1/RM/25, Environment Canada, Ottawa, ON, Canada.
- 649 Fulladosa, E., Murat, J. C., & Villaescusa, I. (2005).Effect of Cadmium(II), Chromium(VI), and Arsenic(V)
650 on long-term viability and growth-inhibition assays using *Vibrio fischeri* marine bacteria. *Archives*
651 *of Environmental Contamination and Toxicology*, 49, 299–306.
- 652 Fulladosa, E., Murat, J. C., Bollinger, J. C., & Villaescusa, I. (2007). Adverse effects of organic arsenical
653 compounds towards *Vibrio fischeri* bacteria. *Science of the Total Environment*, 377, 207-213.
- 654 Geiszinger, A., Bonnineau, C., Faggiano, L., Guasch, H., López-Doval, J. C., Proia, L., Ricart, M.,
655 Ricciardi, F., Romani, A., Rotter, S., Muñoz, I., Schmitt-Jansen, M., & Sabater, S. (2009). The
656 relevance of the community approach linking chemical and biological analyses in pollution
657 assessment. *Trends in Analytical Chemistry*, 28, 619-626.
- 658 Gonzalez-Merchan, C., Perrodin, Y., Barraud, S., Sébastien, C., Becouze-Lareure, C., & Bazin, C. (2014a).
659 Spatial variability of sediment ecotoxicity into a large storm water detention basin. *Environmental*
660 *Science and Pollution Research*, 21, 5357–5366.
- 661 Gonzalez-Merchan, C., Perrodin, Y., Sébastien, C., Bazin, C., Winiarski, T., & Barraud, S. (2014b).
662 Ecotoxicological characterization of sediments from storm water retention basins. *Water Science*
663 *& Technology*, 69, 1045–1051.
- 664 Gopalapillai, Y., Vigneault, B., & Hale, B. A. (2014). Root length of aquatic plant, *Lemna minor* L., as an
665 optimal toxicity endpoint for biomonitoring of mining effluents. *Integrated Environmental*
666 *Assessment and Management*, 10, 493-497.
- 667 Hassan, I., Jabir, N.R., Ahmad, S., Shah, A., & Tabrez, S. (2015). Certain phase I and phase II enzymes as
668 toxicity biomarker: an overview. *Water Air Soil Pollut.* 226, 153.
- 669 Hsieh, C-Y., Tsai, M-H., Ryan, D. K., & Pancorbo, O. C. (2004). Toxicity of the 13 priority pollutant
670 metals to *Vibrio fischeri* in the Microtox chronic toxicity test. *Science of the Total Environment*,
671 320, 37–50.
- 672 Hwang, S-J., Lee, S-W., & Park, Y-S. (2011).Ecological monitoring, assessment, and restoration of
673 running waters in Korea. *International Journal of Limnology: Annales de Limnologie*, 47, S1-S2.
- 674 ISO 11348-1:2009. Water quality — Determination of the inhibitory effect of water samples on the light
675 emission of *Vibrio fischeri* (Luminescent bacteria test) – Part1: Method using freshly prepared
676 bacteria.
- 677 ISO 14371:2012. Water quality- determination of fresh watersediment toxicity to *Heterocypris incongruens*
678 (Crustacea, Ostracoda).
- 679 Ji, K., Kim, Y., Oh, S., Ahn, B., Jo, H., & Choi, K. (2008). Toxicity of Perfluorooctanoic acid on
680 freshwater macro invertebrates (*Daphnia magna* and *Moina macrocopa*) and fish (*Oryzias latipes*).
681 *Environ. Toxicol. Chem.* 27, 2159–2168.
- 682 Joy, M. K., & Death, R. G. (2002). Predictive modelling of freshwater fish as a biomonitoring tool in New
683 Zealand. *Freshwat. Biol.* 47, 2261–2275.
- 684 Katsumata, M., Koike, T., Nishikawa, M., Kazumura, K., & Tsuchiya, H. (2006). Rapid ecotoxicological
685 bioassay using delayed fluorescence in the green alga *Pseudokirchneriella subcapitata*. *Water*
686 *Research*, 40, 3393–3400.
- 687 Kelly, M. (2013). Building capacity for ecological assessment using diatoms in UK rivers. *Journal of*
688 *Ecology and Environment*, 36, 89–94.
- 689 Kim Tiam, S., Fauvelle, V., Morin, S., & Mazzella, N. (2016). Improving toxicity assessment of pesticide
690 mixtures: The use of polar passive sampling devices extracts in microalgae toxicity tests. *Frontiers*
691 *in Microbiology*, 7, 1388.
- 692 Kim, S. B., Kim, W. K., Chounlamany, V., Seo, J., Yoo, J., Jo, H. J., & Jung, J. (2012). Identification of
693 multi-level toxicity of liquid crystal display wastewater toward *Daphnia magna* and *Moina*
694 *macrocopa*. *Journal of Hazardous Materials*, 227–228, 327–333.

- 695 Kim, Y.-J., Han, Y.-S., Kim, E., Jung, J., Kim, S.-H., Yoo, S.-J., Shin, G.-S., Oh, J.-J., Park, A., Choi, H., Kim,
696 M.-S., Brown, M. T., & Han, T. (2015). Application of the *Ulva pertusa* bioassay for a toxicity
697 identification evaluation and reduction of effluent from a wastewater treatment plant. *Frontiers in*
698 *Environmental Science*, 3, 1-9.
- 699 Lainé, M., Morin, S., & Tison-Rosebery, J. (2014). A multicompartiment approach - diatoms, macrophytes,
700 benthic macro-invertebrates and fish-to assess the impact of toxic industrial releases on a small
701 French river. *PLoS ONE*, 9, e102358..
- 702 Lavoie, I., Campeau, S., Zugic-Drakulic, N., Winter, J.G., & Fortin, C. (2014). Using diatoms to monitor
703 stream biological integrity in Eastern Canada: An overview of 10years of index development and
704 ongoing challenges. *Science of the Total Environment*, 475, 187-200.
- 705 Lavoie, I., Hamilton, P., Morin, S., Kim Tiam, S., Gonçalves, S., Falasco, E., Fortin, C., Gontero, B.,
706 Heudre, D., Kahlert, M., Kojadinovic-Sirinelli, M., Manoylov, K., Pandey, L. K., & Taylor, J.
707 (2017). Diatom teratologies as biomarkers of contamination: are all deformities ecologically
708 meaningful? *Ecological Indicators*, 82, 539-550.
- 709 Lavoie, I., Lavoie, M., & Fortin, C. (2012). A mine of information: benthic algal communities as
710 biomonitors of metal contamination from abandoned tailings. *Science of the Total Environment*,
711 425, 231-241.
- 712 Lento, J., Dillon, P. J., Somers, K. M., & Reid, R. A. (2008). Changes in littoral benthic macroinvertebrate
713 communities in relation to water chemistry in 17 Precambrian Shield lakes. *Canadian Journal of*
714 *Fisheries and Aquatic Sciences*, 65, 906-918.
- 715 Li, L., Zheng, B., & Liu, L. (2010). Biomonitoring and bioindicators used for river ecosystems: definitions,
716 approaches and trends. *Procedia Environmental Sciences*, 2, 1510–1524.
- 717 Macken, A., Giltrap, M., Ryall, K., Foley, B., McGovern, E., McHugh, B., & Davoren, M. (2009). A test
718 battery approach to the ecotoxicological evaluation of cadmium and copper employing a battery of
719 marine bioassays. *Ecotoxicology*, 18, 470–480.
- 720 Mankiewicz-Boczek, J., Nałecz-Jawecki, G., Drobnińska, A., Kaza, M., Sumorok, B., Izydorczyk, K.,
721 Zalewski, M., & Sawicki, J. (2008). Application of a microbiotests battery for complete toxicity
722 assessment of rivers. *Ecotoxicology and Environmental Safety*, 71, 830–836.
- 723 M.O.E. Korea, (2007). Toxicity Identification of Ecological Risk Hazards in the Waste Waters around the
724 Industrial Complex from 2015, Ministry of Environment, Korea.
- 725 Marzin, A., Archaimbault, V., Belliard, J., Chauvin, C., Delmas, F., & Pont, D. (2012). Ecological
726 assessment of running waters: do macrophytes, macroinvertebrates, diatoms and fish show similar
727 responses to human pressures? *Ecological Indicators*, 23, 56-65.
- 728 Moreira-Santos, M., Soares, A. M. V. M., & Ribeiro, R. (2004). An in situ bioassay for freshwater
729 environments with the microalga *Pseudokirchneriella subcapitata*. *Ecotoxicology and*
730 *Environmental Safety*, 59, 164–173.
- 731 Morin, S., Duong, T. T., Dabrin, A., Coynel, A., Herlory, O., Baudrimont, M., Delmas, F., Durrieu, G.,
732 Schäfer, J., Winterton, P., Blanc, G., & Coste, M. (2008a). Long-term survey of heavy-metal
733 pollution, biofilm contamination and diatom community structure in the Riou Mort watershed,
734 South-West France. *Environmental Pollution*, 151, 532-542.
- 735 Morin, S., Duong, T.T., Boutry, S., & Coste, M. (2008b). Mitigation of metal toxicity to freshwater
736 biofilms development (Decazeville watershed, SW France). *Cryptogamie Algologie*, 29, 201-216.
- 737 Morin, S., Cordonier, A., Lavoie, I., Arini, A., Blanco, S., Duong, T. T., Tornés, E., Bonet, B., Corcoll, N.,
738 Faggiano, L., Laviale, M., Pérès, F., Becares, E., Coste, M., Feurtet-Mazel, A., Fortin, C., Guasch,
739 H., & Sabater, S. (2012). Consistency in diatom response to metal-contaminated environments. In:
740 Guasch, H., Ginebreda, A., Geiszinger, A. (eds.) *Emerging and Priority Pollutants in Rivers*.
741 Springer-Verlag, Berlin, pp. 117-146.
- 742 Nedeau, E. J., Merritt, R. W., & Kaufman, M. G. (2003). The effect of an industrial effluent on an urban
743 stream benthic community: water quality vs. habitat quality. *Environmental Pollution*, 123, 1–13.
- 744 Nichols, H. W. (1973) Growth media. Freshwater. In: Stein JR (ed.), *Handbook of Phycological Methods*.
745 *Culture Methods and Growth measurements*. Cambridge University Press, London, 7–24.
- 746 Niemirycz, E., Nichthauser, J., Staniszevska, M., Nałecz-Jawecki, G., & Bolałek, J. (2007). The
747 Microtox® biological test: Application in toxicity evaluation of surface waters and sediments in
748 Poland. *Oceanological and Hydrobiological Studies*, 36, 151-163.

- 749 Ntengwe, F. W., & Maseka, K. K. (2006). The impact of effluents containing zinc and nickel metals on
750 stream and river waterbodies: the case of Chambishi and Mwambashi streams in Zambia. *Physics*
751 *and Chemistry of the Earth*, 31, 814–820.
- 752 Oberdorff, T., Pont, D., Hugueny, B., & Porcher, J. P. (2002). Development and validation of a fish-based
753 index (FBI) for the assessment of “river health” in France. *Freshwater Biology*, 47, 1720–1734.
- 754 OECD. (1984). Alga, growth inhibition test. OECD Guideline for Testing of Chemicals, Vol. 201, Paris.
- 755 OECD. (2012). *Daphnia magna* reproduction test. OECD Guideline for the Testing of Chemicals, Vol. 211,
756 Paris.
- 757 OECD. (2013). Fish embryo acute toxicity (FET) test. OECD Guidelines for the testing of chemicals, Vol.
758 236, Paris.
- 759 Panda, S. K., & Upadhyay, R. K. (2003). Salt stress injury induces oxidative alteration and antioxidative
760 defense in the roots of *Lemna minor*. *Biologia Plantarum*, 48, 249–253.
- 761 Pandard, P., Devillers, J., Charissou, A.-M., Poulsen, V., Jourdain, M.-J., Férard, J.-F., Grand, C., & Bispo,
762 A. (2006). Selecting a battery of bioassays for ecotoxicological characterization of wastes. *Science*
763 *of the Total Environment*, 363, 114–125.
- 764 Pandey, L. K., & Bergey, E. A. (2016). Exploring the status of motility, lipid bodies, deformities and size
765 reduction in periphytic diatom community from chronically metal (Cu, Zn) polluted waterbodies
766 as a biomonitoring tool. *Science of the Total Environment*, 550, 372–381.
- 767 Pandey, L. K., Kumar, D., Yadav, A., Rai, J., Gaur, J. P. (2014). Morphological abnormalities in periphytic
768 diatoms as a tool for biomonitoring of heavy metal pollution in a river. *Ecological Indicators*, 36,
769 272–279.
- 770 Pandey, L. K., Lavoie, I., Morin, S., Park, J., Jie, L., Choi, S., Lee, H., & Han, T. (2018). River water
771 quality assessment based on a multi-descriptor approach including chemistry, diatom assemblage
772 structure, and non-taxonomical diatom metrics. *Ecological Indicators*, 84, 140–151.
- 773 Park, A., Kim, Y., Choi, E., Brown, M. T., & Han, T. (2013). A novel bioassay using root re-growth in
774 *Lemna*. *Aquatic Toxicology*, 140–141, 415–424.
- 775 Park, J., Brown, M. T., Depuydt, S., Kim, J. K., Won, D.-S., & Han, T. (2017). Comparing the Acute
776 Sensitivity of Growth and Photosynthetic Endpoints in Three *Lemna* Species Exposed to Four
777 Herbicides. *Environmental Pollution*, 220, 818–827.
- 778 Parvez, S., Venkataraman, C., & Mukherji, S. (2006). A review on advantages of implementing
779 luminescence inhibition test (*Vibrio fischeri*) for acute toxicity prediction of chemicals.
780 *Environment International*, 32, 265–268.
- 781 Persoone, G., Marsalek, B., Blinova, I., Andrea Törökne, A., Zarina, D., Manusadzianas, L., Nalecz-
782 Jawecki, G., Tofan, L., Stepanova, N., Tothova, L., & Kolar, B. (2003). A practical and user-
783 friendly toxicity classification system with microbiotests for natural waters and wastewaters.
784 *Environmental Toxicology*, 18, 395–402.
- 785 Ponader, K. C., Charles, D. F., & Belton, T. J. (2007). Diatom-based TP and TN inference models and
786 indices for monitoring nutrient enrichment of New Jersey streams. *Ecological Indicators*, 7, 79–
787 93.
- 788 Radix, P., Léonard, M., Papantoniou, C., Roman, G., Saouter, E., Gallotti-Schmitt, S., Thiébaud, H., &
789 Vasseur, P. (2000). Comparison of four chronic toxicity tests using algae, bacteria, and
790 invertebrate assessed with sixteen chemicals. *Ecotoxicology and Environmental Safety*, 47, 186–
791 194.
- 792 Rodrigues, E. S., & Umbuzeiro G. A. (2011). Integrating toxicity testing in the wastewater management of
793 chemical storage terminals—a proposal based on a ten-year study. *Journal of Hazardous*
794 *Materials*, 186, 1909–1915.
- 795 Reynoldson, R. H., Norris, V. H., Resh, K. E., Day, D. M., & Rosenberg, T. B. (1997). The reference
796 condition: A comparison of multimetric and multivariate approaches to assess water-quality
797 impairment using benthic macroinvertebrates. *Journal of the North American Benthological*
798 *Society*, 16, 833–852.
- 799 Ruiz, F., Abad, M., Bodergat, A. M., Carbonel, P., Rodríguez-Lázaro, J., González-Regalado, M. L.,
800 Toscano, A., García, E. X., & Prenda, J. (2013). Freshwater ostracods as environmental tracers.
801 *International Journal of Environmental Science and Technology*, 10, 1115–1128.
- 802 Sevilla, J. B., Nakajima, F., & Kasuga, I. (2014). Comparison of aquatic and dietary exposure of heavy
803 metals Cd, Cu, and Zn to benthic ostracod *Heterocypris incongruens*. *Environmental Toxicology*
804 *and Chemistry*, 33, 1624–1630.

805 Shen, K., Shen, C., Lu, Y., Tang, X., Zhang, C., Chen, X., Shi, J., Lin, Q., & Chen, Y. (2009). Hormesis
806 response of marine and freshwater luminescent bacteria to metal exposure. *Biological Research*,
807 *42*, 183-187.

808 Şişman, T., İncekara, Ü., & Yıldız, Y. Ş. (2008). Determination of acute and early life stage toxicity of fat-
809 plant effluent using zebrafish (*Danio rerio*). *Environmental Toxicology*, *23*, 480-486.

810 Small, A. M., Adey, A. H., Lutz, S. M., Reese, E. G., & Roberts, D. L. (1996). A macrophyte-based rapid
811 biosurvey of stream water quality: Restoration at the watershed scale. *Restoration Ecology*, *4*,
812 124-145.

813 Steinberg, R. (1946). Mineral requirement of *Lemna minor*. *Plant Physiology*, *21*, 42-48

814 Stevenson, R. J., Pan, Y., & Vandam, H. (2010). Assessing environmental conditions in rivers and streams
815 with diatoms. In: Smol J. P., and Stoermer E. F., (2nd Eds.), *The diatoms: applications for the*
816 *environmental and earth sciences*. Cambridge university press, London, pp. 57-85.

817 Tabrez, S., Ahmad, M. (2012). Cytochrome P450 system as potential biomarkers of certain toxicants:
818 comparison between plant and animal models. *Environ Monit Assess.* *185*(4), 2977-2987.

819 Tarkpea, M., & Hansson, M. (1989). Comparison between two Microtox test procedures. *Ecotoxicology*
820 *and Environmental Safety*, *18*, 204-210.

821 Thiebaut, G., Guérol, F., & Muller, S. (2002). Are trophic and diversity indices based on macrophyte
822 communities pertinent tools to monitor water quality? *Water Research*, *36*, 3602-3610.

823 Toumi, H., Burga-Perez, K. F., & Ferard, J-F. (2015). Acute and chronic ecotoxicity of carbaryl with a
824 battery of aquatic bioassays. *Journal of Environmental Science and Health, Part B*, *51*, 57-62.

825 USEPA (United States Environmental Protection Agency). (1994). Short-term methods for estimating the
826 chronic toxicity of effluents and receiving waters to freshwater organisms, EPA 600/7-91-
827 002, Washington, DC, USA.

828 Vincze, K., Graf, K., Scheil, V., Köhler, H., & Triebkorn, R. (2014). Embryotoxic and proteotoxic effects
829 of water and sediment from the Neckar River (Southern Germany) to zebrafish (*Danio rerio*)
830 embryos. *Environmental Sciences Europe*, *26*, 1-13.

831 Vörösmarty, C. J., McIntyre, P. B., Gessner, M. O., Dudgeon, D., Prusevich, A., Green, P., Glidden, S.,
832 Bunn, S. E., Sullivan, C. A., Liermann, C. R., & Davies, P. M. (2010). Global threats to human
833 water security and river biodiversity. *Nature*, *467*, 555-561.

834 Watanabe, H., Nakajima, F., Kasuga, I., & Furumai, H. (2008). Toxicity characterization of urban river
835 sediments using bioassay with ostracod. *Journal of Japan Society on Water Environment*, *31*, 671-
836 676.

837 Watanabe, H., Nakajima, K., Kasuga, I., & Furumai, H. (2011). Toxicity evaluation of road dust in the
838 runoff process using a benthic ostracod *Heterocypris incongruens*. *Science of the Total*
839 *Environment*, *409*, 2366-2372.

840 Weyers, A., Sokull-Klütgen, B., Baraibar-Fentanes, J., & Vollmer, G. (2000). Acute toxicity data: a
841 comprehensive comparison of results of fish, Daphnia, and algae tests with new substances
842 notified in the European Union. *Environmental Toxicology and Chemistry*, *19*, 1931-1933.

843 Wolska, L., Sagajdakow, A., Kuczyńska, A., & Namieśnik, J. (2007). Application of ecotoxicological
844 studies in integrated environmental monitoring: possibilities and problems. *Trends in Analytical*
845 *Chemistry*, *26*, 332-344.

846 Yi, X., Kang, S. W., & Jung, J. (2010). Long-term evaluation of lethal and sublethal toxicity of industrial
847 effluents using *Daphnia magna* and *Moinamacrocopa*. *Journal of Hazardous Materials*, *178*, 982-
848 987.




Fig. 1. Site distribution along four major rivers of South Korea.




Fig. 2. Normal and deformed frustules of the diatoms *Ulnaria ulna* (a-c), *Diatoma vulgaris*, (d,e), *Pinnularia biceps* (f) and *Gomphonema parvulum* (g) examined from the 16 sites. 'N' denotes normal frustules.

Table 1. Toxicity classes and attributed scores for each biological descriptor and endpoints.

Bioassay/biological descriptor	Toxicity class and score		
	No toxicity (%)	Moderate toxicity (%)	High toxicity (%)
	Score = 0	Score = 1	Score = 2
% inhibition in bioluminescence of <i>Aliivibrio fischeri</i>	[0-20]	[20-50]	>50
% growth inhibition in <i>Pseudokirchneriella subcapitata</i>	[0-20]	[20-50]	>50
% mortality in <i>Heterocypris incongruens</i>	[0-20]	[20-50]	>50
% growth length inhibition in <i>Heterocypris incongruens</i>	[0-20]	[20-50]	>50
% mortality in <i>Moina macrocopa</i>	[0-20]	[20-50]	>50
% growth length inhibition in <i>Moina macrocopa</i>	[0-20]	[20-50]	>50
% inhibition in reproduction of <i>Moina macrocopa</i>	[0-20]	[20-50]	>50
% mortality in <i>Danio rerio</i> embryos	[0-10]	[10-50]	>50
% inhibition in root elongation of <i>Lemna minor</i>	[0-20]	[20-50]	>50
% deformities in diatoms	<0.5	0.5-1	>1

Table 2. Proposed cross-tabulation approach for the calculation of an overall toxicity category (index) based on (1) the total score obtained by adding individual scores from each test (column), and (2) the number of tests that received a score = 2 suggesting “high toxicity” (rows). Because the number of tests performed may change depending on the project, the values used for this tabulation are reported as percentages.

		Total score (%)				
		[0-20]	[20-40]	[40-60]	[60-80]	[80-100]
% of tests with "high toxicity" score	0					
	[0-20]					
	[20-40]					
	[40-60]					
	[60-80]					

Overall toxicity category (index)
A = no toxicity
B = slightly toxic
C = moderately toxic
D = toxic
E = very toxic

Table 3. Physico-chemical measurements averaged (\pm SD) per river (Nakdong, 3 sites; Yeongsan, 4 sites; Geum, 7 sites; Han, 2 sites).

Parameter	Nakdong	Yeongsan	Geum	Han
pH	8.2 \pm 0.4	8.2 \pm 0.4	8.0 \pm 0.2	8.0 \pm 0.4
Temperature ($^{\circ}$ C)	25.7 \pm 1.4	22.2 \pm 0.7	22.8 \pm 0.4	26.2 \pm 2.1
Conductivity (μ S/cm)	1982 \pm 932	251 \pm 44	458 \pm 45	3293 \pm 1010
Dissolved Oxygen (mg/L)	11.5 \pm 0.0	11.6 \pm 0.2	11.4 \pm 0.1	10.6 \pm 0.4
Total Nitrogen (mg/L)	9.0 \pm 1.9	3.9 \pm 0.8	7.6 \pm 1.6	12.0 \pm 3.0
Total Phosphorus (mg/L)	0.07 \pm 0.03	0.09 \pm 0.03	0.21 \pm 0.04	0.10 \pm 0.00
TDS (mg/L)	1421 \pm 660	223 \pm 25	284 \pm 51	1515 \pm 873
BOD ₅ (mg/L)	1.8 \pm 0.2	1.7 \pm 0.2	1.8 \pm 0.4	6.3 \pm 4.1
Coliforms (CFU/mL)	7 \pm 3	26 \pm 18	16 \pm 6	152 \pm 124
Cu (μ g/L)	39 \pm 13*	nd	nd†	43‡
Ni (μ g/L)	57 \pm 16*	nd	nd	19 \pm 4
Zn (μ g/L)	42 \pm 21	66 \pm 6	71 \pm 6	370 \pm 99

*: Not detected in N1

†: Except in G7: 4 μ g/L

‡: Not detected in H1

Table 4. Results of the six bioassays and the diatom teratology assessment. Inhibition is calculated in comparison to the controls (positive values: inhibition, negative values: stimulation), while mortality rates (number of dead individuals scaled to the initial size of the population) and diatom teratologies are given as raw percentages. Average response (Avg), confidence intervals (CI, 95%), significant differences with the controls (p-values after post-hoc Tukey test; * <0.05, ** <0.01, *** <0.001) and ANOVA outputs are given for the tests that were replicated. Bold indicates toxicity. See tables S1 and S2 for complete information about sites.

Rivers	Bioassay	Bacteria	Microalgae	Ostracods		Cladocera	Fish	Macrophytes	Diatoms
	Sites	Inhibition of bioluminescence	Inhibition of average growth rate	Mortality	Growth inhibition	Inhibition of reproduction	Embryo mortality	Inhibition of root re-growth	Teratologies
		Avg (CI)	Avg (CI)	Avg (CI)	Avg (CI)	Avg (CI)		Avg (CI)	Avg (CI)
	Control	1 (2)	0 (1)	35 (17)	0 (15)	0 (7)	5	0 (22)	0.3 (0.1)
Nakdong	N1	9 (18)	11 (5)**	32 (13)	-31 (23)	3 (7)	5	8 (29)	0.8 (0.7)
	N2	28 (3)***	-7 (3)	27 (20)	-15 (33)	6 (13)	15	40 (9)***	1.5 (0.2)***
	N3	9 (4)	-1 (3)	37 (17)	22 (24)	-31 (7)	10	24 (24)*	0.2 (0.2)
Yeongsan	Y1	0 (0)	-15 (12)***	42 (20)	52 (19)**	-11 (5)	10	2 (14)	2.1 (0.4)***
	Y2	-6 (1)	-28 (5)***	43 (18)	18 (26)	-28 (9)	10	11 (14)	0.2 (0.0)
	Y3	13 (6)	-13 (10)	45 (15)	20 (22)	1 (4)	10	9 (11)	3.1 (0.7)***
	Y4	13 (13)	-32 (1)***	30 (16)	0 (22)	44 (8)**	5	49 (12)***	0.2 (0.0)
Geum	G1	-2 (3)	-21 (4)***	20 (13)	-11 (11)	-60 (6)***	5	41 (7)***	0.2 (0.0)
	G2	16 (3)*	-26 (0)***	40 (18)	44 (19)*	-25 (8)	15	8 (10)	0.4 (0.0)
	G3	-6 (21)	-17 (8)***	56 (24)	45 (37)*	-36 (5)*	40	-1 (20)	0.1 (0.0)
	G4	-11 (4)	-37 (2)***	25 (13)	-13 (17)	-28 (7)	5	-5 (19)	0.3 (0.0)
	G5	-5 (12)	-23 (6)***	10 (7)*	-35 (13)	-20 (6)	25	10 (3)	0.3 (0.0)
	G6	-6 (8)	-9 (8)	5 (7)**	-37 (5)	-33 (7)*	0	-1 (16)	0.4 (0.1)

	G7	2 (9)	-34 (3)***	23 (12)	4 (20)	-44 (9)**	10	17 (6)	0.5 (0.1)
Han	H1	11 (24)	30 (8)***	100 (0)***	/	-77 (5)***	20	52 (11)***	4.9 (0.6)***
	H2	-19 (15)*	7 (3)	15 (15)	-33 (18)	-22 (10)	10	62 (10)***	5.9 (0.9)***
ANOVA		F=4.18, p<0.0001	F=37.38, p<0.0001	F=6.02, p<0.0001	F=7.21, p<0.0001	F=9.10, p<0.0001	/	F=7.20, p<0.0001	F=85.58, p<0.0001

Table 5. Scores obtained for the six bioassays and the diatom teratology assessment after exposure to the 16 water samples tested, and final toxicity class (index). See table 2 for more detail on overall toxicity classes and tables S1 and S2 for complete information about sites.

Rivers	Bioassay	Bacteria	Microalgae	Ostracods	Cladocera	Fish	Macrophytes	Diatoms	Total score	Number of tests with score = 2	Overall toxicity class
	Sites	Bioluminescence	Growth	Mortality	Mortality	Embryo mortality	Root regrowth	Teratologies			
Nakdong	Ayang (N1)	0	0	0	0	0	0	1	1 (7%)	0 (0%)	A
	Dalseo(N2)	1	0	0	0	1	1	2	5 (36%)	1 (14%)	C
	Keumho(N3)	0	0	0	0	1	1	1	3 (21%)	0 (0%)	B
Yeongsan	Gw.gong(Y1)	0	0	2	0	1	0	2	5 (36%)	2 (29%)	C
	Gwangju-1(Y2)	0	0	0	0	1	0	0	1 (7%)	0 (0%)	A
	GJ-2(Y3)	0	0	0	0	1	0	2	3 (21%)	1 (14%)	C
	PD-1(Y4)	0	0	0	1	0	1	0	2 (14%)	0 (0%)	A
Geum	Ohryang(G1)	0	0	0	0	0	1	0	1 (7%)	0 (0%)	A
	Bangchuk(G2)	0	0	1	0	1	0	0	2 (14%)	0 (0%)	A
	Masan(G3)	0	0	1	0	1	0	0	2 (14%)	0 (0%)	A
	Sucheol(G4)	0	0	0	0	0	0	0	0 (0%)	0 (0%)	A
	Miho-5(G5)	0	0	0	0	1	0	0	1 (7%)	0 (0%)	A
	Miho-8(G6)	0	0	0	0	0	0	0	0 (0%)	0 (0%)	A
	Miho-7(G7)	0	0	0	0	1	0	0	1 (7%)	0 (0%)	A
Han	Soyo(H1)	0	1	2	0	1	2	2	8 (57%)	3 (43%)	D
	Daejeon(H2)	0	0	0	0	1	2	2	5 (36%)	2 (29%)	C