

Running title: Land use effects on stream diversity

**LOCAL AND REGIONAL DRIVERS OF TAXONOMIC HOMOGENIZATION IN STREAM COMMUNITIES
ALONG A LAND USE GRADIENT**

William R. Budnick¹, Thibault Leboucher², Jerome Beilard³, Janne Soininen⁴, Isabelle Lavoie⁵,
Katrina Pound¹, Aurélien Jamoneau², Juliette Tison-Rosebery², Evelyne Tales³, Virpi Pajunen⁴,
Stephane Campeau⁶, and Sophia I. Passy^{1*}

¹Department of Biology, University of Texas at Arlington, Arlington, Texas 76019, USA

²National Research Institute of Science and Technology for Environment and Agriculture
(Irstea), Aquatic Ecosystems and Global Changes Research Unit, Bordeaux, France

³Irstea, Hydrosystems under Change (HYCAR) Research Unit, Antony, France

⁴Department of Geosciences and Geography, University of Helsinki, Helsinki, Finland

⁵Institut national de la recherche scientifique, centre Eau Terre Environnement, 490 rue de la
Couronne, Québec City, Québec G1K 9A9, Canada

⁶Université du Québec à Trois-Rivières, 3351 boul. des Forges, Trois-Rivières, Québec G9A
5H7, Canada

*Corresponding author: sophia.passy@uta.edu

ACKNOWLEDGEMENTS

We thank W. Xu for helpful comments on the null model. Support by the National Science Foundation (grant NSF DEB-1745348) to SIP is gratefully acknowledged.

Author Biosketch: William R. Budnick is a Ph.D. candidate studying macroecology at the University of Texas at Arlington. His primary research interests lie with the application of cutting-edge numerical and spatial techniques to study and predict how global change forces will affect macroecological patterns particular to aquatic ecosystems. His interests also include integrating basic ecological theory with fisheries science to assist with the global conservation of threatened fish and crayfish fauna.